

Ecological site R042BB004NM

Gyp Interdune (Wet), Desert Shrub

Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Ecological site concept

This Ecological Site (ES) occurs exclusively on Whitesands National Monument and Whitesands Missile Range properties which are located within the Tulerosa basin. ES characteristics are found in the interdunes of barchnoid dunes. Where interdune has a watertable that fluctuates seasonally from surface to a few feet from soil surface. Facultative wet herbaceous vegetation occurs on this site. As one travels in the direction of the prevailing wind (SW to NE) along the Dunefield, this wet interdune ES is situated predominately prior to deflated barchnoid dune and barial of the interdune.

Associated sites

R042BB002NM	Gyp Duneland Barren, Desert Shrub The Gyp Interdune (wet) ecological site is associated with the Gyp Duneland Barren ecological site. They occur in association and can change places spatially as the dunes migrate downwind across the interdune.
-------------	---

Table 1. Dominant plant species

Tree	(1) <i>Populus fremontii</i>
Shrub	(1) <i>Yucca elata</i>
Herbaceous	(1) <i>Sporobolus airoides</i>

Physiographic features

This site occurs as the interdunal areas within the barchanoid type dunefields. During years with unusually high precipitation these interdunes may pond water for varying periods of time. Slope of the interdune landform averages 0 to 3 percent. Elevation ranges from approximately 3900 to 4200 feet above sea level.

Table 2. Representative physiographic features

Landforms	(1) Interdune
Flooding frequency	None
Ponding duration	Very brief (4 to 48 hours) to long (7 to 30 days)
Ponding frequency	Rare to occasional
Elevation	1,189–1,280 m

Slope	0–3%
Water table depth	0–102 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual average precipitation ranges from 7 to 12 inches. Wide fluctuations from year to year are common. At least one-half of the annual precipitation comes in the form of rainfall during July, August, and September. Precipitation in the form of snow or sleet averages less than 4 inches annually. The average annual air temperature is about 60 degree F. Summer maximums can exceed 100 degrees F. and winter minimums can go below zero. The average frost-free season exceeds 200 days and extends from April 1 to November 1. Both the temperature regime and rainfall distribution favor warm-season perennial plants on this site. Spring moisture conditions are only occasionally adequate to cause significant growth during this period of year. High winds from the west and southwest are common from March to June, which further tends to create poor soil moisture conditions in the springtime

Climate data was obtained from

<http://www.wrcc.dri.edu/summary/climsmnm.html>

Table 3. Representative climatic features

Frost-free period (average)	205 days
Freeze-free period (average)	227 days
Precipitation total (average)	305 mm

Influencing water features

This site is not influenced by water from a wetland or stream.

The interdunes have a water table that averages 30 to 40 inches in depth. During years that experience above average precipitation, water may pond for periods of several days to several weeks.

Soil features

The surface and underlying soils of this site are white gypsiferous sands. Depth to the water table averages 30 to 40 inches, but fluctuates seasonally. The soils are strongly saline and exhibit very rapid permeability.

Characteristic Soils:

NM-688 MU-426 Transformer-Duneland Association, 0-90 percent slopes

NM-688 MU-427 Duneland-Transformer Association, 0-90 percent slopes

Table 4. Representative soil features

Parent material	(1) Eolian sands–rock gypsum (2) Eolian deposits–rock gypsum
Surface texture	(1) Gypsiferous sand
Family particle size	(1) Sandy
Drainage class	Moderately well drained to somewhat poorly drained
Permeability class	Very rapid
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%

Available water capacity (0-101.6cm)	7.11–7.62 cm
Electrical conductivity (0-101.6cm)	15–24 mmhos/cm
Soil reaction (1:1 water) (0-101.6cm)	8–8.5
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics


Overview:

Plant community composition, distribution, and production within the interdunes are controlled by numerous factors. Some of which include depth to ground water, quality and amount of water, and the erosional and depositional effects of wind.

Changes from a grass/shrub state to a barren state may be initiated by ponding, an increase in amount of salts present on the site and or changes in wind flow patterns. Salts can accumulate on the surface of the Interdunes by the evaporation of shallow saline groundwater brought near the surface via capillary rise. Wind erosion can inhibit seedling establishment and can create interdune surfaces with features that enhance capillary rise. The grassland state of this ecological site is susceptible to invasion by saltcedar. Domination of the interdune by saltcedar is unlikely due to shallow depth to water. Saltcedar establishing on the interdunes can act as a seed source for the associated dunes, and adjoining areas.

State and transition model

R042XB004NM Gyp Interdune (Wet)



State 1 Grass / Shrub

This state is mix of midgrasses and various shrubs. This state tends to occur where the interdunes are least susceptible to ponding and increased salts. Plant densities also tend to be higher on the narrower interdunes and in the protected areas near the leeward side of the associated barchanoid dunes.

Community 1.1 Midgrass-Shrub community



Figure 4. Midgrass/Shrub



Figure 5. Midgrass/Shrub 2



Figure 6. Transformer Soil Pit

This community type is common on many of the interdunes within the barchanoid dunefields. Alkali sacaton is the dominant grass species. Other common grasses include Indian ricegrass, and New Mexico bluestem. Typical shrubs include ephedra, Trans-Pecos false claddaisy, yucca, and rabbitbrush.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	101	275	404
Shrub/Vine	22	123	196
Forb	39	62	106
Total	162	460	706

State 2

Exotic-Invaded State

The presence of saltcedar characterizes this state. Saltcedar densities are typically low, occurring as widely scattered individual plants.

Community 2.1

Midgrass/Shrub/Saltcedar



Figure 8. Saltcedar-Invaded

This community type is typically composed of alkali sacaton and various shrubs with saltcedar occurring as a minor component. Other grasses can include Indian ricegrass, NM bluestem, and sandhill muhly. Shrubs include yucca, ephedra, skunkbush sumac, frosted mint, and rabbitbrush.

State 3

Barren

This state is characterized by the lack of, or very sparse vegetation. Species if present usually consist of widely scattered alkali sacaton, and/or Iodinebush. These areas are subject to ponding following high intensity rainfall events.

Community 3.1

Sparse Sacaton/Iodinebush



Figure 9. Sparse Sacaton/Iodinebush

This community phase is typically composed of a sparse cover of alkali sacaton and or Iodinebush. Salinity levels probably help to limit species diversity within the community. This community phase often intergrades with the bare phase and may represent the topographic high spots within the interdunes.


Community 3.2
Bare



Figure 10. Bare

This community phase is characterized by the lack of vegetation.

Pathway 3.1A
Community 3.1 to 3.2



The community pathway to a bare or non vegetated community is believed to result from extensive erosion. This may be due in part to the orientation and size of the interdunes and associated dunes, and or actual dune movement. Additionally, extended periods of water ponding within the interdune may also cause this community pathway.

Pathway 3.2A
Community 3.2 to 3.1



Bare



Sparse Sacaton/Iodinebush

Seed dispersal from adjoining sites may provide a pathway from the bare community phase to a sparsely vegetated community. Deposition of gypsiferous sands across the interdune may favor seedling germination and survival.

Transition T1A
State 1 to 2

Introduction of seed is the primary factor in the cause of this transition. Due to the presence of a shallow water table, soils on this site usually contain adequate soil moisture necessary for seed germination. Ironically this shallow depth to water may naturally inhibit domination by saltcedar. Saltcedar seem to dominate on soils where depth to the water table is between 5 and 18 feet from the surface. Locations adjacent to areas with an abundant seed source or underground streams may also increase the likelihood of this transition.

Restoration pathway R2A
State 2 to 1

Brush management by chemical, mechanical or a combination can be used to remove saltcedar and return to the Grass/Shrub state. Manually removing and treating the cut stumps with “Garlon” is effective in the treatment of saltcedar. Additionally, spraying with “Arsenal” also has exhibited high kill rates. Regardless of the means of treatment, monitoring and follow-up will be necessary to insure missed plants and regrowth is addressed.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Warm season Mid/Tall grasses			90–364	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	90–280	–
	little bluestem	SCSCS	<i>Schizachyrium scoparium</i> var. <i>scoparium</i>	0–84	–
2	Cool season short grasses			11–39	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	11–39	–
Forb					
3	Forbs			39–106	
	greenthread	THELE	<i>Thelesperma</i>	17–45	–
	purple sand verbena	ABAN	<i>Abronia angustifolia</i>	17–39	–
	Forb, annual	2FA	<i>Forb, annual</i>	6–17	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	6–17	–
Shrub/Vine					
4	Shrubs			17–163	
	Trans-Pecos false clappaisy	PSAR	<i>Pseudocappia arenaria</i>	0–50	–
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	11–45	–
	rabbitbrush	CHRY9	<i>Chrysothamnus</i>	0–28	–
	James' seaheath	FRJA	<i>Frankenia jamesii</i>	0–11	–
5	Succulents			6–34	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	6–34	–
	soaptree yucca	YUEL	<i>Yucca elata</i>	6–34	–

Animal community

Characteristic wildlife includes little striped whiptail, White Sands prairie lizard, bleached earless lizard, spadefoot toad, Apache pocket mouse, White sands woodrat, spotted ground squirrel, camel cricket, tiger beetle, lycosid spider, and scorpions. Ord's kangaroo rat and spotted ground squirrel are also common to this site.

Hydrological functions

The dunefields and interdunes exhibit a relatively shallow ground water table. The water table depth varies seasonally and depending on precipitation and season ranges from the surface to a few feet deep. The shallow groundwater controls dune sediment accumulation. Additionally the moisture helps to stabilize the sediments resisting erosion and movement.

Recreational uses

This site and the associated Gypsum dune sites currently provide extensive recreational opportunities on the White Sands National Monument, and Guadalupe State Park. Where this site occurs on White Sands Missile Range access is restricted.

Wood products

This site provides no significant wood products.

Other products

Type locality

Location 1: Otero County, NM	
UTM zone	N
UTM northing	3640218
UTM easting	0381789
General legal description	Otero Co. 13.3 miles east and 1.6 miles north of junction of range roads 7&10

Other references

Other References:

Bennett, J., Wilder, D., 2009. Physical resources foundation report, White Sands National Monunent, Natural Resource Report NPS/NRPC/NRR—2009/166. National Park Service, Fort Collins, Colorado.

Di Tomaso, J. M., 1998. Impact, Biology, and Ecology of Saltcedar (*Tamarix* spp.) in the Southwestern United States. *Weed Technology*. 12: 326-336.

Fryberger, S.G., 2000. Geological overview of White Sands National Monument.
<http://www.nature.nps.gov/geology/parks/whsa/geows/index.htm>

Kocurek, G., Carr, M., Ewing, R., Havholm, K.G., Nagar, Y.C., Singhvi, a.K. 2007. White Sands Dune Field, New Mexico: Age, dune dynamics and recent accumulations. *Sedimentary Geology* 197 (2007) 313-331

Langford, R.P., Rose, J.M., White, D.E., 2009. Groundwater salinity as a control on development of eolian landscape: An example from White Sands of New Mexico. *Geomorphology* 105(2009) 39-49.

Muldavin, E., Chauvin, Y., Harper, G., 2000 The Vegetation of White Sands Missile Range, New Mexico. Volume 1: Handbook of Vegetation Communities. Final Report, New Mexico Natural Heritage Program. University of New Mexico. Albuquerque, New Mexico.

Contributors

David Trujillo

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-