

Ecological site R042CY755TX Sandstone Hill

Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R042CY151NM	Limestone Hills Limestone Hills occurs on limestone hills and footslopes of higher mountains. Slopes range from 10 to 60 percent. Soils are shallow, gravelly, and loamy. The reference plant community is mixed prairie grassland with scattered shrubs, forbs, and trees.
R042CY152NM	Shallow Shallow occurs on gravelly alluvial fans of higher limestone mountains. Slopes range from 5 to 20 percent. Soils are loamy, gravelly, and very shallow to a petrocalcic horizon also known as a root restricting layer. The reference plant community is mixed prairie grassland with scattered shrubs, trees, and forbs.
R042CY156NM	Gravelly Gravelly occurs on gravelly alluvial fans of higher limestone mountains. Slopes range from 5 to 20 percent. Soils are deep gravelly loams. The reference plant community is mixed prairie grassland with scattered shrubs, trees, and forbs.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

The site occurs on sandstone hills, hillslopes, and ridges. Slopes range from 10 to 60 percent.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Ridge
Elevation	1,372–1,676 m
Slope	5–20%
Aspect	N, S

Climatic features

The climate of this area is "semi-arid continental."

Annual average precipitation ranges from 11 to 19 inches. Variations of 5 inches, more or less, are not uncommon. Approximately 70 percent of the precipitation occurs from May through October. Most of the summer rain comes in the form of high-intensity, short- uration thunderstorms. Winter moisture is usually negligible.

Temperatures are characterized by distinct seasonal changes and large annual and diurnal temperature changes. The average annual temperature ranges from 55 degrees to 60 degrees, with extremes of 20 degrees below zero in the winter to 110 degrees in the summer not uncommon.

The average frost-free season is 170 to 189 days. The last killing frost is in early April and the first killing frost is in mid October.

Both temperature (especially south slope)and precipitation favor warm-season species. However, approximately 40 percent of the precipitation (and temperature on north slopes) is favorable to to cool season growth at the middle to higher elevations. This could allow the cool season plants to occupy a very important part of this complex plant community. Due to the shallow soil profile, vegetation responds well to light rains. Moisture can also be stored relatively deep in the seams and cracks of fractured limestone. This moisture is also available for plant use. Slope aspect is also important when the strong winds from the west and southwest blow. These winds, which blow from February to June, cause the soil to dry during the critical growth stage for many cool season species. The north slopes are somewhat protected and do not dry as fast as the south slopes.

Climate data was obtained from http://www.wrcc.sage.dri.edu/summary/climsmnm.html web site. Data interpreted utilizing NM NRCS Climate Summarizer spreadsheet.

Table 3. Representative climatic features

Frost-free period (average)	189 days
Freeze-free period (average)	211 days
Precipitation total (average)	483 mm

Influencing water features

None.

Soil features

The site consists of very shallow or shallow, well drained, and moderately permeable soils over very slowly permeable fractured sandstone bedrock. Parent material is loamy residuum and colluvium weathered from Permian sandstone. Typically, the surface layer is light brown, very channery sandy loam about 3.5 inches thick. The subsurface layer ranges from 3.5 to 11 inches in light brown extremely gravelly sandy loam. The underlying material from 11 inches to 21 inches is fractured sandstone bedrock.

The soil components and associated map units correlated to this site are:

Guadalupe Mountains National Park Soil Survey: Bonespring very gravelly loam, 1 to 8 percent slopes

Table 4. Representative soil features

Parent material	(1) Residuum-sandstone
Surface texture	(1) Very channery sandy loam(2) Very channery fine sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained

Permeability class	Very slow
Soil depth	8–46 cm
Surface fragment cover <=3"	10–35%
Surface fragment cover >3"	10–25%
Available water capacity (0-101.6cm)	0–5.08 cm
Calcium carbonate equivalent (0-101.6cm)	0–13%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	30–40%
Subsurface fragment volume >3" (Depth not specified)	4–15%

Ecological dynamics

The reference or current potential plant community for the Sandstone Hill ecological site is a mid and shortgrass and predominantly warm-season grama grassland interspersed with mixed shrubs. Grass, shrub and forb canopy cover ranges from 65-75, 5-20, and 1-5 percent, respectively. Surface fragment cover can range up to 75 percent. The scale of the plant community for measurement is the associated soil component within the soil map units.

Fluctuations in yearly weather conditions is probably the most influential factor that helps shape the plant communities. Prolonged droughts will dramatically reduce herbaceous cover and production while wet summers will favor black grama dominance and wet winters will favor more mesic and cool-season grasses (Ludwig et al. 2000). Aspect, elevation, soil variability, and percentage of rock outcrop also play a role in shape the inherent plant community. Lightning induced fires do occur in the area but the exact frequency is unknown.

It is unknown whether bison utilized this site historically. Slope, the rough topography, and lack of abundant perennial water may limited or prevented them from utilizing the site. Desert bighorn sheep probably utilized this site historically for part of their habitat needs. Other native fauna such as insects, rodents, mule deer, and pronghorn antelope are still helping influence the current plant communities.

The site is suited for prescribed grazing practices. Heavy continuous grazing, especially during dry years, over long periods will cause decreases in plants such as blue grama, black grama and increases in less palatable grasses such as sideoats grama, hairy grama, tridens, fluffgrass, and threeawns. In addition, woody plants may increase in areas where the seed source is available. Ecological processes such as water and nutrient cycles may also be affected by improper grazing management.

An inclusion plant community that is significantly different is also included in this ecological site due to its managerial importance within Guadalupe Mountains National Park. This Savannah State does not transition or restore to the Grassland or Shrubland State. Currently, this state has not been described outside the park. It is a gray oak – alligator juniper savanna with pine muhly, New Mexico muhly, cane bluestem, sideoats grama, New Mexico feathergrass, and blue grama as common grasses. This community grows in a wetter and cooler climate than the rest of the described Sandstone Hill ecological site. It grows in a more mesic portion of the ecological site is not a part of the dynamics of the warm-season grassland with shrubs that characterizes the majority of the Sandstone Hill site. The fire management plan of the park identifies this area as Madrean Evergreen Woodland. Fire plays an important role in maintaining an open savanna. The Woodland – Grassland dynamic potentially exists in this system.

The following diagram suggests general pathways that the vegetation on this site might follow. There are other plant communities and states not shown on the diagram. This information is intended to show what might happen in a given set of circumstances; it does not mean that this would happen the same way in every instance. Local professional guidance should always be sought before pursuing a treatment scenario.

Sandstone Hill

State and Transition Model:

State and transition model



Legend

1.1A, 2.1A, 3.1A, 3.2B: Heavy continuous grazing, exacerbated by drought 1.2A, 2.2A, 3.2A, 3.3A: Prescribed grazing and/or favorable precipitation T1A: Heavy continuous grazing, fire suppression, woody plant seed dispersal R2A: Brush control and/or prescribed fire.

State 1 Grassland State

The Grama/Shrubs plant community phase is the reference or current potential vegetative community. Characteristic grasses include sideoats grama, black grama, blue grama, Warnock's grama, curlyleaf muhly, and New Mexico feathergrass. Shifts in above average winter or summer precipitation can shift dominance from either black grama or sideoats to cool-season grasses. It is unknown if the higher shrub cover in some locations is attributed to past management or it is part of the natural variability of the site. Shrub canopy cover over 25 percent will categorize the site in the Shrubland State. Annual production ranges from 600 to 1400 pounds per acre. The Threeawns/Shrubs Community is a response to heavy continuous grazing. Drought will only exacerbate the situation. Perennial grass cover is lower and dominated by mainly perennial threeawns, hairy grama, and slim tridens. Both annual and perennial forbs will also increase in relative composition. Palatable grasses such as black grama and blue grama have decreased in cover. Shrubs may begin to encroach in certain areas.

Community 1.1 Gramas/Shrubs Community



Figure 4. 1.1 Gramas/Shrubs Community

The Grama/Shrubs Community is the reference or current potential vegetative community. Characteristic grasses include sideoats grama, black grama, blue grama, Warnock's grama, curlyleaf muhly, and New Mexico feathergrass. Shifts in above average winter or summer precipitation can shift dominance from either black grama or sideoats to cool-season grasses. In addition, the more mesic grasses such as sideoats grama, blue grama, and curlyleaf muhly will inherently be more abundant on north facing slopes while black grama will dominate south-facing slope. A variety of shrubs occurs sporadically and is characterized by winterfat, skeletonleaf goldeneye, ephedra, mariola, mountain mahogany, and New Mexico agave. Canopy cover of shrubs can range from 5-20 percent. It is unknown if the higher shrub cover in some locations is attributed to past management or it is part of the natural variability of the site. Shrub canopy cover over 25 percent will categorize the site in the Shrubland State. Annual production ranges from 600 to 1400 pounds per acre. The site is suited for livestock grazing and provides important wildlife habitat. Heavy continuous grazing over several years can reduce and potentially eliminate black and blue grama. Sideoats grama, threeawns, slim tridens, and hairy grama will increase. If there is an available seed source, shrubs may begin to encroach on the site.

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	673	919	1345
Shrub/Vine	84	157	168
Forb	28	45	56
Tree	_	-	-
Total	785	1121	1569

Table 5. Annual production by plant type

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	5-20%
Grass/grasslike foliar cover	50-65%
Forb foliar cover	1-5%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%

Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	-	0-2%	5-10%	1-3%
>0.15 <= 0.3	-	1-5%	30-35%	1-2%
>0.3 <= 0.6	-	1-5%	25-30%	-
>0.6 <= 1.4	-	3-10%	-	-
>1.4 <= 4	-	-	-	-
>4 <= 12	-	-	-	-
>12 <= 24	-	-	-	-
>24 <= 37	-	_	-	_
>37	-	_	-	_

Figure 6. Plant community growth curve (percent production by month). TX0030, Mixed Grama/Muhly Grassland Community. Mixed grama/muhly grassland characterized by blue grama, sideoats, grama, black grama, and curly leaf muhly.

Jai	n Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	1	3	3	10	15	25	25	10	5	1

Community 1.2 Threeawns/Shrubs Community

This plant community is a response to heavy continuous grazing. Drought will only exacerbate the situation. Perennial grass cover is lower and dominated by mainly perennial threeawns, hairy grama, and slim tridens. Both annual and perennial forbs will also increase in relative composition. Palatable grasses such as black grama and blue grama have decreased in cover. Shrubs may begin to encroach in certain areas.

Pathway 1.1A Community 1.1 to 1.2

Heavy continuous grazing is the major driver causing this transition. Drought will only hasten the transition. Palatable grasses favored by livestock such as blue grama and black grama have decreased in cover while less palatable grasses such as tridens, perennial threeawns, fluffgrass, sideoats grama, and hairy grama have increased in cover. Time period for this transition is probably less than 10 years.

Pathway 1.2A Community 1.2 to 1.1

A combination of favorable rainfall and conservation practices such as prescribed grazing and/or planned grazing system can help facilitate the recovery of grasses more palatable to livestock. Rate of recovery will depend on the extent to which the site has been disturbed.

Conservation practices

Prescribed Grazing

State 2 Shrubland State

The Shrubs/Grama Community is noted for lower perennial grass cover and is dominated by sideoats grama, slim tridens and threeawns. Both annual and perennial forbs will also increase in relative composition. Palatable grasses such as black grama and blue grama have decreased in cover. Shrub canopy cover is greater than 25 percent. The Shrubs/Threeawn Community has unknown past reference concerning whether if this community is attributed to past disturbances such as heavy continuous grazing or if these areas are within the inherent variability of the site. Nonetheless, shrubs can comprise greater than 25 percent cover in areas. Common shrubs include sotol, lechuguilla, sacahuista, and creosotebush. This community can provide habitat for some wildlife that the grassland community is not able to provide.

Community 2.1 Shrubs/Gramas Community

It is unknown if this community is attributed to past disturbances such as heavy continuous grazing or if these areas are within the inherent variability of the site. Nonetheless, shrubs can comprise greater than 25 percent cover in areas. Common shrubs include sotol, lechuguilla, sacahuista, and creosotebush. This community can provide habitat for some wildlife that the grassland community is not able to provide.

Community 2.2 Shrubs/Threeawns Community

This plant community is a response to heavy continuous grazing. Drought will only exacerbate the situation. Perennial grass cover is lower and dominated by sideoats grama, slim tridens and threeawns. Both annual and perennial forbs will also increase in relative composition. Palatable grasses such as black grama and blue grama have decreased in cover. Shrub canopy cover is greater than 25 percent.

Pathway 2.1A Community 2.1 to 2.2

Heavy continuous grazing is the major driver causing this transition. Drought will only hasten the transition. Palatable grasses favored by livestock such as blue grama and black grama have decreased in cover while less palatable grasses such as tridens, perennial threeawns, fluffgrass, sideoats grama, and hairy grama have increased in cover. Time period for this transition is probably less than 10 years.

Pathway 2.2A Community 2.2 to 2.1

A combination of favorable rainfall and conservation practices such as prescribed grazing and/or planned grazing system can help facilitate the recovery of grasses more palatable to livestock. Rate of recovery will depend on the extent to which the site has been disturbed.

Conservation practices

Prescribed Grazing

State 3 Savanna State (Inclusion State)

The Muhlys-Gramas/Oak-Juniper plant community is an inclusion that occurs in a cooler and wetter soil temperature regime. Soils still remain shallow to mostly sandstone with some areas derived from limestone. The reference plant community is a gray oak – alligator juniper savanna with abundant warm- and cool-season perennial bunchgrasses such as pine muhly, bull muhly, sideoats grama, and New Mexico feathergrass. Scattered shrubs include skunkbush sumac, cholla, and wavyleaf oak. Annual production ranges from 1350 to 2600 pounds per acre. The Threeawns/Oak-Juniper plant community phase results from overgrazing. A shift from decreaser grasses (palatable) to increaser (unpalatable) dominance has occurred. The community is more susceptible to woody plant

encroachment due the overall decline of herbaceous plants and the fuel to carry fire. Threeawns and hairy grama are common increaser grasses. Decreaser grasses include blue grama and the various muhlenbergia species. The Woodland/Mixed-grass plant community results from fire suppression and/or livestock overgrazing. A shift from decreaser grasses (palatable) to increaser (unpalatable) dominance has occurred. Further woody plant encroachment occurs due to the overall decline of herbaceous plants and the lack of fuel to carry fire. Threeawns and hairy grama are common increaser grasses. Decreaser grasses include blue grama and the various muhlenbergia species.

Community 3.1 Muhlys-Gramas/Oak-Juniper Community



Figure 7. 3.1 Muhlys-Gramas/Oak-Juniper Community

This plant community is an inclusion within the Bonespring map unit that occurs in a cooler and wetter soil temperature regime. Soils still remain shallow to mostly sandstone with some areas derived from limestone. At Guadalupe Mountains National Park, it is found on the north facing slopes of Pine Spring Canyon and the north face slopes of the foothills of Frijole Ridge. According to the park's Fire Management Plan, this area is referred to as the Madrean Evergreen Woodland. The reference plant community is a gray oak – alligator juniper savanna with abundant warm- and cool-season perennial bunchgrasses such as pine muhly, bull muhly, sideoats grama, and New Mexico feathergrass. Scattered shrubs include skunkbush sumac, cholla, and wavyleaf oak. Annual production ranges from 1350 to 2600 pounds per acre. Similar open savanna communities such as this one are typically maintained by a fire interval of 10-30 years (Abbott 1998). The few scattered woody plants allows for maximum herbaceous production that ultimately provides the fuel to carry fires. Fire suppression either alone or in combination with overgrazing will allow woody plants to increase at the expense of herbaceous plants. Conservation practices such as prescribed fire and prescribed grazing will help maintain an open savanna or oak woodland.

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1121	1569	2018
Tree	224	448	560
Shrub/Vine	112	224	336
Forb	56	84	112
Total	1513	2325	3026

Table 8. Annual production by plant type

Community 3.2 Threeawns/Oak-Juniper Community

This plant community phase results from overgrazing. A shift from decreaser grasses (palatable) to increaser (unpalatable) dominance has occurred. The community is more susceptible to woody plant encroachment due the overall decline of herbaceous plants and the fuel to carry fire. Threeawns and hairy grama are common increaser grasses. Decreaser grasses include blue grama and the various muhlenbergia species.

Community 3.3 Woodland/Mixed-grass Community

This plant community phase results from fire suppression and/or livestock overgrazing. A shift from decreaser grasses (palatable) to increaser (unpalatable) dominance has occurred. Further woody plant encroachment occurs due to the overall decline of herbaceous plants and the lack of fuel to carry fire. Threeawns and hairy grama are common increaser grasses. Decreaser grasses include blue grama and the various muhlenbergia species.

Pathway 3.1A Community 3.1 to 3.2

The Muhlys-Gramas/Oak-Juniper Community transitions into the Threeawns/Oak-Juniper Community through the use of heavy continuous grazing and exacerbated drought conditions.

Pathway 3.2A Community 3.2 to 3.1

The shift from the Threeawns/Oak-Juniper Community to the Muhlys-Gramas/Oak-Juniper Community occurs when Prescribed Grazing is applied and/or favorable precipitation exists.

Conservation practices

Prescribed Grazing

Pathway 3.2B Community 3.2 to 3.3

The Threeawns/Oak-Juniper Community transitions into the Woodland/Mixed-grass Community through the use of heavy continuous grazing and exacerbated drought conditions.

Pathway 3.3A Community 3.3 to 3.2

The shift from the Woodland/Mixed-grass Community to the Threeawns/Oak-Juniper Community occurs when Prescribed Grazing and/or favorable precipitation is applied.

Conservation practices

Prescribed Grazing

Transition T1A State 1 to 2

The grassland state transitions to the shrubland state when shrub cover is higher than 25 percent. It is unknown if the transition is attributed to past disturbances such as heavy continuous grazing or if these areas are within the inherent variability of the site. Nonetheless, shrubs can comprise greater than 25 percent cover in areas. Common increaser shrubs include sotol, broomweed, sacahuista, and mariola.

Restoration pathway R2A State 2 to 1

Rangeland conservation practices such as brush management and possibly prescribed burning (where amount of fine fuel is adequate) can restore the Shrubland State into the Grassland State.

Conservation practices

Brush Management

Prescribed Grazing

Additional community tables

Table 9. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike		<u>.</u>		
1	Warm-season stolonifero	us grasses	3	224–336	
	black grama	BOER4	Bouteloua eriopoda	224–336	_
2	Dominant warm-season b	unchgras	Ses	224–336	
	sideoats grama	BOCU	Bouteloua curtipendula	112–224	-
	blue grama	BOGR2	Bouteloua gracilis	56–168	-
	Warnock's grama	BOWA	Bouteloua warnockii	56–168	_
3	Subdominant warm-sease	on bunchg	rasses	84–112	
	hairy grama	BOHI2	Bouteloua hirsuta	17–34	-
	common wolfstail	LYPH	Lycurus phleoides	17–34	-
	Hall's panicgrass	PAHA	Panicum hallii	17–34	-
	hairy woollygrass	ERPI5	Erioneuron pilosum	11–28	-
	threeawn	ARIST	Aristida	11–28	-
	sand dropseed	SPCR	Sporobolus cryptandrus	11–28	-
	slim tridens	TRMU	Tridens muticus	11–28	-
	curlyleaf muhly	MUSE	Muhlenbergia setifolia	6–17	_
	plains lovegrass	ERIN	Eragrostis intermedia	6–17	-
	streambed bristlegrass	SELE6	Setaria leucopila	6–17	-
4	Cool-season bunchgrasse	es		84–112	
	southwestern needlegrass	ACEM4	Achnatherum eminens	67–168	-
	New Mexico feathergrass	HENE5	Hesperostipa neomexicana	17–34	-
Shrub	/Vine				
5	Shrubs			34–67	
	viscid acacia	ACNE4	Acacia neovernicosa	6–22	_
	desert ceanothus	CEGR	Ceanothus greggii	6–22	_
	alderleaf mountain mahogany	CEMO2	Cercocarpus montanus	6–22	-
	Guadalupe rabbitbrush	CHSP3	Chrysothamnus spathulatus	6–22	-
	javelina bush	COER5	Condalia ericoides	6–22	-
	rough jointfir	EPAS	Ephedra aspera	6–22	_
	eggleaf silktassel	GAOV	Garrya ovata	6–22	-
	Pinchot's juniper	JUPI	Juniperus pinchotii	6–22	_
	winterfat	KRLA2	Krascheninnikovia lanata	6–22	_
	Big Bend barometerbush	LEMI4	Leucophyllum minus	6–22	_
	algerita	MATR3	Mahonia trifoliolata	6–22	
	Rio Grande saddlebush	MOSC	Mortonia scabrella	6–22	_
	mariola	PAIN2	Parthenium incanum	6–22	-

6 Subshrubs ········· 6-11 1 Italelerplume DAFO Jalea formosa		skunkbush sumac	RHTR	Rhus trilobata	6–22	-				
Instant Part Part Part Part Part Part Part Par	6	Subshrubs	•		6–11					
Ittelear ratanyKRERKrameria erectaCallCallwody crinklematTCACTquilia canescens var. canescensCallCallFibrourSucculentsICACValuia canescens var. canescensCallCallParly's agaveAGPANAgave party sp. neomexicanaCallCallI'ee chollaCIYIIIValuro puntia imbricata var. indicataCallCallgreen sotolDAL22Dasylrion leiophyllumCallCallTexas sacchuistaNOTENolina texanaCallCallpricklypearOPUNTOpuntia calcalaCallCalla parter succaYUBAVucca baccataCallCallsapatre succaYUBAVucca alcalaCallCallb forbFortsValura datasithifolaCallCallc forb perennialSPABBahia absinthifolaCallCalla lainterwedGRO5Coton potistiCallCalla lainterwedGUSAGuierzi sarothraCallCalla laintersional fullBakaJalacarosa var. polymenidesCalla laintersional fullMacMoMacoraCalla laintersional fullGUSAGuierzi sarothraCalla laintersional fullMacoraMacoraCalla laintersional fullMacoraCallCalla laintersional fullMacoraCallCalla laintersional fullMacoraCallCalla laintersional fullGuierzi sar		featherplume	DAFO	Dalea formosa	2–6	-				
woody crinklematTICACTiquilia canescens var. canescens2-67Fibrous/Succulents17-342Parry's agaveAGPAN6Agave parryi ssp. neomexicana6-174tree chollaCYIMICylindropuntia imbricata var. imbricata var. imbricata6-175green sotolDALE2Dasylirion leiophyllum6-176Texas sacahuistaNOTENolina texana6-177pricklypearOPUNTOpuntia6-176banana yuccaYUBAYucca baccata6-177banana yuccaYUELYucca baccata6-177banana yuccaYUELYucca elata6-177banana yuccaYUELYucca leata6-177banana yuccaYUELYucca leata6-177banana yuccaYUELYucca leata6-177banana yuccaYUELYucca leata6-177banana yuccaYUELYucca leata6-178ForbForb, perennial6-179ForbSorbonSorbon11-39Forb, perennialSABBahia absinthifolia1-310IeathenweedGROSGroton pottsi1-311BABBahia absinthifolia1-312buckwheatEGUSGuiterrzia sarothrae1-3		littleleaf ratany	KRER	Krameria erecta	2–6	-				
7Fibrous/SucculentsAGPAN6Agave parryi sap. neomexicana1.17-341Parry's agaveAGPAN6Agave parryi sap. neomexicana6.17.1tee chollaCYIM1Cylindropunti nibricata var. mibricata6.17.1green sotolDALE2Dasylirion leiophyllum6.171.1Texas sacahuistaNOTENotne texana6.171.1Texas sacahuistaNOTENotne texana6.171.1picktypearOPUNTOpuntia.6.171.2banan yuccaYUE0Yucca baccata.6.1617.3baran yuccaYUE0Yucca elata.6.1617.5ForbForb6Forbs7PorbsStata absintifolia8Forbs9Porby soundeflowerDECODesmathus cooleyi10blokwheatERIOSGuienroza sarchirae11BardongueMEL2Melamodum leucanthum10blokwheatERIOSGuienroza sarchirae <td></td> <td>woody crinklemat</td> <td>TICAC</td> <td>Tiquilia canescens var. canescens</td> <td>2–6</td> <td>-</td>		woody crinklemat	TICAC	Tiquilia canescens var. canescens	2–6	-				
Parny's agaveAGPAN6Agave parnyl ssp. neomexicana6-17Iree chollaCYIMICylindropuntia imbricata var. imbricata6-17green sotolDALE2Dasylirion leiophyllum6-17Texas sacahuistaNOTENolina texana6-17pricklypearOPUNTOpuntia6-17soture vuccaYUEAYucca baccata6-17soture vuccaYUEAYucca elata6-17Forb8Forb8Forbs11-391BaABBaha absinthifolia1-31hairyseed bahiaBAABBahia absinthifolia1-31IeatherweedCRPO5Croton pottsii1-31boom snakeweedGUSA2Gutierrezia sarothrae1-31needleleaf bluetHOACP2Houstoria acerosa var. polypremoides1-31nenodoraMELE2Menodora1-31nenodoraMENDMenodora1-31polygalaPOLYGPolygala1-31noseburnTRAGITragia1-39ForbsTragia1-31-39ForbsTragia Polygala1-31-39ForbTragia polygala0-141-39ForbTragia polygala0-151-39ForbTragia polygala0-111-39ForbPLA2Plantago polygala0-119ForbTragia1-30-1<	7	Fibrous/Succulents	-		17–34					
Itee chollaCYIMI CYIMI imbricataCylindropuntia imbricata var. imbricata6-17-green sotolDALE2Dasylirion leiophyllum6-17-Texas sacahuistaNOTENolina texana6-17-pricklypearOPUNTOpuntia6-17-banana yuccaYUBAYucca elata6-17-soatree yuccaYUEAYucca elata6-17-Forb111-39Porbs111-39PorbSoatree yuccaCRPO5Croton portsii1-hairyseed bahiaBAABBahia absinthifolia1<13		Parry's agave	AGPAN6	Agave parryi ssp. neomexicana	6–17	-				
green sololDALE2Dasylirion leiophyllum6–17Texas sacahuistaNOTENolina texana6–17pricklypearOPUNTOpuntia6–17banana yuccaYUBAYucca baccata6–17soaptree yuccaYUELYucca elata6–17ForbsSoaptree yuccaYUELYucca elata6–17ForbsSoaptree yuccaZEPForb, perennial7–17ForbForb, perennial2FPForb, perennial7–17Ihairyseed bahiaBAABBahia absinthifolia1IleatherweedCRPO5Croton pottsii1IbuckwheatERIOCDesmanthus cooleyi1Ibroom snakeweedGUSA2Gutierrezia sarothrae1Ipeldeleaf bluetMELE2Melampodium leucanthum1IpolygalaPOLYGPolygalaInoseburnTRAGITragiaIForbsTradiTragiaIForb, annual2FAForb, annualISartereetSartereetIISartereetSartereet		tree cholla	CYIMI	Cylindropuntia imbricata var. imbricata	6–17	-				
Texas sacahuistaNOTENolina texana6-17-pricklypearOPUNTOpuntia6-17-banana yuccaYUBAYucca baccata6-617-soaptree yuccaYUELYucca elata6-617-ForbForbs11-39Intersect8 Forbs11-39Intersect11-39IntersectIntersectIntersectIntersectCooley's bundleflowerDECO2Desmanthus cooleyi11-39Intersect <td <="" colspan="4" td=""><td></td><td>green sotol</td><td>DALE2</td><td>Dasylirion leiophyllum</td><td>6–17</td><td>-</td></td>	<td></td> <td>green sotol</td> <td>DALE2</td> <td>Dasylirion leiophyllum</td> <td>6–17</td> <td>-</td>					green sotol	DALE2	Dasylirion leiophyllum	6–17	-
pricklypearOPUNT $Opuntia$ $Opuntia$ $6-17$ $-$ banana yuccaYUBAYucca baccata $6-17$ $-$ soaptree yuccaYUELYucca elata $6-17$ $-$ ForbForbs8Forbs 5 5 5 $11-39$ $-$ 10hairyseed bahiaBAABBahia absinthifolia $1-3$ $ -$ 11elatherweedCRPO5Croton pottsii $1-3$ $ -$ 11buckwheatDECO2Desmanthus cooleyi $1-3$ $ -$ 11buckwheatERIOGEriogonum $1-3$ $ -$ 11broom snakeweedGUSA2Gutierrezia sarothrae $1-3$ $ -$ 11plains blackfootMELE2Melampodium leucanthum $1-3$ $ -$ 11polygalaPOLYGPolygala $ -$ 11SeaburnTRAGITraja $ -$ 12Forb, annualEFAForb, annual $ -$ 13DestructureGUSA2Gutierrezia sarothrae $ -$ 14blains blackfootMELE2Melampodium leucanthum $ -$ </td <td></td> <td>Texas sacahuista</td> <td>NOTE</td> <td>Nolina texana</td> <td>6–17</td> <td>-</td>		Texas sacahuista	NOTE	Nolina texana	6–17	-				
banana yuccaYUBAYucca baccata6-17soaptree yuccaYUELYucca elata6-17-Forbs8Forbs11-391Forb, perennial2FPForb, perennial7-17-1hairyseed bahiaBAABBahia absinthifolia11-31leatherweedCRPO5Croton pottsii11-32Cooley's bundleflowerDEC02Desmanthus cooleyi11-34buckwheatERIOGEriogonum11-35broom snakeweedGUSA2Gutierrezia sarothrae11-36plains blackfootMELE2Melampodium leucanthum11-37plains blackfootMELE2Penstemon9polygalaPUYGPolygala9ForbsTRAGITraja9Forb, annual2FAForb, annual0-119wolly plantainPLA2Planago patagonica0-119wolly plantainPLA2Planago patagonica0-111wolly plantainPLA2Planago patagonica0-11		pricklypear	OPUNT	Opuntia	6–17	-				
soaptree yuccaYUELYucca elata6-17-Forb8Forbs11-399Forb, perennial2FPForb, perennial7-1710hairyseed bahiaBAABBahia absinthifolia1-310leatherweedCRPO5Croton pottsii1-310Cooley's bundleflowerDEC02Desmanthus cooleyi1-310buckwheatERIOGEriogonum1-311broom snakeweedGUSA2Gutierrezia sarothrae1-311needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1-311plains blackfootMELE2Melampodium leucanthum1-311polygalaPENSTPenstemon1-311ingstalk greenthreadTHLOThelesperma longipes1-311ForbsTragia1-3-11Forb, annualZFAForb, annual0-1111Volly plantainPLPA2Plantago patagonica0-11		banana yucca	YUBA	Yucca baccata	6–17	-				
Forbs11-398Forb, perennial2FPForb, perennial7-777-71hairy seed bahiaBAABBahia absinthifolia1-37-71leatherweedCRPO5Croton pottsii1-37-71leatherweedDEC02Desmanthus cooleyi1-37-72Cooley's bundleflowerDEC02Desmanthus cooleyi1-37-72buckwheatERIOGEriogonum1-37-73broom snakeweedGUSA2Gutierrezia sarothrae1-37-74plains blackfootMEL02Meutonia acerosa var. polypremoides1-37-74plains blackfootMEL22Melampodium leucanthum1-37-74plains blackfootMEL02Mendora1-37-75polygalaPENSTPenstermon Ingipes1-37-79ForbsTHLOTragia7-77-77forb, annual2FAForb, annual0-117-79Wolly plantainPLPA2Plantago palagonica0-117-7		soaptree yucca	YUEL	Yucca elata	6–17	-				
8Forbs11-391Forb, perennial2FPForb, perennial7-171hairyseed bahiaBAABBahia absinthifolia1-31leatherweedCRPO5Croton pottsii1-31-31beckwheatDECO2Desmanthus cooleyi1-31-31buckwheatERIOGFriogonum1-31-31broom snakeweedGUSA2Gutierrezia sarothrae1-31-31needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1-31-31plains blackfootMELE2Melampodium leucanthum1-31-31beardtonguePENSTPensternon1-31-31Iongstalk greenthreadTHLOThelesperma longipes1-31-39ForbsTragia1-31-31-39Forb, annual2FAForb, annual0-111-31Voolly plantainPLPA2Plantago patagonica0-111-3	Forb		-							
Forb, perennial2FPForb, perennial7-17-hairyseed bahiaBAABBahia absinthifolia1-31-3leatherweedCRPO5Croton pottsii1-31-3Cooley's bundleflowerDEC02Desmanthus cooleyi1-31-3buckwheatERIOGEriogonum1-31-3broom snakeweedGUSA2Gutierrezia sarothrae1-31-3needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1-31-3menodoraMELE2Melampodium leucanthum1-31-3beardtonguePENSTPenstemon1-31-3polygalaPOLYGPolygala1-31-3polygalaTHLOThelesperma longipes1-31-39ForbsTragia1-31-39Forb, annual2FAForb, annual0-111-3woolly plantainPLPA2Plantago patagonica0-111-3woolly plantainPLPA2Plantago patagonica0-111-3	8	Forbs			11–39					
hairyseed bahiaBAABBahia absinthifolia13leatherweedCRPO5Croton pottsii1313Cooley's bundleflowerDEC02Desmanthus cooleyi1313buckwheatERIOGEriogonum1313broom snakeweedGUSA2Gutierrezia sarothrae1313needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1313plains blackfootMELE2Melampodium leucanthum13beardtonguePENSTPenstemon13polygalaPOLYGPolygala13onseburnTRAGITragia13forb, annual2FAForb, annual0-11woolly plantainPLPA2Plantago patagonica011		Forb, perennial	2FP	Forb, perennial	7–17	-				
leatherweedCRPO5Croton pottsii1-3Cooley's bundleflowerDECO2Desmanthus cooleyi1-3buckwheatERIOGEriogonum1-3broom snakeweedGUSA2Gutierrezia sarothrae1-3needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1-3menodoraMELE2Melampodium leucanthum1-3menodoraMENODMenodora1-3beardtonguePENSTPenstemon1-3polygalaPOLYGPolygala1-3noseburnTRAGITragia1-3Forb, annual2FAForb, annual0-11-woolly plantainPLPA2Plantago patagonica0-11-woolly plantainPLPA2Plantago patagonica0-11-		hairyseed bahia	BAAB	Bahia absinthifolia	1–3	-				
Cooley's bundleflowerDECO2Desmanthus cooleyi1–31–3buckwheatERIOGEriogonum1–31–31–3broom snakeweedGUSA2Gutierrezia sarothrae1–31–3needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1–31–3plains blackfootMELE2Melampodium leucanthum1–31–3menodoraMENODMenodora1–31–3beardtonguePENSTPenstemon1–31–3polygalaPOLYGPolygala1–31–3onseburnTRAGITragia1–31–39ForbsZFAForb, annual0–111–3woolly plantainPLPA2Plantago patagonica0–111–3		leatherweed	CRPO5	Croton pottsii	1–3	-				
buckwheatERIOGEriogonum1-31-3broom snakeweedGUSA2Gutierrezia sarothrae1-31-3needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1-31-3plains blackfootMELE2Melampodium leucanthum1-31-3menodoraMENODMenodora1-31-3beardtonguePENSTPenstemon1-31-3polygalaPOLYGPolygala1-31-3poseburnTRAGITragia1-31-39ForbsTragia1-31-39Forb, annual2FAForb, annual0-111-3woolly plantainPLPA2Plantago patagonica0-111-3		Cooley's bundleflower	DECO2	Desmanthus cooleyi	1–3	-				
broom snakeweedGUSA2Gutierrezia sarothrae1–3–needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1–31–3–plains blackfootMELE2Melampodium leucanthum1–3–menodoraMENODMenodora1–3–beardtonguePENSTPenstemon1–3–polygalaPOLYGPolygala1–3–iongstalk greenthreadTHLOThelesperma longipes1–3–9ForbsTragia1–3–9Forb, annual2FAForb, annual0–11–woolly plantainPLPA2Plantago patagonica0–11–		buckwheat	ERIOG	Eriogonum	1–3	-				
needleleaf bluetHOACP2Houstonia acerosa var. polypremoides1-313plains blackfootMELE2Melampodium leucanthum13-menodoraMENODMenodora13-beardtonguePENSTPenstemon13-polygalaPOLYGPolygala13-longstalk greenthreadTHLOThelesperma longipes13-9Forbs7-177-17-Forb, annual2FAForb, annual0-11-woolly plantainPLPA2Plantago patagonica0-11-		broom snakeweed	GUSA2	Gutierrezia sarothrae	1–3	-				
plains blackfootMELE2Melampodium leucanthum1-3menodoraMENODMenodora1-3beardtonguePENSTPenstemon1-3polygalaPOLYGPolygala1-3longstalk greenthreadTHLOThelesperma longipes1-39ForbsTragia7-177-17Forb, annual2FAForb, annual0-11woolly plantainPLPA2Plantago patagonica0-11		needleleaf bluet	HOACP2	Houstonia acerosa var. polypremoides	1–3	-				
menodoraMENODMenodora1-3-beardtonguePENSTPenstemon1-3-polygalaPOLYGPolygala1-3-longstalk greenthreadTHLOThelesperma longipes1-3-noseburnTRAGITragia1-3-9Forbs7-177-17Forb, annual2FAForb, annual0-11-woolly plantainPLPA2Plantago patagonica0-11-		plains blackfoot	MELE2	Melampodium leucanthum	1–3	-				
beardtonguePENSTPenstemon1-3polygalaPOLYGPolygala1-3longstalk greenthreadTHLOThelesperma longipes1-3noseburnTRAGITragia1-39Forbs7-177-17Forb, annual2FAForb, annual0-11-woolly plantainPLPA2Plantago patagonica0-11-		menodora	MENOD	Menodora	1–3	-				
polygalaPOLYGPolygala1-3-longstalk greenthreadTHLOThelesperma longipes1-3-noseburnTRAGITragia1-3-9Forbs7-177-171Forb, annual2FAForb, annual0-11-woolly plantainPLPA2Plantago patagonica0-11-		beardtongue	PENST	Penstemon	1–3	-				
Iongstalk greenthreadTHLOThelesperma longipes1–3–noseburnTRAGITragia1–3–9Forbs7–177–171Forb, annual2FAForb, annual0–11–1woolly plantainPLPA2Plantago patagonica0–11–		polygala	POLYG	Polygala	1–3	-				
noseburnTRAGITragia1-3-9Forbs7-177Forb, annual2FAForb, annual0-11-woolly plantainPLPA2Plantago patagonica0-11-		longstalk greenthread	THLO	Thelesperma longipes	1–3	-				
9 Forbs 7–17 Forb, annual 2FA Forb, annual 0–11 – woolly plantain PLPA2 Plantago patagonica 0–11 –		noseburn	TRAGI	Tragia	1–3	-				
Forb, annual2FAForb, annual0–11–woolly plantainPLPA2Plantago patagonica0–11–	9	Forbs			7–17					
woolly plantain PLPA2 Plantago patagonica 0-11 -		Forb, annual	2FA	Forb, annual	0–11	-				
		woolly plantain	PLPA2	Plantago patagonica	0–11	-				

Table 10. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)			
Grass	Grasslike							
1	Muhlenbergia spp.			504–908				
	pine muhly	MUDU	Muhlenbergia dubia	140–336	-			
	bullgrass	MUEM	Muhlenbergia emersleyi	140–336	-			
	New Mexico muhly	MUPA2	Muhlenbergia pauciflora	112–280	_			
	slimflower muhly	MUTE	Muhlenbergia tenuiflora	112–280	-			
2	Midgrasses			336–605				
	cane bluestem	BOBA3	Bothriochloa barbinodis	84–280	_			

	sideoats grama	BOCU	Bouteloua curtipendula	84–280	-
	blue grama	BOGR2	Bouteloua gracilis	84–280	-
	plains lovegrass	ERIN	Eragrostis intermedia	84–280	-
	common wolfstail	LYPH	Lycurus phleoides	56–168	_
3	Shortgrasses	-		168–303	
	threeawn	ARIST	Aristida	56–112	_
	hairy grama	BOHI2	Bouteloua hirsuta	56–112	_
	hairy woollygrass	ERPI5	Erioneuron pilosum	56–112	_
4	Cool-season bunchgra	sses	-	112–202	
	New Mexico feathergrass	HENE5	Hesperostipa neomexicana	84–168	_
	pinyon ricegrass	PIFI	Piptochaetium fimbriatum	28–101	-
Shrub	/Vine	-	-		
5	Shrubs			84–224	
	white sagebrush	ARLUM2	Artemisia ludoviciana ssp. mexicana	28–84	Ι
	brickellbush	BRICK	Brickellia	28–84	-
	featherplume	DAFO	Dalea formosa	28–84	-
	catclaw mimosa	MIACB	Mimosa aculeaticarpa var. biuncifera	28–84	_
	pungent oak	QUPU	Quercus pungens	28–84	-
	fragrant sumac	RHAR4	Rhus aromatica	28–84	-
	littleleaf sumac	RHMI3	Rhus microphylla	28–84	Ι
6	Fibrous/Succulents	-		28–112	
	tree cholla	CYIMI	Cylindropuntia imbricata var. imbricata	6–28	-
	Texas sacahuista	NOTE	Nolina texana	6–28	-
	tulip pricklypear	OPPH	Opuntia phaeacantha	11–28	-
	soaptree yucca	YUEL	Yucca elata	6–28	-
Forb		-			
7	Forbs			56–112	
	Forb, perennial	2FP	Forb, perennial	22–56	-
	awnless bushsunflower	SICA7	Simsia calva	11–28	_
	Forb, annual	2FA	Forb, annual	11–28	-
	vervain	VERBE	Verbena	3–11	_
	milkvetch	ASTRA	Astragalus	3–11	-
Tree					
8	Trees			224–560	
	alligator juniper	JUDE2	Juniperus deppeana	112–448	_
	gray oak	QUGR3	Quercus grisea	112–448	-
	Texas madrone	ARXA80	Arbutus xalapensis	56–224	_

Animal community

The reference plant community is suited for conservative livestock grazing by cattle, horses, burros, and sheep and goats. Livestock should be stocked in proportion to the grazeable grass, forbs, and browse. Improper grazing

management, especially during droughts, causes a gradual decline in range health, reducing livestock nutrition and habitat quality for some wildlife.

Wildlife that use this site for at least a portion of their overall habitat needs include mule deer, pronghorn antelope, desert bighorn sheep, coyotes, black-tailed jackrabbits, cottontails, mice, and ground squirrels. Birds that use this site as for at least a portion of their habitat needs include scaled quail, dove, raptors, and numerous song birds. Insects and reptiles such as rattlesnakes and lizards also frequent the area.

Plant Preference by Animal Kind:

These preferences are somewhat general in nature as the preferences for plants is dependent upon grazing experience, time of year, availability of choices, and total forage supply.

Legend: P=Preferred D=Desirable U=Undesirable N=Not Consumed T=Toxic X=Used, but not degree of utilization unknown

Preferred - Percentage of plant in animal diet is greater than it occurs on the land

Desirable - Percentage of plant in animal diet is similar to the percentage composition on the land

Undesirable - Percentage of plant in animal diet is less than it occurs on the land

Not Consumed – Plant would not be eaten under normal conditions. Only consumed when other forages are not available.

Toxic – Rare occurrence in diet and, if consumed in any tangible amounts results in death or severe illness in animal

Hydrological functions

The site is located in both a surface water run-in and runoff position. Surface rock fragments and a high grass cover help reduce runoff potential especially on hill slopes. A reduction in grass and ground cover will impair the hydrologic function of the site by increasing surface runoff and decreasing water infiltration.

Recreational uses

The site can be used for hiking and hunting.

Wood products

None.

Other products

None.

Other information

None.

Inventory data references

Information presented here has been developed from NRCS clipping, composition, plant cover, soils data and ecological interpretations gained by field observation.

Other references

Reviewers: Mark Moseley, Rangeland Management Specialist, NRCS, Boerne, TX

Contributors

Michael Margo

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: