

Ecological site F043AX957MT

Lower Subalpine Frigid Coniferous western redcedar (*Thuja plicata*)- western hemlock (*Tsuga heterophylla*)

Last updated: 9/08/2023
Accessed: 04/20/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043A–Northern Rocky Mountains

This MLRA is located in Montana (43 percent), Idaho (34 percent), and Washington (23 percent). It makes up about 31,435 square miles (81,460 square kilometers). It has no large cities or towns. It has many national forests, including the Okanogan, Colville, Kootenai, Lolo, Flathead, Coeur d'Alene, St. Joe, Clearwater, and Kaniksu National Forests.

This MLRA is in the Northern Rocky Mountains Province of the Rocky Mountain System. It is characterized by rugged, glaciated mountains; thrust- and block-faulted mountains; and hills and valleys. Steep-gradient rivers have cut deep canyons. Natural and manmade lakes are common.

The major Hydrologic Unit Areas (identified by four-digit numbers) that make up this MLRA are: Kootenai-Pend Oreille-Spokane (1701), 67 percent; Upper Columbia (1702), 18 percent; and Lower Snake (1706), 15 percent. Numerous rivers originate in or flow through this area, including the Sanpoil, Columbia, Pend Oreille, Kootenai, St. Joe, Thompson, and Flathead Rivers.

This area is underlain primarily by stacked slabs of layered sedimentary or metasedimentary bedrock. The bedrock formations range from Precambrian to Cretaceous in age. The rocks consist of shale, sandstone, siltstone, limestone, argillite, quartzite, gneiss, schist, dolomite, basalt, and granite. The formations have been faulted and stacked into a series of imbricate slabs by regional tectonic activity. Pleistocene glaciers carved a rugged landscape that includes sculpted hills and narrow valleys filled with till and outwash. Continental glaciation overrode the landscape in the northern half of the MLRA while glaciation in the southern half was confined to montane settings.

The average annual precipitation is 25 to 60 inches (635 to 1,525 millimeters) in most of this area, but it is as much as 113 inches (2,870 millimeters) in the mountains and is 10 to 15 inches (255 to 380 millimeters) in the western part of the area. Summers are dry. Most of the precipitation during fall, winter, and spring is snow. The average annual temperature is 32 to 51 degrees F (0 to 11 degrees C) in most of the area, decreasing with elevation. In most of the area, the freeze-free period averages 140 days and ranges from 65 to 215 days. It is longest in the low valleys of Washington, and it decreases in length with elevation. Freezing temperatures occur every month of the year on high mountains, and some peaks have a continuous cover of snow and ice.

The dominant soil orders in this MLRA are Andisols, Inceptisols, and Alfisols. Many of the soils are influenced by Mount Mazama ash deposits. The soils in the area have a frigid or cryic soil temperature regime; have an ustic, xeric, or udic soil moisture regime; and dominantly have mixed mineralogy. They are shallow to very deep, are very poorly drained to well drained, and have most of the soil texture classes. The soils at the lower elevations include Udivitrands, Vitrixerands and Haplustalfs. The soils at the higher elevations include Dystrocrypts, Eutrocrypts, Vitricryands, and Haplocryalfs. Cryorthents, Cryepts, and areas of rock outcrop are on ridges and peaks above timberline

This area is in the northern part of the Northern Rocky Mountains. Grand fir, Douglas-fir, western red cedar, western hemlock, western larch, lodgepole pine, subalpine fir, ponderosa pine, whitebark pine, and western white pine are the dominant overstory species, depending on precipitation, temperature, elevation, and landform aspect. The understory vegetation varies, also depending on climatic and landform factors. Some of the major wildlife species in this area are whitetailed deer, mule deer, elk, moose, black bear, grizzly bear, coyote, fox, and grouse. Fish, mostly in the trout and salmon families, are abundant in streams, rivers, and lakes.

More than one-half of this area is federally owned and administered by the U.S. Department of Agriculture, Forest Service. Much of the privately-owned land is controlled by large commercial timber companies. The forested areas are used for wildlife habitat, recreation, watershed, livestock grazing, and timber production. Meadows provide summer grazing for livestock and big game animals. Less than 3 percent of the area is cropland.

LRU notes

This ecological site resides in MLRA 43A in the Livingston-Lewis-Apgar Mountains which includes the bulk of Glacier National Park (GNP) and the lower western valley portions along the Flathead River. The landscape is mountains and landforms include glaciated mountains with associated features such as U-shaped valleys, mountain slopes, alpine ridges, cirques, valley floors and moraines. Glaciation of this area was in the form of alpine, icecaps and valley outlet glaciers. It also includes associated alluvium and outwash features. This area includes low valleys to tall mountains with elevation ranging 989-2,762 m (3,250-9,050 ft.). The climate is cold and wet with mean annual air temperature of 3 degrees Celsius (37 degrees F), mean frost free days of 65 days and mean annual precipitation of 1,295 mm (51 in.) and relative effective annual precipitation is 169 cm (66 in.). The soil temperature regime is cryic and the soil moisture regime is udic. The geology of this area is dominated by metasedimentary rocks of the Belt Supergroup (Grinnell argillite and Siyeh limestone) with minor Tertiary sediments. Soils are generally weakly developed on mountain slopes within U-shaped valleys. Parent materials are commonly of colluvium, till, and residuum from metasedimentary rocks. Limestone bedrock within this part of the Belt Supergroup is not highly calcareous and due to high precipitation received in this area most carbonates at mid and upper elevations have been leached from the soil profiles. Bedrock depth varies greatly with location, landform and slope position. Volcanic ash is often found in the soil surface with various degrees of mixing. Thicker volcanic ash can be found on more stable positions on mid and upper elevation slopes that are protected from wind erosion. Volcanic ash is not typically found in low elevation areas on stream and outwash terraces associated with streams and rivers. There are numerous large lakes including St. Mary, Bowman, Kintla, Lake Sherburne, Logging, Upper Waterton and numerous creeks (

Classification relationships

This ecological site relates to the USFS Habitat Type THSE/CLUN2 & THPL/CLUN2. This site relates to the USFS Habitat Type Group 5 and Fire Group 11. Both of these classification guides are specifically for the western Montana and northern Idaho region.

This ecological site relates to the National Park Service, NatureServe classification, *Thuja plicata/Clintonia uniflora* Forest (CEGL000474)

Ecological site concept

Ecological Site Concept

The lower subalpine frigid coniferous site is found west of the Continental Divide in moist areas within this maritime climate in glacial valley wall and lateral moraine landforms on back and foot slope positions, and on all aspects, spanning elevations from 1,000 to 1,500 meters (3,280-4,920 feet). The Reference Community is dominated by western cedar and western hemlock, with early seral tree species constrained to less than 3% of the overstory canopy. The ground cover consists predominantly of duff (63%) with high cover of moss (29%), and low cover of embedded litter (5%) and trace stones. The vegetation structure is that of very tall trees from 42-92 feet tall of western larch, western redcedar and western hemlock. The understory is multistoried though fairly sparse. The tallest understory layer is 51-102 cm or 20-40 inches tall and can include common ladyfern (*Athyrium filix-femina*), common snowberry (*Symphoricarpos albus*), white spirea (*Spiraea betulifolia*), and thinleaf huckleberry (*Vaccinium membranaceum*). The lowest layer is less than 25 cm or 10 inches tall and can include fireweed (*Chamerion angustifolium*), pipsissewa (*Chimaphila umbellata*), queencup beadlily (*Clintonia uniflora*), western rattlesnake plantain (*Goodyera oblongifolia*), twinflower (*Linnaea borealis*), threeleaf foamflower (*Tiarella trifoliata*), and Oregon

boxleaf (*Paxistima myrsinites*). The most common seral tree species is western larch. The understory is depauperate of species and cover is very low. Species are shade-loving and include prince's plume, queencup beadlily, western rattlesnake plantain, and twinflower with a thick cover of moss. Soils associated with this ecological site are very deep ash-capped soils over rocky subsoils. Ash thickness on these soils varies from 20cm to 50+ cm and provides a boost in water-holding capacity to these sites. These soils generally belong in the Andic Eutrudept soil subgroup, but also in the Typic Hapludands and Andic Dystrudepts. Diagnostic features include an ochric epipedon, andic soil properties and a cambic horizon. There is a thin organic layer present in these soils, generally less than 5 cm thick.

Associated sites

F043AX952MT	<p>Lower Subalpine Cool Moist Coniferous subalpine fir-Engelmann spruce/Rocky Mountain maple-thinleaf huckleberry/thimbleberry</p> <p>The 43A Lower Subalpine Frigid Coniferous ecological site is associated with the 43A Subalpine Coniferous Cool Moist Ashy Very Deep ecological site. The 43A Lower Subalpine Coniferous Cool Moist Ashy Very Deep is found in cool, moister mid-elevations that span the lower subalpine to subalpine. This ecological site is found on back, foot and toeslope positions, on glacial valley wall and moraine landforms, on all slopes, at elevations ranging 1,000 to 2,100 meters (3,280-6,890 feet). The 43A Lower Subalpine Frigid Coniferous ecological site is associated with the 43A Subalpine Coniferous Cool Moist Ashy Very Deep ecological site. The 43A Lower Subalpine Coniferous Cool Moist Ashy Very Deep has soils associated with this Ecological Site that are very deep, well drained or somewhat excessively drained and have subsoils with abundant rock fragments. The parent material is volcanic ash over glacial till from metasedimentary rock. In Soil Taxonomy, these soils classify primarily as Inceptisols soil order and more specifically as the Andic Haplocryepts taxonomic subgroup. The 43A Lower Subalpine Frigid Coniferous ecological site is associated with the 43A Subalpine Coniferous Cool Moist Ashy Very Deep ecological site. The 43A Lower Subalpine Coniferous Cool Moist Ashy Very Deep has a reference vegetation community of Subalpine fir-Engelmann spruce overstory and an understory of Rocky Mountain maple, thinleaf huckleberry, thimbleberry, wild Sarsaparilla, threeleaf foamflower and queencup bead lily.</p>
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Thuja plicata</i> (2) <i>Tsuga heterophylla</i>
Shrub	(1) <i>Chimaphila umbellata</i>
Herbaceous	(1) <i>Clintonia uniflora</i> (2) <i>Goodyera oblongifolia</i>

Physiographic features

The lower subalpine frigid coniferous site is found west of the Continental Divide in moist areas within this maritime climate in glacial valley wall and lateral moraine landforms on back and foot slope positions, and on all aspects, spanning elevations from 1,000 to 1,500 feet. In Montana, in general, the site is found at elevations ranging from 2,000-5,000 feet. The annual precipitation generally is above 32" (USFS H.T. Guide, 1977).

Table 2. Representative physiographic features

Landforms	(1) Mountains > Lateral moraine (2) Mountains > Glacial-valley wall (3) Mountains > Alluvial fan (4) Mountains > Outwash terrace
Elevation	3,280–4,921 ft
Slope	5–35%
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

This ecological site is found in the frigid soil temperature regime and the udic soil moisture regime. The soils that support this native plant community occur in the frigid soil temperature regime (average annual temperature less

than 8 degrees C, with more than 5 degrees C summer-winter fluctuation). An udic soil moisture regime denotes that the rooting zone is usually moist throughout the winter and the majority of summer. This site is found on the west side of the Continental Divide and has more maritime weather influences.

Table 3. Representative climatic features

Frost-free period (characteristic range)	17-57 days
Freeze-free period (characteristic range)	76-117 days
Precipitation total (characteristic range)	20-26 in
Frost-free period (actual range)	6-68 days
Freeze-free period (actual range)	66-127 days
Precipitation total (actual range)	20-28 in
Frost-free period (average)	37 days
Freeze-free period (average)	97 days
Precipitation total (average)	23 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) POLEBRIDGE 1 N [USC00246618], Essex, MT
- (2) POLEBRIDGE [USC00246615], Essex, MT
- (3) WEST GLACIER [USC00248809], Kalispell, MT

Influencing water features

This ecological site is not influenced by wetland or riparian water features but may be found on stream terraces or adjacent to wetland and riparian areas.

Soil features

Soils associated with this ecological site are very deep ash-capped soils over rocky subsoils. Ash thickness on these soils varies from 20cm to 50+ cm and provides a boost in water-holding capacity to these sites. These soils generally belong in the Andic Eutrudept soil subgroup, but also in the Typic Hapludands and Andic Dystrudepts. Diagnostic features include an ochric epipedon, andic soil properties and a cambic horizon. There is a thin organic layer present in these soils, generally less than 5 cm thick.

“Mount Mazama (Crater Lake, Oregon) violently erupted around 6,700 years ago. The massive plume of volcanic ash from the eruption drifted in a northwest direction through northwest Montana. Deposition was widespread throughout western Montana, but is greatest in the northwest corner of the state. The ash likely fell uniformly across the landscape and was then re-distributed by wind and water erosion. The resulting distribution, given the dominate winds from the south and southwest, favors thicker deposits of ash on slopes with north-facing aspects. Little ash is usually found on south-facing slopes except at the higher elevations in cirque basins.

Volcanic ash has a large impact on overstory and understory plant productivity due to the increase in water-holding capacity that it adds to the soil. The ash is typically found as a surface mantle overlying whatever existing soil or parent material it was deposited on top of. The ash mantles exhibit varying amounts of mixing with the underlying material, but tend to lack the rock fragments commonly found in the sub-soils of the park. Ashy soil layers generally tend to have brighter colors than the underlying sub-soils and have a soft and very friable consistency.” (J. Skovlin, personal communication, 2015). (Soil Survey Staff, 2015). For more information on soil taxonomy, please follow this link:

http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/class/?cid=nrcs142p2_053580

CORRELATED SOIL SERIES & TAXONOMIC CLASS NAME

Backroad Loamy-skeletal, mixed, superactive, frigid Andic Dystrudepts

Pasturecreek Loamy-skeletal, isotic, frigid Andic Eutrudepts

Sunroad Coarse-loamy, isotic, frigid Andic Eutrudepts

Typic Hapludands Medial over sandy or sandy-skeletal, amorphic over isotic, frigid Typic Hapludands



Figure 7. Soils for this ecological site.

Table 4. Representative soil features

Parent material	(1) Till–metasedimentary rock (2) Colluvium–metasedimentary rock (3) Alluvium–metasedimentary rock
Surface texture	(1) Gravelly, ashy loam (2) Ashy loam
Family particle size	(1) Loamy-skeletal

Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderate
Soil depth	60–100 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (2.7-7.2in)	Not specified
Soil reaction (1:1 water) (4.5-6.5in)	Not specified

Ecological dynamics

Ecological Dynamics of the Site

This site is found west of the Continental Divide in moderately cool and moist mid-elevations sites. Within Glacier National Park (NP), this ecological site has its greatest expression on the eastern side of Lake McDonald. While primary data was collected in Glacier NP, this habitat type also spans into the adjacent US Forest Service (USFS) land Flathead National Forest (NF), and in the Kootenai NF.

Management

Various management strategies can be employed for this ecological site, depending upon the ownership of the particular land and which value is prioritized. The management of the forest determines the composition of the stand and the amount of fuel loading. A stand will be managed differently and look differently if it is managed for timber or ecological services like water quality and quantity, old growth, or endangered species. If a stand is managed for timber, then it may be missing certain attributes necessary for lynx habitat. If a stand is managed for lynx habitat, it may have increased fuels and therefore an increased risk of wildfires.

The USFS Habitat Type Guide (1973) states that the basal area on the western side of the Continental Divide for western hemlock/queencup beadlily is 267+/-55 ft² per acre. The fifty-year site index for western white pine is 62, western larch is 80, spruce is 77, and grand fir is 50. Basal area for western redcedar/queencup beadlily is 305+/-96 ft² per acre and the site index for Douglas-fir is 66, western larch is 63, PICEA is 72+/-14, grand fir is 61, and subalpine fir is 74.

Timber production on these sites is very high, particularly in the seral phases of this ecological site.

Each national forest has a specific management plan. The management plan for the Flathead N.F. also has an Appendix B that gives specific management guidelines for habitat types (which relate to our forested ecological sites) found on the forest in relation to current and historic data on forest conditions (Flathead N.F. Plan, 2001 and Appendix B). Another guiding USFS document is the Green et al. document (2005) which defines "Old Growth" forest for the northern Rocky Mountains. This document provides an ecologically-based classification of old growth based on forest stand attributes including numbers of large trees, snags, downed logs, structural canopy layers, canopy cover, age, and basal area. While this document finds that the bulk of the pre-settlement upland old growth in the northern Rockies was in the lower elevation, ground-fire maintained ponderosa pine/western larch/Douglas-fir types (Losensky, 1992), it does not mean that other types were not common or not important. This could apply to some of the areas of this ecological site.

State 1.0

Western White Pine (*Pinus monticola*)(Western redcedar (*Thuja plicata*)-Western hemlock (*Tsuga heterophylla*))/pipsissewa (*Chimaphila umbellata*)-queencup beadlily (*Clintonia uniflora*).

Historically western white pine would have been within Flathead County, which encompasses the Flathead N.F. and in lower elevations, west of the Continental Divide in Glacier NP. The historic extent of western white pine in Glacier National Park was primarily along the western border. Originally, western white pine covered five million acres in the Inland Northwest. Western white pine is incredibly productive for timber with a very high growth rate, tall and deep-rooted, and competes best on highly variable, high resource sites. As well, it is tolerant to the native root rot diseases and other native forest pests. Western white pine is susceptible to Armillaria root disease only when young, and to mountain pine beetle largely at advanced ages (over 140 years). It also has the capability to thrive in

a wide variety of sites and environments, which means it has high ecological flexibility. It is a long-living seral species that tolerated intense timber harvesting practices and severe fire disturbance by its ability to regenerate heavily on mineral soil and full sunlight. Fire greatly influences the composition, structure, and function of vegetation across the landscape. Historically, it was mixed severity fire between severe stand replacement fires. Western larch and western white pine are long-lived, fire-adapted, shade-intolerant tree species that historically thrived. Also present in significant amounts particularly in young stands, but declined through time due to effects of insects and pathogens, were the shorter-lived, shade-intolerant, fire-adapted tree species such as Douglas-fir and lodgepole pine. Shade-tolerant, fire-intolerant tree species such as western cedar, western hemlock, grand fir, Engelmann spruce, and subalpine fir were present, but rarely survived long enough to dominate stands except in areas where the interval between fires was unusually long and where root disease was not severe.

Prior to the 20th century, western white pine was a major component in forested ecosystems of the inland northwest U.S. It has been greatly reduced in distribution and abundance by white pine blister rust, mountain pine beetles, and anthropogenic fire exclusion (Tomback and Achuff, 2010). Western white pine has been replaced by Douglas-fir, grand fir, and western hemlock. Douglas-fir and grand fir are susceptible to a greater variety of insect and disease problems and hemlock is more sensitive to drought and decay. More stands have also progressed to the climax species-dominated phase, which previously were rarely achieved due to the fire rotations and susceptibility of these species to disease and forest pests. In a study of pathogens and insects effects on forests within the Inland Empire found that, excluding fire, there were more than 90% of sample stands changed to a different cover type, structure stage, or both during a 40-year period that was coincident with the blister rust epidemic and fire suppression policy. Root pathogens, white pine blister rust, and bark beetle were the cause of most changes, and this accelerated succession of western white pine, ponderosa pine, and lodgepole pine to later successional, more shade-tolerant species. Structure was reduced in stand density or prevented canopy closure. Grand fir, Douglas-fir, and subalpine fir were the predominant cover types at the end of the period, and were highly susceptible to root diseases, bark beetles, fire, and drought. It is estimated that there will be continuation of this trend occurring in low-density mature stands and younger pole-sized stands that result from root disease and bark beetle-caused mortality (Byler and Hagel, 2000). These stands also are less productive in terms of timber. They are dominated by species with high nutrient demands, and therefore nutrient storage and cycling rates are increasingly depressed. This will likely lead to ever-increasing stress and destabilization by pests and diseases. Drought can further exacerbate the situation by stressing trees.

The Inland Empire Tree Improvement Cooperative and the USFS have a breeding program for blister-resistant western white pine. Approximately 5 percent of the original acre range was re-planted with rust-resistant stock. Currently, the modified stock shows about 60 percent resistance to blister rust. A study modeling the effects of climate change found that warming temperatures would favor increased abundance of western white pine over existing climax and shade-tolerant species in Glacier NP, mainly because warmer conditions potentiate fire dynamics, including increased wildfire frequency and extent, which facilitates regeneration (Loehman, et al., 2011).

State 2.0

Western redcedar (*Thuja plicata*)-Western hemlock (*Tsuga heterophylla*)/pipsissewa (*Chimaphila umbellata*)-queencup beadlily (*Clintonia uniflora*)-western rattlesnake plantain (*Goodyera oblongifolia*)-twinflower (*Linnaea borealis*)/moss.

State 2 is different than State 1 in that western white pine no longer plays a significant role in the seral communities. It has been dramatically reduced in numbers and area by the epidemics of white pine blister rust and western spruce budworm, and by dramatic fire suppression. Therefore, climax species have been able to fill the seral role that western white pine once held. As well, more forests are progressing to the climax or Reference Phase than historically, when most forests were in the fire-maintained western white pine-dominated seral phase. State 2 forests are now dominated by the shade-tolerant climax species western redcedar and western hemlock. While there is a tremendous effort to bolster the numbers of western white pine, it currently covers only 5 percent of its historic range.

This ecological site is described as having moderately cool and moist site conditions. The Reference state is dominated by western redcedar and western hemlock, both of which are shade-tolerant climax conifers that grow in similar environments. Western redcedar has a larger geographic extent in Montana, but western hemlock usually is capable of attaining dominance over western redcedar and other species at climax because it is better able to

reproduce under a dense forest canopy. Western redcedar is able to maintain itself indefinitely as a minor climax species because of its shade tolerance, longevity (often 600-1,000 years), and apparent ability to regenerate vegetatively (USFS H.T. Guide, 1973). Within Glacier NP, these species are co-dominant in nearly all of the sites visited. The seral successional stages have very diverse overstory tree composition and can be very productive in terms of basal area. Douglas-fir, western larch and, to a lesser extent, spruce are often dominants in seral stands with lodgepole, western white pine, and paper birch as minor components. Grand fir and subalpine fir can be either minor seral or climax components. Western redcedar and western hemlock will regenerate after disturbance along with seral species, and it will take centuries for these species to gain dominance in the overstory over the seral species. The early successional phase can be dominated by fireweed (*Chamerion angustifolium*). The understory in seral successional phases have moisture-loving forbs or shrubs including Scouler's willow, thimbleberry, serviceberry, rocky mountain maple, thinleaf huckleberry, and snowbush ceanothus. The historic fire regime of these forests is one of low fire frequency, but fire severity can be highly variable. It can be low due to the most common moist conditions, but can be severe during times of drought. Fire return intervals range from 50 to greater than 200 years, but include mixed severity fires on 50-85 year intervals, as well as stand replacement fires on 150-250 year intervals. Western redcedar can thrive for centuries on this ecological site without disturbance. The Northern Rocky Mountain mesic montane mixed-conifer forest-cedar groves are in fire regime group 5 and had a fire interval of 334 years, with 87 percent of fires classified as of replacement severity and 13 percent of fires classed as mixed severity and none as low severity (USDA, USFS, FEIS, Fire Regime). Fuel loadings for this ecological site can be very high due to deadfall and natural thinning of small and medium-sized branches. In early and intermediate successional phases, the understory can have high cover adding to fuel loadings. Due to the generally moist conditions, fire return intervals can be long. In general, the variability in fire regime and the high diversity of tree species present in most stands, except the Reference, allow this ecological site to form a diverse mosaic landscape with varying dominance or mixes of seral species.

The general fire succession process is that after stand replacement fires, the community reverts to an herbaceous one, then to shrubland. If fire is reoccurring in this phase, then the phase is maintained for a long time. The herbaceous community can be dominated by the disturbance-loving fireweed, beargrass, or numerous other species, depending upon the seedbank at the site and beyond. Duration of the herbaceous or shrubland phase is also dependent upon the availability of tree seed. If serotinous lodgepole pine seeds are available, then the site will become dominated by it and a lodgepole pine stand will develop for about 10-25 years (Habeck, 1968). After that time, other species become established including western larch and other conifers. If serotinous lodgepole pine seeds are not present, then the seedlings are a very diverse mixture of conifers. These seedlings form a thick carpet on the site shared with shrub species such as Scouler's willow, white spirea, thinleaf huckleberry, thimbleberry, and Oregon boxwood. Forbs present include ferns, beargrass and fireweed. Moss cover can be variable. If fire does not occur, the seedlings will grow to saplings and then pole-sized trees of diverse seral species. Low to moderate fires in this stage would favor fire-tolerant seral species over western redcedar, grand fir, or western hemlock, which are less fire-resistant. Severe fires will return these to the herbaceous or shrubland phase. In the pole-sized phase, seral species are abundant and western redcedar and western hemlock are just becoming established and usually have low cover (3-15 percent of the stand). Without further disturbance, this phase will continue to the maturing forest in which western redcedar and western hemlock become more evident in the stand and eventually have higher cover than the seral tree species. Western larch may survive severe fires in the maturing or mature phases. These trees would then provide seed for the stand initiation phase after a fire. As well, after frequent low to moderate fires in the mature phase, a relict western larch stand could occur. Reference stands in which only western redcedar and western hemlock occur can be rare, as seral species are long-lived and fire occurs frequently enough that stands seldom develop beyond the mature phase. Along the shores of Lake McDonald in Glacier NP, there were abundant stands that were in the Reference phase. The Robert fire in 2003 heavily impacted some areas on the west side of Lake McDonald, but other areas close to the shore were not affected. Reference stands may withstand low fires that thin the stand, but moderate or severe fires would return the site to the herbaceous or shrubland phase. Significant fires that have occurred on the west side of the Continental Divide that have affected this ecological site are the Robert Fire in 2003 that burned 54,191 acres, the Moose Fire in 2001 that burned 66,688 acres, and the Middle Fork Complex Fire in 2003 that burned 11,996 acres. There were historic fires within the area of this ecological site that burned significant portions in 1735 and another portion in 1926 (NPS Stand Age spatial layer).

Both western hemlock and western redcedar are subjected to a variety of diseases and insect pests including Armillaria root disease, Annosus root disease, pouch fungus, red belt fungus, pini rot, metallic wood borers, and roundheaded borers. Western redcedar also is susceptible to cedar laminated butt rot, cedar brown pocket rot, and cedar bark beetles. Western hemlock is also susceptible to Indian paint fungus.

A good tool to use to discern the levels of insects and diseases, the damage patterns, and whether these are at endemic or epidemic levels is aerial photography. These maps capture only moments in time, and infestations grow and move from location to location following their preferred habitat, so repeated photography can be necessary. Specifically for the northern region, the USFS Stand Health map shows, via many very large polygons throughout the area, that the major impact is defoliation by western spruce budworm. The defoliation was categorized as mostly of low severity (equal to or less than 50% defoliation) and some of high severity (with greater than 50% defoliation) on *Abies* species, and the damage is contiguous or nearly continuous. The forest type was categorized as W. Fir-Spruce. There also was defoliation by western spruce budworm on Douglas-fir, but to a much lesser degree. Larch casebearer, a defoliator of western larch, and generalized needlecast disease of western larch also was found, to a much lesser degree. Scattered small polygons showing damage were found throughout the region, including mortality from mountain pine beetle on lodgepole pines, Douglas-fir beetle on Douglas-fir, spruce beetle on Engelmann spruce, fir engravers and woolly adelgid on *Abies* species, and general *Abies* species mortality. These would affect the seral tree species of this ecological site and field notes corroborate these findings.

Community Phase 2.1: Reference

Western redcedar-western hemlock/prince's plume-queencup beadlily-western rattlesnake plantain-twinflower/moss
Structure: Multistory with small gap dynamics

The Reference Community is dominated by western cedar and western hemlock, with seral tree species constrained to less than 3% of the overstory canopy. The ground cover consists predominantly of duff with fairly high cover of moss and trace cover of embedded litter and stones. The vegetation structure is that of very tall trees from 500-1,100 inches tall of western larch, western redcedar and western hemlock. The understory is multistoried though fairly sparse. Species occurring with the highest frequency of occurrence include thinleaf huckleberry, darkwoods violet, twinflower, western rattlesnake plantain, queencup beadlily and prince's plume (15 sites of canopy cover data). Foliar cover at six sites indicate that the foliar cover is fairly high (50.8%) and the ground cover is primarily litter, which includes woody litter, litter or duff (total is 74%, duff is 63%) and moss (30%). The tallest understory layer is 20-40 inches tall and can include common ladyfern (*Athyrium filix-femina*), common snowberry (*Symphoricarpos albus*), white spirea (*Spiraea betulifolia*), and thinleaf huckleberry (*Vaccinium membranaceum*). The lowest layer is less than 10 inches tall and can include fireweed (*Chamerion angustifolium*), pipsissewa (*Chimaphila umbellata*), queencup beadlily (*Clintonia uniflora*), western rattlesnake plantain (*Goodyera oblongifolia*), twinflower (*Linnaea borealis*), threeleaf foamflower (*Tiarella trifoliata*), and Oregon boxleaf (*Paxistima myrsinites*). The most common seral tree species is western larch. The understory is depauperate of species and cover is very low. Species are shade-loving and include prince's plume, queencup beadlily, western rattlesnake plantain, and twinflower with a thick cover of moss.

Community Phase Pathway 2.1A

This pathway represents a major stand-replacement fire disturbance such as a high-intensity fire, large scale wind event, or major insect infestation.

Forest overstory summarization table.

FOREST OVERSTORY

Forest canopy Canopy cover Average= 60%, Range= 45-90%

Average basal area Total 260-380 ft²/acre

Site Index at 100 yrs.: THPL (64-86); TSHE (65-91)

Community Phase 2.2: Stand Initiation. Regen: Lodgepole pine-western larch (mixed seral species)/

Scouler's willow-white spirea-thinleaf huckleberry/thimbleberry-Oregon boxwood/fireweed-beargrass/moss.

Post-fire 1-5 years

Structure: Initially this post fire disturbance community is dominated by herbaceous and shrub species most commonly fireweed, species with a resident seedbank or disturbance loving species.

Structure: Continuous cover of regeneration-single story mixed tree species.

Post-fire 5-50 years

Structure: This is a forest in the stand initiation phase, possibly with scattered remnant mature trees; the composition of the seedlings depends on the natural seed sources available. Habeck found that in the vicinity of

Lake McDonald in Glacier N.P., the dominant seral tree species is Lodgepole pine for 10-25 years post-fire. Afterwards, Western Larch will co-dominant from 25-50 years post-fire with other seral tree species at lower cover. Throughout the entire area of this ecological site, the regeneration will probably be a mixture. Overstory canopy cover is generally less than 10%, but the regeneration tree cover is very high forming a thick carpet. It is a mixture of species including: Lodgepole pine, Western larch, subalpine fir, Paper birch, Engelmann spruce, Western white pine, Black cottonwood, Quaking aspen, Douglas fir, Western redcedar and Western hemlock. The understory is a diverse mixture of herbaceous and shrub species including tall willow species, particularly Scouler's willow, medium statured shrubs white spirea and thinleaf huckleberry, the low statured shrubs thimbleberry and Oregon boxwood. Herbaceous species include: fireweed, beargrass. Moss cover is variable.

Community Phase Pathway 2.2A

This pathway represents growth over time with no further significant disturbance. The areas of regeneration pass through the typical stand phases-competitive exclusion, maturation, understory reinitiating-until they resemble the old-growth structure of the reference community.

Community Phase 2.3: Intermediate Aged Forest.

Lodgepole pine-Douglas fir-Engelmann spruce-western larch-paper birch-subalpine fir-western white pine (western redcedar-western hemlock)/white spirea-snowberry-thinleaf huckleberry/thimbleberry/prince's plume-queencup beadlily-twinflower-beargrass/moss.

This community phase is dominated by seral tree species that have matured to pole size and are in the competitive exclusion phase of forest succession. Overstory tree canopy is dense and competition for resources is very high. Canopy cover averages 50%. This community is incredibly diverse in tree species including: Lodgepole pine, Douglas fir, Engelmann spruce, Western larch, Paper birch, subalpine fir, Western white pine, Western redcedar and Western hemlock. The overstory canopy of Western redcedar and Western hemlock is less than 3% as they are just beginning to become established. The understory can have high cover of the medium sized shrubs white spirea, snowberry, Oregon boxleaf, common snowberry and thinleaf huckleberry. The short statured thimbleberry can have high cover. The herbaceous layer is diverse, with medium statured beargrass occurring frequently and sometime in high cover. Other herbaceous species include the short statured prince's plume, queencup beadlily, trailplant and twinflower.

Figure 7. Plant Community 2.3 Intermediate Aged Forest, Dense Thick Pole Sized Trees.

Community Phase Pathway 2.3A

This pathway represents continued growth over time with no further major disturbance.

Community Phase Pathway 2.3B

This pathway represents a major stand-replacement fire disturbance such as a high-intensity fire, large scale wind event or major insect infestation.

Community Phase 2.4: Maturing Forest.

Western redcedar-western hemlock-subalpine fir-lodgepole pine-Engelmann spruce-western white pine/thinleaf huckleberry-snowberry-white spirea/wild sarsaparilla-heartleaf arnica-queencup beadlily-twinflower-beargrass/moss. Structure:

This community is a maturing forest with vertical differentiation in the overstory tree canopy. Canopy cover averages 60%. This community has diverse tree species with Western redcedar and Western hemlock ranging 3-15% each and other seral tree species about equally distributed. These species include: Subalpine fir, Lodgepole pine, Engelmann spruce, Western white pine. The understory has patchy medium sized shrubs including: thinleaf huckleberry, snowberry, white spirea. There is a diverse understory of herbaceous species including: wild sarsaparilla, heartleaf arnica, queencup beadlily, twinflower and beargrass. There can be high cover of moss.

Community Phase Pathway 2.4A

This pathway represents continued growth over time with no further major disturbance.

Community Phase Pathway 2.4B

This pathway represents a major stand-replacement fire disturbance, such as a major insect outbreak, or major fire event which leads to the stand initiation phase of forest development.

Community Phase 2.5: Mature Forest.

Western redcedar-western hemlock (remnant seral species)/thinleaf huckleberry/threeleaf foamflower-prince's

plume-queencup beadlily-western rattlesnake plantain/moss.

Structure: Mature forest with vertical differentiation in the stand. Overstory is dominated by Western redcedar and Western hemlock although seral tree species are present and can have up to 15% cover each. Overstory canopy cover ranges 50-80%. Western larch is the most common seral species but others include: Grand fir, subalpine fir, Paper birch, Lodgepole pine, Western white pine, and Douglas fir. The understory is diverse but generally has low overall cover. Thinleaf huckleberry occurs in clumps and queencup beadlily and western rattlesnake plantain are common.

Figure 9. Plant Community 2.5 Mature forest with some small gap dynamics, remnant seral tree species and western redcedar and western hemlock dominant.

Community Phase Pathway 2.5A

This pathway represents no further major disturbance. Continued growth over time, as well as ongoing mortality, leads to continued vertical diversification. The community begins to resemble the structure of the reference community, with small pockets of regeneration and a more diversified understory.

Community Phase Pathway 2.5B


This pathway represents a major stand-replacement fire disturbance leading to the stand initiation phase of forest development.

State 3.0


Another disease affecting this ecological site is root rot. While Douglas-fir, grand fir, and subalpine fir are most susceptible, western redcedar and western hemlock can be affected as well. Armillaria root disease is the most common root disease fungus in this region, and is especially prevalent west of the Continental Divide. It may be difficult to detect until it has killed enough trees to create large root disease pockets or centers, ranging in size from a fraction of an acre to hundreds of acres. The root disease spreads from an affected tree to its surrounding neighbors through root contact. The root disease effects the tree species most susceptible first, leaving less susceptible tree species that mask its presence. When root rot is severe, the pocket has abundant regeneration or dense brush growth in the center. Western redcedar is moderately resistant to Armillaria root rot in Idaho and Montana. The common disease expression is some mortality in saplings, and residuals of partial harvests often develop severe infections but are very slow to die (Hagel, 2010). There has been a link determined between parent material and susceptibility to root disease (Kimsey et al., 2012). Metasedimentary parent material is thought to increase the risk of root disease. Glacier National Park is dominated by metasedimentary parent material and may be more at risk than other areas to root disease (Kimsey et al., 2012). If a stand sustains very high levels of root disease mortality, then a coniferous stand could cross a threshold and become a shrubland, once all conifers are gone (Kimsey et al., 2012). Management tactics include to identify the type of Armillaria root disease, and manage for pines and larch. Pre-commercial thinning may improve growth and survival of pines and larch. Avoid harvests that leave susceptible species (usually Douglas-fir or true firs) as crop trees (Hagel, 2010).

State and transition model


Ecosystem states




State 1 submodel, plant communities




Communities 1 and 5 (additional pathways)




State 2 submodel, plant communities



Communities 1, 5 and 2 (additional pathways)



State 3 submodel, plant communities



State 1

Historical Reference State Western white pine (western hemlock-western red cedar)/pipsissewa-Clintonia

Historically western white pine would have been within Flathead County, which encompasses the Flathead N.F. and in lower elevations, west of the Continental Divide in Glacier NP. The historic extent of western white pine in Glacier National Park was primarily along the western border. Originally, western white pine covered five million acres in the Inland Northwest. Western white pine is incredibly productive for timber with a very high growth rate, tall and deep-rooted, and competes best on highly variable, high resource sites. As well, it is tolerant to the native root rot diseases and other native forest pests. Western white pine is susceptible to Armillaria root disease only when young, and to mountain pine beetle largely at advanced ages (over 140 years). It also has the capability to thrive in a wide variety of sites and environments, which means it has high ecological flexibility. It is a long-living seral species that tolerated intense timber harvesting practices and severe fire disturbance by its ability to regenerate heavily on mineral soil and full sunlight. Fire greatly influences the composition, structure, and function of vegetation across the landscape. Historically, it was mixed severity fire between severe stand replacement fires. Western larch and western white pine are long-lived, fire-adapted, shade-intolerant tree species that historically thrived. Also present in significant amounts particularly in young stands, but declined through time due to effects of insects and pathogens, were the shorter-lived, shade-intolerant, fire-adapted tree species such as Douglas-fir and lodgepole pine. Shade-tolerant, fire-intolerant tree species such as western cedar, western hemlock, grand fir, Engelmann spruce, and subalpine fir were present, but rarely survived long enough to dominate stands except in areas where the interval between fires was unusually long and where root disease was not severe.

Community 1.1

Reference Community Subalpine fir- Engelmann spruce overstory. Minor western white pine-western larch-grand fir

Reference phase of multi-storied forest canopy dominated by western redcedar and western hemlock.

Community 1.2

Western white pine-western larch- (lodgepole pine) seedlings

Post fire disturbance community of herb and shrub species.

Community 1.3

western white pine-western larch-(subalpine fir-grand fir-Engelmann spruce)

Intermediate aged forest, dense thick pole sized trees.

Community 1.4

Western white pine- (western hemlock-western red cedar grand fir)

Maturing forest phase of seral tree species and western redcedar and western hemlock.

Community 1.5

Western white pine- (western hemlock-western red cedar-grand fir

Mature forest with some small gap dynamics, remnant seral trees species and western red cedar and western hemlock dominant.

Pathway 1.1A

Community 1.1 to 1.2

A major stand-replacement disturbance, such as a major insect outbreak or major fire event, which leads to the stand initiation phase of forest development.

Pathway 1.2A

Community 1.2 to 1.3

Continued growth over time with no further major disturbance to dense single story pole sized stand.

Pathway 1.3A

Community 1.3 to 1.4

Continued growth over time with no further major disturbance to mature stand with all size classes.

Pathway 1.3B

Community 1.3 to 1.5

Continued growth over time with no further major disturbance with patches of regeneration.

Pathway 1.4A

Community 1.4 to 1.1

Continued growth over time with no further major disturbance with patches of regeneration.

Pathway 1.5A

Community 1.5 to 1.1

Continued growth over time with no further major disturbance with patches of regeneration.

State 2

Current Reference State

Western redcedar (*Thuja plicata*)-Western hemlock (*Tsuga heterophylla*)/pipsissewa (*Chimaphila umbellata*)-queencup beadlily (*Clintonia uniflora*)-western rattlesnake plantain (*Goodyera oblongifolia*)-twinflower (*Linnea borealis*)/moss. State 2 is different than State 1 in that western white pine no longer plays a significant role in the seral communities. It has been dramatically reduced in numbers and area by the epidemics of white pine blister rust and western spruce budworm, and by dramatic fire suppression. Therefore, climax species have been able to fill the seral role that western white pine once held. As well, more forests are progressing to the climax or Reference Phase than historically, when most forests were in the fire-maintained western white pine-dominated seral phase. State 2 forests are now dominated by the shade-tolerant climax species western redcedar and western hemlock. While there is a tremendous effort to bolster the numbers of western white pine, it currently covers only 5 percent of its historic range. This ecological site is described as having moderately cool and moist site conditions. The Reference state is dominated by western redcedar and western hemlock, both of which are shade-tolerant climax conifers that grow in similar environments. Western redcedar has a larger geographic extent in Montana, but western hemlock usually is capable of attaining dominance over western redcedar and other species at climax because it is better able to reproduce under a dense forest canopy. Western redcedar is able to maintain itself indefinitely as a minor climax species because of its shade tolerance, longevity (often 600-1,000 years), and apparent ability to regenerate vegetatively (USFS H.T. Guide, 1973). Within Glacier NP, these species are co-dominant in nearly all of the sites visited. The seral successional stages have very diverse overstory tree composition and can be very productive in terms of basal area. Douglas-fir, western larch and, to a lesser extent, spruce are often dominants in seral stands with lodgepole, western white pine, and paper birch as minor components. Grand fir and subalpine fir can be either minor seral or climax components. Western redcedar and western hemlock will regenerate after disturbance along with seral species, and it will take centuries for these species to gain dominance in the overstory over the seral species. The early successional phase can be dominated by fireweed (*Chamerion angustifolium*). The understory in seral successional phases have moisture-loving forbs or shrubs including Scouler's willow, thimbleberry, serviceberry, rocky mountain maple, thinleaf huckleberry, and snowbush ceanothus. The historic fire regime of these forests is one of low fire frequency, but fire severity can be highly variable. It can be low due to the most common moist conditions, but can be severe during times of drought. Fire return intervals range from 50 to greater than 200 years, but include mixed severity fires on 50-85 year intervals, as well as stand replacement fires on 150-250 year intervals. Western redcedar can thrive for centuries on this ecological site without disturbance. The Northern Rocky Mountain mesic montane mixed-conifer forest-cedar groves are in fire regime group 5 and had a fire interval of 334 years, with 87 percent of fires classified as of replacement severity and 13 percent of fires classed as mixed severity and none as low severity (USDA, USFS,

FEIS, Fire Regime). Fuel loadings for this ecological site can be very high due to deadfall and natural thinning of small and medium-sized branches. In early and intermediate successional phases, the understory can have high cover adding to fuel loadings. Due to the generally moist conditions, fire return intervals can be long. In general, the variability in fire regime and the high diversity of tree species present in most stands, except the Reference, allow this ecological site to form a diverse mosaic landscape with varying dominance or mixes of seral species.

Community 2.1

Reference Community Western redcedar-western hemlock/prince's plume-queencup beadlily-western rattlesnake plantain-twinflower/moss



Figure 8. Vegetation community in the reference phase.



Figure 9. Vegetation at a reference site, noting multi-storied understory.

The Reference Community is dominated by western cedar and western hemlock, with seral tree species constrained to less than 3% of the overstory canopy. The ground cover consists predominantly of duff with fairly high cover of moss and trace cover of embedded litter and stones. The vegetation structure is that of very tall trees from 500-1,100 inches tall of western larch, western redcedar and western hemlock. The understory is multistoried though fairly sparse. Species occurring with the highest frequency of occurrence include thinleaf huckleberry, darkwoods violet, twinflower, western rattlesnake plantain, queencup beadlily and prince's plume (15 sites of canopy cover data). Foliar cover at six sites indicate that the foliar cover is fairly high (50.8%) and the ground cover is primarily litter, which includes woody litter, litter or duff (total is 74%, duff is 63%) and moss (30%). The tallest understory layer is 20-40 inches tall and can include common ladyfern (*Athyrium filix-femina*), common snowberry (*Symphoricarpos albus*), white spirea (*Spiraea betulifolia*), and thinleaf huckleberry (*Vaccinium membranaceum*). The lowest layer is less than 10 inches tall and can include fireweed (*Chamerion angustifolium*), pipsissewa (*Chimaphila umbellata*), queencup beadlily (*Clintonia uniflora*), western rattlesnake plantain (*Goodyera oblongifolia*), twinflower (*Linnaea borealis*), threeleaf foamflower (*Tiarella trifoliata*), and Oregon boxleaf (*Paxistima myrsinites*). The most common seral tree species is western larch. The understory is depauperate of species and cover is very low. Species are shade-loving and include prince's plume, queencup beadlily, western rattlesnake plantain, and twinflower with a thick cover of moss.

Community 2.2

Stand Initiation Regen: Lodgepole pine-western larch (mixed seral species)/Scouler's willow-white spirea-thinleaf huckleberry/thimbleberry-Oregon boxwood/fireweed-beargrass/moss



Figure 10. Plant Community 2.2 Post Fire Disturbance Community of Herbaceous and Shrub Species.

Structure: This is a forest in the stand initiation phase, possibly with scattered remnant mature trees; the composition of the seedlings depends on the natural seed sources available. Habeck found that in the vicinity of Lake McDonald in Glacier N.P., the dominant seral tree species is Lodgepole pine for 10-25 years post-fire. Afterwards, Western Larch will co-dominant from 25-50 years post-fire with other seral tree species at lower cover. Throughout the entire area of this ecological site, the regeneration will probably be a mixture. Overstory canopy cover is generally less than 10%, but the regeneration tree cover is very high forming a thick carpet. It is a mixture of species including: Lodgepole pine, Western larch, subalpine fir, Paper birch, Engelmann spruce, Western white pine, Black cottonwood, Quaking aspen, Douglas fir, Western redcedar and Western hemlock. The understory is a diverse mixture of herbaceous and shrub species including tall willow species, particularly Scouler's willow, medium statured shrubs white spirea and thinleaf huckleberry, the low statured shrubs thimbleberry and Oregon boxwood. Herbaceous species include: fireweed, beargrass. Moss cover is variable.

Community 2.3

Intermediate Aged Forest Lodgepole pine-Douglas fir-Engelmann spruce-western larch-paper birch-subalpine fir-western white pine (western redcedar-western hemlock)/white spirea-snowberry-thinleaf huckleberry/thimbleberry/prince's plume-queencup beadlily



Figure 11. Plant Community 2.3 Intermediate Aged Forest, Dense Thick Pole Sized Trees.

This community phase is dominated by seral tree species that have matured to pole size and are in the competitive exclusion phase of forest succession. Overstory tree canopy is dense and competition for resources is very high. Canopy cover averages 50%. This community is incredibly diverse in tree species including: Lodgepole pine, Douglas fir, Engelmann spruce, Western larch, Paper birch, subalpine fir, Western white pine, Western redcedar and Western hemlock. The overstory canopy of Western redcedar and Western hemlock is less than 3% as they are just beginning to become established. The understory can have high cover of the medium sized shrubs white spirea, snowberry, Oregon boxleaf, common snowberry and thinleaf huckleberry. The short statured thimbleberry can have high cover. The herbaceous layer is diverse, with medium statured beargrass occurring frequently and sometime in high cover. Other herbaceous species include the short statured prince's plume, queencup beadlily, trailplant and twinflower.

Community 2.4

Maturing Forest Western redcedar-western hemlock-subalpine fir-lodgepole pine-Engelmann spruce-western white pine/thinleaf huckleberry-snowberry-white spirea/wild sarsaparilla-heartleaf arnica-queencup beadlily-twinflower-beargrass/moss



Figure 12. Plant Community 2.4 Maturing Forest Phase of Seral Tree Species and Western Redcedar and Western Hemlock

Structure: This community is a maturing forest with vertical differentiation in the overstory tree canopy. Canopy cover averages 60%. This community has diverse tree species with Western redcedar and Western hemlock ranging 3-15% each and other seral tree species about equally distributed. These species include: Subalpine fir, Lodgepole pine, Engelmann spruce, Western white pine. The understory has patchy medium sized shrubs including: thinleaf huckleberry, snowberry, white spirea. There is a diverse understory of herbaceous species including: wild sarsaparilla, heartleaf arnica, queencup beadlily, twinflower and beargrass. There can be high cover of moss.

Community 2.5

Mature Forest Western redcedar-western hemlock (remnant seral species)/thinleaf huckleberry/threeleaf foamflower-prince's plume-queencup beadlily-western rattlesnake plantain/moss



Figure 13. Plant Community 2.5 Mature forest with some small gap dynamics, remnant seral tree species and western redcedar and western hemlock dominant.

Structure: Mature forest with vertical differentiation in the stand. Overstory is dominated by Western redcedar and Western hemlock although seral tree species are present and can have up to 15% cover each. Overstory canopy cover ranges 50-80%. Western larch is the most common seral species but others include: Grand fir, subalpine fir, Paper birch, Lodgepole pine, Western white pine, and Douglas fir. The understory is diverse but generally has low overall cover. Thinleaf huckleberry occurs in clumps and queencup beadlily and western rattlesnake plantain are common.

**Pathway 2.1A
Community 2.1 to 2.2**



Reference Community
Western redcedar-western hemlock/prince's plume-queencup beadlily-western rattlesnake plantain-twinflower/moss



Stand Initiation Regen:
Lodgepole pine-western larch (mixed seral species)/Scouler's willow-white spirea-thinleaf huckleberry/thimbleberry-Oregon boxwood/fireweed-beargrass/moss

This pathway represents a major stand-replacement disturbance such as a high-intensity fire, large scale wind event, or major insect infestation.

**Pathway 2.2A
Community 2.2 to 2.3**



Stand Initiation Regen:
Lodgepole pine-western larch (mixed seral species)/Scouler's willow-white spirea-thinleaf huckleberry/thimbleberry-Oregon boxwood/fireweed-beargrass/moss



Intermediate Aged Forest
Lodgepole pine-Douglas fir-Engelmann spruce-western larch-paper birch-subalpine fir-western white pine (western redcedar-western hemlock)/white spirea-snowberry-thinleaf huckleberry/thimbleberry/prince's plume-queencup beadlily

This pathway represents growth over time with no further significant disturbance. The areas of regeneration pass through the typical stand phases-competitive exclusion, maturation, understory reinitiating-until they resemble the old-growth structure of the reference community.

Pathway 2.3B Community 2.3 to 2.2



Intermediate Aged Forest
Lodgepole pine-Douglas fir-Engelmann spruce-western larch-paper birch-subalpine fir-western white pine (western redcedar-western hemlock)/white spirea-snowberry-thinleaf huckleberry/thimbleberry/prince's plume-queencup beadlily



Stand Initiation Regen:
Lodgepole pine-western larch (mixed seral species)/Scouler's willow-white spirea-thinleaf huckleberry/thimbleberry-Oregon boxwood/fireweed-beargrass/moss

This pathway represents a major stand-replacement disturbance such as a high-intensity fire, large scale wind event, or major insect infestation.

Pathway 2.3A Community 2.3 to 2.4



Intermediate Aged Forest
Lodgepole pine-Douglas fir-Engelmann spruce-western larch-paper birch-subalpine fir-western white pine (western redcedar-western hemlock)/white spirea-snowberry-thinleaf huckleberry/thimbleberry/prince's plume-queencup beadlily



Maturing Forest Western redcedar-western hemlock-subalpine fir-lodgepole pine-Engelmann spruce-western white pine/thinleaf huckleberry-snowberry-white spirea/wild sarsaparilla-heartleaf arnica-queencup beadlily-twinflower-beargrass/moss

This pathway represents continued growth over time with no further major disturbance.

Pathway 2.4B Community 2.4 to 2.2



Maturing Forest Western redcedar-western hemlock-subalpine fir-lodgepole pine-Engelmann spruce-western white pine/thinleaf huckleberry-snowberry-white spirea/wild sarsaparilla-heartleaf arnica-queencup beadlily-twinflower-beargrass/moss



Stand Initiation Regen:
Lodgepole pine-western larch (mixed seral species)/Scouler's willow-white spirea-thinleaf huckleberry/thimbleberry-Oregon boxwood/fireweed-beargrass/moss

This pathway represents a major stand-replacement fire disturbance, such as a major insect outbreak, or major fire event which leads to the stand initiation phase of forest development.

Pathway 2.4A Community 2.4 to 2.5



Maturing Forest Western
redcedar-western hemlock-
subalpine fir-lodgepole pine-
Engelmann spruce-western
white pine/thinleaf
huckleberry-snowberry-white
spirea/wild sarsaparilla-
heartleaf arnica-queencup
beadlily-twinflower-
beargrass/moss



Mature Forest Western
redcedar-western hemlock
(remnant seral
species)/thinleaf
huckleberry/threeleaf
foamflower-prince's plume-
queencup beadlily-western
rattlesnake plantain/moss

This pathway represents continued growth over time with no further major disturbance.

Pathway 2.5A Community 2.5 to 2.1



Mature Forest Western
redcedar-western hemlock
(remnant seral
species)/thinleaf
huckleberry/threeleaf
foamflower-prince's plume-
queencup beadlily-western
rattlesnake plantain/moss



Reference Community
Western redcedar-western
hemlock/prince's plume-
queencup beadlily-western
rattlesnake plantain-
twinflower/moss

This pathway represents no further major disturbance. Continued growth over time, as well as ongoing mortality, leads to continued vertical diversification. The community begins to resemble the structure of the reference community, with small pockets of regeneration and a more diversified understory.

Pathway 2.5B Community 2.5 to 2.2



Mature Forest Western
redcedar-western hemlock
(remnant seral
species)/thinleaf
huckleberry/threeleaf
foamflower-prince's plume-
queencup beadlily-western
rattlesnake plantain/moss



Stand Initiation Regen:
Lodgepole pine-western larch
(mixed seral
species)/Scouler's willow-
white spirea-thinleaf
huckleberry/thimbleberry-
Oregon boxwood/fireweed-
beargrass/moss

This pathway represents a major stand-replacement fire disturbance leading to the stand initiation phase of forest development.

State 3 Armillaria Root Rot Shrubland

Another disease affecting this ecological site is root rot. While Douglas-fir, grand fir, and subalpine fir are most susceptible, western redcedar and western hemlock can be affected as well. Armillaria root disease is the most common root disease fungus in this region, and is especially prevalent west of the Continental Divide. It may be difficult to detect until it has killed enough trees to create large root disease pockets or centers, ranging in size from

a fraction of an acre to hundreds of acres. The root disease spreads from an affected tree to its surrounding neighbors through root contact. The root disease effects the tree species most susceptible first, leaving less susceptible tree species that mask its presence. When root rot is severe, the pocket has abundant regeneration or dense brush growth in the center. Western redcedar is moderately resistant to *Armillaria* root rot in Idaho and Montana. The common disease expression is some mortality in saplings, and residuals of partial harvests often develop severe infections but are very slow to die (Hagle, 2010). There has been a link determined between parent material and susceptibility to root disease (Kimsey et al., 2012). Metasedimentary parent material is thought to increase the risk of root disease. Glacier National Park is dominated by metasedimentary parent material and may be more at risk than other areas to root disease (Kimsey et al., 2012). If a stand sustains very high levels of root disease mortality, then a coniferous stand could cross a threshold and become a shrubland, once all conifers are gone (Kimsey et al., 2012). Management tactics include to identify the type of *Armillaria* root disease, and manage for pines and larch. Pre-commercial thinning may improve growth and survival of pines and larch. Avoid harvests that leave susceptible species (usually Douglas-fir or true firs) as crop trees (Hagle, 2010).

Community 3.1 Shrub dominated area

Armillaria root rot induced shrubland state

Transition T1A State 1 to 2

Substantial loss of western white pine as a major seral tree species

Restoration pathway R2A State 2 to 1

Western white pine restored as a major seral tree species

Transition T2A State 2 to 3

Significant loss of susceptible tree species at a site due to *Armillaria* root rot and conversion of the forest to a shrubland

Restoration pathway R3A State 3 to 2

Conversion of the *Armillaria* root rot induced shrubland to forest, generally of less susceptible seral tree species and eventually to climax tree species

Additional community tables

Table 5. Community 2.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
pinegrass	CARU	<i>Calamagrostis rubescens</i>	–	–	0.5
Forb/Herb					
threeleaf foamflower	TITR	<i>Tiarella trifoliata</i>	–	–	0.5–15
green false hellebore	VEVI	<i>Veratrum viride</i>	–	–	3
darkwoods violet	VIOR	<i>Viola orbiculata</i>	–	–	0.5–3
common beargrass	XETE	<i>Xerophyllum tenax</i>	–	–	0.5–3
greenflowered wintergreen	PYCH	<i>Pyrola chlorantha</i>	–	–	0.5–3
claspleaf twistedstalk	STAM2	<i>Streptopus amplexifolius</i>	–	–	3

pipsissewa	CHUM	<i>Chimaphila umbellata</i>	–	–	0.5–3
bride's bonnet	CLUN2	<i>Clintonia uniflora</i>	–	–	0.5–3
American trailplant	ADBI	<i>Adenocaulon bicolor</i>	–	–	3
wild sarsaparilla	ARNU2	<i>Aralia nudicaulis</i>	–	–	3
western rattlesnake plantain	GOOB2	<i>Goodyera oblongifolia</i>	–	–	0.5–3
twinflower	LIBO3	<i>Linnaea borealis</i>	–	–	0.5–3
feathery false lily of the valley	MARA7	<i>Maianthemum racemosum</i>	–	–	3
naked miterwort	MINU3	<i>Mitella nuda</i>	–	–	3
sidebells wintergreen	ORSE	<i>Orthilia secunda</i>	–	–	0.5–3
bunchberry dogwood	COCA13	<i>Cornus canadensis</i>	–	–	0.5
mountain lady's slipper	CYMO2	<i>Cypripedium montanum</i>	–	–	0.5
lesser roundleaved orchid	PLOR4	<i>Platanthera orbiculata</i>	–	–	0.5
Pacific trillium	TROV2	<i>Trillium ovatum</i>	–	–	0.5
Fern/fern ally					
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	–	–	0.5–37.5
Shrub/Subshrub					
Woods' rose	ROWO	<i>Rosa woodsii</i>	–	–	0.5–3
common snowberry	SYAL	<i>Symphoricarpos albus</i>	–	–	0.5–3
devilsclub	OPHO	<i>Oplopanax horridus</i>	–	–	3
white spirea	SPBE2	<i>Spiraea betulifolia</i>	–	–	0.5–3
thinleaf huckleberry	VAME	<i>Vaccinium membranaceum</i>	–	–	0.5–3
grouse whortleberry	VASC	<i>Vaccinium scoparium</i>	–	–	0.5
thimbleberry	RUPA	<i>Rubus parviflorus</i>	–	–	0.5
Oregon boxleaf	PAMY	<i>Paxistima myrsinites</i>	–	–	0.5
creeping barberry	MARE11	<i>Mahonia repens</i>	–	–	0.5
Tree					
western hemlock	TSHE	<i>Tsuga heterophylla</i>	–	0–39	3–85
western redcedar	THPL	<i>Thuja plicata</i>	–	0–39	0.5–37.5
Pacific yew	TABR2	<i>Taxus brevifolia</i>	–	–	0.5–3
western larch	LAOC	<i>Larix occidentalis</i>	–	–	3
western white pine	PIMO3	<i>Pinus monticola</i>	–	0–39	3
subalpine fir	ABLA	<i>Abies lasiocarpa</i>	–	0–39	3
Rocky Mountain maple	ACGL	<i>Acer glabrum</i>	–	–	3
paper birch	BEPA	<i>Betula papyrifera</i>	–	–	0.5

Table 6. Community 2.2 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
bluejoint	CACA4	<i>Calamagrostis canadensis</i>	–	–	15
pinegrass	CARU	<i>Calamagrostis rubescens</i>	–	–	15
western fescue	FEOC	<i>Festuca occidentalis</i>	–	–	3
mountain brome	BRMA4	<i>Bromus marginatus</i>	–	–	3
Geyer's sedge	CAGE2	<i>Carex geyeri</i>	–	–	0.5
Forb/Herb					

twinflower	LIBO3	<i>Linnaea borealis</i>	–	–	0.5–37.5
fireweed	CHAN9	<i>Chamerion angustifolium</i>	–	–	0.5–15
bride's bonnet	CLUN2	<i>Clintonia uniflora</i>	–	–	0.5–15
western redcedar	THPL	<i>Thuja plicata</i>	–	0–39	0.5–15
common beargrass	XETE	<i>Xerophyllum tenax</i>	–	–	3–15
darkwoods violet	VIOR	<i>Viola orbiculata</i>	–	–	0.5–3
threeleaf foamflower	TITR	<i>Tiarella trifoliata</i>	–	–	3
bunchberry dogwood	COCA13	<i>Cornus canadensis</i>	–	–	3
western showy aster	EUCO36	<i>Eurybia conspicua</i>	–	–	3
Scouler's woollyweed	HISC2	<i>Hieracium scouleri</i>	–	–	3
narrowleaf hawkweed	HIUM	<i>Hieracium umbellatum</i>	–	–	0.5–3
pipsissewa	CHUM	<i>Chimaphila umbellata</i>	–	–	0.5–3
Canada thistle	CIAR4	<i>Cirsium arvense</i>	–	–	3
American trailplant	ADBI	<i>Adenocaulon bicolor</i>	–	–	0.5–3
western pearly everlasting	ANMA	<i>Anaphalis margaritacea</i>	–	–	0.5–3
pussytoes	ANTEN	<i>Antennaria</i>	–	–	3
spreading dogbane	APAN2	<i>Apocynum androsaemifolium</i>	–	–	3
heartleaf arnica	ARCO9	<i>Arnica cordifolia</i>	–	–	0.5–3
wild sarsaparilla	ARNU2	<i>Aralia nudicaulis</i>	–	–	0.5–3
feathery false lily of the valley	MARA7	<i>Maianthemum racemosum</i>	–	–	3
streambank wild hollyhock	ILRI	<i>Iliamna rivularis</i>	–	–	0.5–3
miterwort	MITEL	<i>Mitella</i>	–	–	3
sweetcicely	OSBE	<i>Osmorhiza berteroi</i>	–	–	0.5–3
Oregon boxleaf	PAMY	<i>Paxistima myrsinites</i>	–	–	0.5–3
goldenrod	SOLID	<i>Solidago</i>	–	–	3
common dandelion	TAOF	<i>Taraxacum officinale</i>	–	–	0.5–3
western meadow-rue	THOC	<i>Thalictrum occidentale</i>	–	–	0.5
greenflowered wintergreen	PYCH	<i>Pyrola chlorantha</i>	–	–	0.5
common yarrow	ACMI2	<i>Achillea millefolium</i>	–	–	0.5
strawberry	FRAGA	<i>Fragaria</i>	–	–	0.5
woodland strawberry	FRVE	<i>Fragaria vesca</i>	–	–	0.5
Virginia strawberry	FRVI	<i>Fragaria virginiana</i>	–	–	0.5
fragrant bedstraw	GATR3	<i>Galium triflorum</i>	–	–	0.5
autumn dwarf gentian	GEAM3	<i>Gentianella amarella</i>	–	–	0.5
western rattlesnake plantain	GOOB2	<i>Goodyera oblongifolia</i>	–	–	0.5
Pacific trillium	TROV2	<i>Trillium ovatum</i>	–	–	0.5
sidebells wintergreen	ORSE	<i>Orthilia secunda</i>	–	–	0.5
Fern/fern ally					
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	–	–	0.5–15
brackenfern	PTERI	<i>Pteridium</i>	–	–	15
Pacific oakfern	GYDI2	<i>Gymnocarpium disjunctum</i>	–	–	3
common ladyfern	ATFI	<i>Athyrium filix-femina</i>	–	–	3
Shrub/Subshrub					
kinnikinnick	ARUV	<i>Arctostaphylos uva-ursi</i>	–	–	0.5–37.5

snowbrush ceanothus	CEVE	<i>Ceanothus velutinus</i>	–	–	15–37.5
willow	SALIX	<i>Salix</i>	–	–	3–37.5
Scouler's willow	SASC	<i>Salix scouleriana</i>	–	–	0.5–37.5
white spirea	SPBE2	<i>Spiraea betulifolia</i>	–	–	3–15
common snowberry	SYAL	<i>Symphoricarpos albus</i>	–	–	3–15
thimbleberry	RUPA	<i>Rubus parviflorus</i>	–	–	0.5–15
Sitka alder	ALVIS	<i>Alnus viridis ssp. sinuata</i>	–	–	3–15
redstem ceanothus	CESA	<i>Ceanothus sanguineus</i>	–	–	15
thinleaf huckleberry	VAME	<i>Vaccinium membranaceum</i>	–	–	0.5–15
grouse whortleberry	VASC	<i>Vaccinium scoparium</i>	–	–	3
dwarf bilberry	VACE	<i>Vaccinium cespitosum</i>	–	–	3
Saskatoon serviceberry	AMAL2	<i>Amelanchier alnifolia</i>	–	–	0.5–3
oceanspray	HODI	<i>Holodiscus discolor</i>	–	–	3
dwarf red blackberry	RUPU	<i>Rubus pubescens</i>	–	–	3
Alberta saxifrage	SAOC4	<i>Saxifraga occidentalis</i>	–	–	3
sticky currant	RIVI3	<i>Ribes viscosissimum</i>	–	–	3
rose	ROSA5	<i>Rosa</i>	–	–	3
Woods' rose	ROWO	<i>Rosa woodsii</i>	–	–	0.5–3
American red raspberry	RUID	<i>Rubus idaeus</i>	–	–	3
russet buffaloberry	SHCA	<i>Shepherdia canadensis</i>	–	–	3
Greene's mountain ash	SOSC2	<i>Sorbus scopulina</i>	–	–	3
creeping barberry	MARE11	<i>Mahonia repens</i>	–	–	3
rusty menziesia	MEFE	<i>Menziesia ferruginea</i>	–	–	0.5–3
Tree					
lodgepole pine	PICO	<i>Pinus contorta</i>	–	0–39	3–62.5
Engelmann spruce	PIEN	<i>Picea engelmannii</i>	–	0–39	0.5–37.5
western larch	LAOC	<i>Larix occidentalis</i>	–	0–39	0.5–37.5
western hemlock	TSHE	<i>Tsuga heterophylla</i>	–	0–39	0.5–37.5
quaking aspen	POTR5	<i>Populus tremuloides</i>	–	0–39	0.5–15
Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	–	0–39	0.5–15
paper birch	BEPA	<i>Betula papyrifera</i>	–	–	0.5–15
subalpine fir	ABLA	<i>Abies lasiocarpa</i>	–	0–39	0.5–3
Rocky Mountain maple	ACGL	<i>Acer glabrum</i>	–	–	0.5–3
western white pine	PIMO3	<i>Pinus monticola</i>	–	0–39	0.5–3
black cottonwood	POBAT	<i>Populus balsamifera ssp. trichocarpa</i>	–	0–39	3
Nonvascular					
Moss	2MOSS	<i>Moss</i>	–	–	3

Table 7. Community 2.3 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
pinegrass	CARU	<i>Calamagrostis rubescens</i>	–	–	0.5–15
mountain brome	BRMA4	<i>Bromus marginatus</i>	–	–	0.5–3
intermediate wheatgrass	THIN6	<i>Thinopyrum intermedium</i>	–	–	0.5

intermediate wildgrass	THNU	Thimopyrum intermedium	-	-	0.5
Forb/Herb					
common beargrass	XETE	<i>Xerophyllum tenax</i>	-	-	0.5-37.5
twinflower	LIBO3	<i>Linnaea borealis</i>	-	-	0.5-37.5
feathery false lily of the valley	MARA7	<i>Maianthemum racemosum</i>	-	-	0.5-15
starry false lily of the valley	MAST4	<i>Maianthemum stellatum</i>	-	-	0.5-15
wild sarsaparilla	ARNU2	<i>Aralia nudicaulis</i>	-	-	3-15
bride's bonnet	CLUN2	<i>Clintonia uniflora</i>	-	-	0.5-15
bunchberry dogwood	COCA13	<i>Cornus canadensis</i>	-	-	3-15
redosier dogwood	COSES	<i>Cornus sericea ssp. sericea</i>	-	-	3-15
American trailplant	ADBI	<i>Adenocaulon bicolor</i>	-	-	0.5-15
heartleaf arnica	ARCO9	<i>Arnica cordifolia</i>	-	-	3-15
threeleaf foamflower	TITR	<i>Tiarella trifoliata</i>	-	-	0.5-15
claspleaf twistedstalk	STAM2	<i>Streptopus amplexifolius</i>	-	-	0.5-15
arrowleaf ragwort	SETR	<i>Senecio triangularis</i>	-	-	3
western meadow-rue	THOC	<i>Thalictrum occidentale</i>	-	-	0.5-3
darkwoods violet	VIOR	<i>Viola orbiculata</i>	-	-	0.5-3
green false hellebore	VEVI	<i>Veratrum viride</i>	-	-	0.5-3
sweetcicely	OSBE	<i>Osmorhiza berteroi</i>	-	-	0.5-3
liverleaf wintergreen	PYAS	<i>Pyrola asarifolia</i>	-	-	3
wintergreen	PYROL	<i>Pyrola</i>	-	-	3
broadleaf arnica	ARLA8	<i>Arnica latifolia</i>	-	-	0.5-3
spreading dogbane	APAN2	<i>Apocynum androsaemifolium</i>	-	-	3
yellow avalanche-lily	ERGR9	<i>Erythronium grandiflorum</i>	-	-	3
pipsissewa	CHUM	<i>Chimaphila umbellata</i>	-	-	0.5-3
miterwort	MITEL	<i>Mitella</i>	-	-	3
narrowleaf hawkweed	HIUM	<i>Hieracium umbellatum</i>	-	-	0.5-3
northern bedstraw	GABO2	<i>Galium boreale</i>	-	-	0.5-3
fragrant bedstraw	GATR3	<i>Galium triflorum</i>	-	-	3
western rattlesnake plantain	GOOB2	<i>Goodyera oblongifolia</i>	-	-	0.5-3
Scouler's woollyweed	HISC2	<i>Hieracium scouleri</i>	-	-	3
western blue virginsbower	CLOC2	<i>Clematis occidentalis</i>	-	-	3
raceme pussytoes	ANRA	<i>Antennaria racemosa</i>	-	-	3
woodland strawberry	FRVE	<i>Fragaria vesca</i>	-	-	3
arnica	ARNIC	<i>Arnica</i>	-	-	3
white thistle	CIHO	<i>Cirsium hookerianum</i>	-	-	0.5
Asian forget-me-not	MYAS2	<i>Myosotis asiatica</i>	-	-	0.5
cream pea	LAOC2	<i>Lathyrus ochroleucus</i>	-	-	0.5
strawberry	FRAGA	<i>Fragaria</i>	-	-	0.5
summer coralroot	COMA25	<i>Corallorhiza maculata</i>	-	-	0.5
woodland pinedrops	PTAN2	<i>Pterospora andromedea</i>	-	-	0.5
western sweetroot	OSOC	<i>Osmorhiza occidentalis</i>	-	-	0.5
common plantain	PLMA2	<i>Plantago major</i>	-	-	0.5
lesser roundleaved orchid	PLOR4	<i>Platanthera orbiculata</i>	-	-	0.5

sidebells wintergreen	ORSE	<i>Orthilia secunda</i>	–	–	0.5
pioneer violet	VIGL	<i>Viola glabella</i>	–	–	0.5
Pacific trillium	TROV2	<i>Trillium ovatum</i>	–	–	0.5
Fern/fern ally					
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	–	–	3–37.5
common ladyfern	ATFI	<i>Athyrium filix-femina</i>	–	–	3
Pacific oakfern	GYDI2	<i>Gymnocarpium disjunctum</i>	–	–	0.5
Shrub/Subshrub					
thimbleberry	RUPA	<i>Rubus parviflorus</i>	–	–	0.5–37.5
Oregon boxleaf	PAMY	<i>Paxistima myrsinites</i>	–	–	0.5–15
white spirea	SPBE2	<i>Spiraea betulifolia</i>	–	–	0.5–15
common snowberry	SYAL	<i>Symphoricarpos albus</i>	–	–	0.5–15
thinleaf huckleberry	VAME	<i>Vaccinium membranaceum</i>	–	–	0.5–15
Woods' rose	ROWO	<i>Rosa woodsii</i>	–	–	3–15
rusty menziesia	MEFE	<i>Menziesia ferruginea</i>	–	–	0.5–15
creeping barberry	MARE11	<i>Mahonia repens</i>	–	–	0.5–3
Utah honeysuckle	LOUT2	<i>Lonicera utahensis</i>	–	–	0.5–3
Sitka alder	ALVIS	<i>Alnus viridis ssp. sinuata</i>	–	–	3
Saskatoon serviceberry	AMAL2	<i>Amelanchier alnifolia</i>	–	–	0.5–3
russet buffaloberry	SHCA	<i>Shepherdia canadensis</i>	–	–	3
Greene's mountain ash	SOSC2	<i>Sorbus scopulina</i>	–	–	0.5–3
grouse whortleberry	VASC	<i>Vaccinium scoparium</i>	–	–	0.5–3
dwarf red blackberry	RUPU	<i>Rubus pubescens</i>	–	–	3
sticky currant	RIVI3	<i>Ribes viscosissimum</i>	–	–	0.5–3
rose	ROSA5	<i>Rosa</i>	–	–	0.5–3
Alberta saxifrage	SAOC4	<i>Saxifraga occidentalis</i>	–	–	3
oceanspray	HODI	<i>Holodiscus discolor</i>	–	–	3
prickly currant	RILA	<i>Ribes lacustre</i>	–	–	0.5
Tree					
Pacific yew	TABR2	<i>Taxus brevifolia</i>	–	–	3–62.5
lodgepole pine	PICO	<i>Pinus contorta</i>	–	0–39	0.5–37.5
Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	–	–	0.5–37.5
subalpine fir	ABLA	<i>Abies lasiocarpa</i>	–	–	0.5–37.5
western larch	LAOC	<i>Larix occidentalis</i>	–	0–39	0.5–37.5
Rocky Mountain maple	ACGL	<i>Acer glabrum</i>	–	–	0.5–15
paper birch	BEPA	<i>Betula papyrifera</i>	–	–	0.5–15
Engelmann spruce	PIEN	<i>Picea engelmannii</i>	–	0–39	0.5–15
western white pine	PIMO3	<i>Pinus monticola</i>	–	0–39	3
western hemlock	TSHE	<i>Tsuga heterophylla</i>	–	0–39	0.5–3
western redcedar	THPL	<i>Thuja plicata</i>	–	0–39	0.5–3
grand fir	ABGR	<i>Abies grandis</i>	–	0–39	0.5–3
quaking aspen	POTR5	<i>Populus tremuloides</i>	–	0–39	0.5
Nonvascular					
Moss	2MOSS	<i>Moss</i>	–	–	3–62.5

Moss	2MOSS	<i>Moss</i>	–	–	15
groundcedar	LYCO3	<i>Lycopodium complanatum</i>	–	–	3

Table 8. Community 2.4 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
pinegrass	CARU	<i>Calamagrostis rubescens</i>	–	–	3–15
mountain brome	BRMA4	<i>Bromus marginatus</i>	–	–	0.5–3
sedge	CAREX	<i>Carex</i>	–	–	0.5
Forb/Herb					
common beargrass	XETE	<i>Xerophyllum tenax</i>	–	–	0.5–37.5
twinflower	LIBO3	<i>Linnaea borealis</i>	–	–	3–37.5
starry false lily of the valley	MAST4	<i>Maianthemum stellatum</i>	–	–	3–15
bride's bonnet	CLUN2	<i>Clintonia uniflora</i>	–	–	0.5–15
bunchberry dogwood	COCA13	<i>Cornus canadensis</i>	–	–	0.5–15
heartleaf arnica	ARCO9	<i>Arnica cordifolia</i>	–	–	3–15
wild sarsaparilla	ARNU2	<i>Aralia nudicaulis</i>	–	–	0.5–15
roughfruit fairybells	PRTR4	<i>Prosartes trachycarpa</i>	–	–	15
claspleaf twistedstalk	STAM2	<i>Streptopus amplexifolius</i>	–	–	3–15
western meadow-rue	THOC	<i>Thalictrum occidentale</i>	–	–	3
threeleaf foamflower	TITR	<i>Tiarella trifoliata</i>	–	–	3
greenflowered wintergreen	PYCH	<i>Pyrola chlorantha</i>	–	–	3
aster	ASTER	<i>Aster</i>	–	–	3
northern bedstraw	GABO2	<i>Galium boreale</i>	–	–	0.5–3
common cowparsnip	HEMA80	<i>Heracleum maximum</i>	–	–	3
green false hellebore	VEVI	<i>Veratrum viride</i>	–	–	3
violet	VIOLA	<i>Viola</i>	–	–	0.5–3
darkwoods violet	VIOR	<i>Viola orbiculata</i>	–	–	0.5–3
American trailplant	ADBI	<i>Adenocaulon bicolor</i>	–	–	3
woodland strawberry	FRVE	<i>Fragaria vesca</i>	–	–	3
pipsissewa	CHUM	<i>Chimaphila umbellata</i>	–	–	0.5–3
western blue virginsbower	CLOC2	<i>Clematis occidentalis</i>	–	–	3
feathery false lily of the valley	MARA7	<i>Maianthemum racemosum</i>	–	–	3
western rattlesnake plantain	GOOB2	<i>Goodyera oblongifolia</i>	–	–	3
narrowleaf hawkweed	HIUM	<i>Hieracium umbellatum</i>	–	–	3
sweetcicely	OSBE	<i>Osmorhiza berteroi</i>	–	–	0.5
fragrant bedstraw	GATR3	<i>Galium triflorum</i>	–	–	0.5
common yarrow	ACMI2	<i>Achillea millefolium</i>	–	–	0.5
Fern/fern ally					
horsetail	EQUIS	<i>Equisetum</i>	–	–	3
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	–	–	3
common ladyfern	ATFI	<i>Athyrium filix-femina</i>	–	–	0.5–3
western showy aster	EUCO36	<i>Eurybia conspicua</i>	–	–	0.5–3
Pacific oakfern	GYDI2	<i>Gymnocarpium disiunctum</i>	–	–	3

Shrub/Subshrub					
Utah honeysuckle	LOUT2	<i>Lonicera utahensis</i>	–	–	3–15
Oregon boxleaf	PAMY	<i>Paxistima myrsinites</i>	–	–	3–15
white spirea	SPBE2	<i>Spiraea betulifolia</i>	–	–	3–15
thimbleberry	RUPA	<i>Rubus parviflorus</i>	–	–	3–15
thinleaf huckleberry	VAME	<i>Vaccinium membranaceum</i>	–	–	0.5–15
common snowberry	SYAL	<i>Symphoricarpos albus</i>	–	–	3–15
Greene's mountain ash	SOSC2	<i>Sorbus scopulina</i>	–	–	3–15
kinnikinnick	ARUV	<i>Arctostaphylos uva-ursi</i>	–	–	3
Sitka alder	ALVIS	<i>Alnus viridis ssp. sinuata</i>	–	–	0.5–3
dwarf red blackberry	RUPU	<i>Rubus pubescens</i>	–	–	3
prickly currant	RILA	<i>Ribes lacustre</i>	–	–	3
sticky currant	RIVI3	<i>Ribes viscosissimum</i>	–	–	0.5–3
Woods' rose	ROWO	<i>Rosa woodsii</i>	–	–	0.5–3
devilsclub	OPHO	<i>Oplopanax horridus</i>	–	–	3
rusty menziesia	MEFE	<i>Menziesia ferruginea</i>	–	–	3
creeping barberry	MARE11	<i>Mahonia repens</i>	–	–	3
Saskatoon serviceberry	AMAL2	<i>Amelanchier alnifolia</i>	–	–	3
redosier dogwood	COSES	<i>Cornus sericea ssp. sericea</i>	–	–	3
grouse whortleberry	VASC	<i>Vaccinium scoparium</i>	–	–	0.5
Tree					
Pacific yew	TABR2	<i>Taxus brevifolia</i>	–	–	3–37.5
subalpine fir	ABLA	<i>Abies lasiocarpa</i>	–	–	3–37.5
lodgepole pine	PICO	<i>Pinus contorta</i>	–	0–39	3–37.5
Engelmann spruce	PIEN	<i>Picea engelmannii</i>	–	0–39	3–37.5
western white pine	PIMO3	<i>Pinus monticola</i>	–	0–39	3–37.5
Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	–	0–39	0.5–15
western larch	LAOC	<i>Larix occidentalis</i>	–	0–39	3–15
paper birch	BEPA	<i>Betula papyrifera</i>	–	–	0.5–15
western redcedar	THPL	<i>Thuja plicata</i>	–	0–39	3–15
western hemlock	TSHE	<i>Tsuga heterophylla</i>	–	0–39	3–15
black cottonwood	POBAT	<i>Populus balsamifera ssp. trichocarpa</i>	–	0–39	15
Rocky Mountain maple	ACGL	<i>Acer glabrum</i>	–	–	0.5–3
grand fir	ABGR	<i>Abies grandis</i>	–	–	0.5–3
Nonvascular					
Moss	2MOSS	Moss	–	–	3–15

Table 9. Community 2.5 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
mountain brome	BRMA4	<i>Bromus marginatus</i>	–	–	0.5–3
pinegrass	CARU	<i>Calamagrostis rubescens</i>	–	–	3
Forb/Herb					
common beargrass	YETE	<i>Xerophyllum tenax</i>	–	–	37.5

Common Beargrass		<i>Asophrum tenua</i>	–	–	37.5
threeleaf foamflower	TITR	<i>Tiarella trifoliata</i>	–	–	3–15
wild sarsaparilla	ARNU2	<i>Aralia nudicaulis</i>	–	–	15
bride's bonnet	CLUN2	<i>Clintonia uniflora</i>	–	–	0.5–15
twinflower	LIBO3	<i>Linnaea borealis</i>	–	–	0.5–15
arnica	ARNIC	<i>Arnica</i>	–	–	0.5–3
cream pea	LAOC2	<i>Lathyrus ochroleucus</i>	–	–	3
starry false lily of the valley	MAST4	<i>Maianthemum stellatum</i>	–	–	0.5–3
sidebells wintergreen	ORSE	<i>Orthilia secunda</i>	–	–	3
sweetcicely	OSBE	<i>Osmorhiza berteroi</i>	–	–	3
lesser roundleaved orchid	PLOR4	<i>Platanthera orbiculata</i>	–	–	3
bunchberry dogwood	COCA13	<i>Cornus canadensis</i>	–	–	3
fragrant bedstraw	GATR3	<i>Galium triflorum</i>	–	–	3
western rattlesnake plantain	GOOB2	<i>Goodyera oblongifolia</i>	–	–	0.5–3
fireweed	CHAN9	<i>Chamerion angustifolium</i>	–	–	3
pipsissewa	CHUM	<i>Chimaphila umbellata</i>	–	–	0.5–3
American trailplant	ADBI	<i>Adenocaulon bicolor</i>	–	–	3
greenflowered wintergreen	PYCH	<i>Pyrola chlorantha</i>	–	–	3
claspleaf twistedstalk	STAM2	<i>Streptopus amplexifolius</i>	–	–	3
darkwoods violet	VIOR	<i>Viola orbiculata</i>	–	–	3
green false hellebore	VEVI	<i>Veratrum viride</i>	–	–	0.5
violet	VIOLA	<i>Viola</i>	–	–	0.5
mountain lady's slipper	CYMO2	<i>Cypripedium montanum</i>	–	–	0.5
feathery false lily of the valley	MARA7	<i>Maianthemum racemosum</i>	–	–	0.5
Fern/fern ally					
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	–	–	3–37.5
common ladyfern	ATFI	<i>Athyrium filix-femina</i>	–	–	3
Pacific oakfern	GYDI2	<i>Gymnocarpium disjunctum</i>	–	–	3
Shrub/Subshrub					
thinleaf huckleberry	VAME	<i>Vaccinium membranaceum</i>	–	–	0.5–15
Woods' rose	ROWO	<i>Rosa woodsii</i>	–	–	3
Saskatoon serviceberry	AMAL2	<i>Amelanchier alnifolia</i>	–	–	3
rusty menziesia	MEFE	<i>Menziesia ferruginea</i>	–	–	3
Utah honeysuckle	LOUT2	<i>Lonicera utahensis</i>	–	–	3
devilsclub	OPHO	<i>Oplopanax horridus</i>	–	–	0.5–3
Oregon boxleaf	PAMY	<i>Paxistima myrsinites</i>	–	–	0.5–3
creeping barberry	MARE11	<i>Mahonia repens</i>	–	–	0.5
white spirea	SPBE2	<i>Spiraea betulifolia</i>	–	–	0.5
Tree					
Pacific yew	TABR2	<i>Taxus brevifolia</i>	–	–	37.5–62.5
western redcedar	THPL	<i>Thuja plicata</i>	–	0–39	3–37.5
western hemlock	TSHE	<i>Tsuga heterophylla</i>	–	0–39	15–37.5
western white pine	PIMO3	<i>Pinus monticola</i>	–	0–39	15
Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	–	0–39	15

western larch	LAOC	<i>Larix occidentalis</i>	–	0–39	0.5–15
paper birch	BEPA	<i>Betula papyrifera</i>	–	–	0.5–3
grand fir	ABGR	<i>Abies grandis</i>	–	0–39	3
subalpine fir	ABLA	<i>Abies lasiocarpa</i>	–	0–39	3
Rocky Mountain maple	ACGL	<i>Acer glabrum</i>	–	–	3
lodgepole pine	PICO	<i>Pinus contorta</i>	–	0–39	3
Nonvascular					
Moss	2MOSS	<i>Moss</i>	–	–	37.5–85

References

. 2001. Flathead National Forest Plan Appendix B..

Arno, S. 2001. Forest Regions of Montana. USDA Forest Service Research Paper INT-218. USFS, USDA.

Arno, S. and R. Hammerly. 2007. Northwest Trees. in The Mountaineers Books.

Arno, S., D. Parsons, and R. Keane. 2000. Mixed-Severity Fire Regimes in the Northern Rocky Mountains: Consequences of Fire Exclusion and Options for the Future. USDA Forest Service Proceedings RMRS-P-15-VOL-5.2000. USDA.

Barrett, S.W., S. Arno, and C. Key. 1991. Fire regimes of western larch-lodgepole pine forests in Glacier National Park, Montana.

Bollenbacher, B., P. Kolb, and J. Morrison. 2013. Review Draft: Vulnerability, exposure, and sensitivity in restoring and maintaining the adaptive capacity of forest landscapes in the northern region of the Northern Rocky Mountains..

Byler, J. and S. Hagle. 2000. Succession functions of pathogens and insects.

Fins, L., J. Byler, D. Ferguson, A. Harvey, M. Frances Mahalovich, G. McDonald, D. Miller, J. Schwandt, and A. Zack. 2000. Return of the Giants.

Fischer, W. and A. Bradley. 2000. Fire Ecology of Western Montana Forest Habitat Types. Intermountain Research Station. GTR-INT-223. US Department of Agriculture. Forest Service..

Garrison-Johnston, R., R. Lewis, and L. Johnson. 2007. Northern Idaho and Western Montana Nutrition Guidelines by Rock Type. Intermountain Forest Tree Nutrition Cooperative. Forest Resources Department, University of Idaho.

Green, P., D. Sirucek, W. Hann, A. Zack, and B. Naumann. 1992. Old Growth Criteria. R-1 SES 4/92. Northern Region. USDA USFS.

Habeck J R. 1968. Forest succession in the Glacier Park cedar-hemlock forests. Ecology. Ecology. 872–880.

Hagle, S. 2008. Management Guide for Armillaria Root Disease. USFS, Forest Health Protection and State Forestry

Organizations. USFS, USDA.

- Haig, Irvine, Davis, R. Weidman, and R. Kenneth. 1941. Natural regeneration in the western white pine type. Technical Bulletin no. 767. Northern Rocky mountain Forest and Range Experiment Station. USFS.
- Harvey, A., J. Byler, G. McDonald, L. Neuenschwander, and J. Tonn. 2008. Death of an Ecosystem: Perspectives on Western White Pine Ecosystems of North America at the End of the Twentieth Century. USDA Forest Service RMRS-GTR-208. USDA Forest Service.
- Hoff, R.J., G.I. McDonald, and R.T. Bingham. 1976. Mass selection for blister rust resistance: a method for natural regeneration of western white pine. US Dept. of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station.
- Jones, G.P. 2004 (Date accessed). U.S. National Vegetation Classification Standard: Terrestrial Ecological Classifications.
- Kimsey, M., T. Shaw, M. Johnson, and P. McDaniel. 2012. Ecological and physiological overview of volcanic soils and their influence on tree growth and vegetation.
- Kimsey, M. 2012. Geospatial tools for estimating and maintaining soil-site productivity. Pages in Northwest Forest Soils Council Meeting, February 28, 2012. Intermountain Forest Tree Nutrition Cooperative.
- Loehman, R.A., J.A. Clark, and R.E. Keane. 2011. Modeling Effects of Climate Change and Fire Management on Western White Pine (*Pinus monticola*) in the Northern Rocky Mountains, USA. *Forests* 2011:832–860.
- Losensky, J.L. 1992. Personal communication. Jack Losensky.
- Mahalovich, M.F. 1995. The role of genetics in improving forest health.
- McDonald, G.L., A.E. Harvey, and J.R. Tonn. 2000. Fire, Competition, and Forest Pests: Landscape Treatment to Sustain Ecosystem Functions, The Joint Fire Science Conference and Workshop.. Pages 195–211 in Proceedings from the Joint Fire Science Conference and Workshop: crossing the millennium: integrating spatial technologies and ecological principles for a new age in fire management.
- McKenzie, D. and D. Tinker. 2012. Fire-induced shifts in overstory tree species composition and associated understory plant composition in Glacier National Park, Montana. *Plant Ecology*. *Plant Ecology*. 207–224.
- Pfister, R., B. Kovalchik, S. Arno, and R. Presby. 1977. Forest Habitat Types of Montana. USDA Forest Service General Technical Report INT-34. Intermountain Forest and Range Experiment Station, US Department of Agriculture. Intermountain Forest and Range Experiment Station, US Department of Agriculture.
- Soil Survey Staff. 2015. Illustrated guide to soil taxonomy. National Soil Survey Center. U.S. Department of Agriculture, Natural Resources Conservation Service, Lincoln, Nebraska.
- Tomback, D.F. and P. Achuff. 2010. Blister rust and western forest biodiversity: ecology, values and outlook for white pines.. *Forest Pathology*. *Forest Pathology*. 186–225.

Zack, A. 2005. Region One, Vegetation Classification, mapping, inventory and analysis report. U.S. Department of Agriculture, US Forest Service, Northern Region. Report 09-08 v1.0. 1997, revised 2005. U.S. Department of Agriculture, US Forest Service, Northern Region.

Other references

References

Arno, S. Forest Regions of Montana. USDA Forest Service Research Paper INT-218. USFS. USDA.

Arno, S. and R. Hammerly. Northwest Trees, by Stephen F. Arno and Ramona P. Hammerly. Anniversary Edition, the Mountaineers Books, 2007.

Arno S., D. Parsons and R. Keane. Mixed-Severity Fire Regimes in the Northern Rocky Mountains: Consequences of Fire Exclusion and Options for the Future. USDA Forest Service Proceedings RMRS-P-15-VOL-5.2000.

Barrett, S., S. Arno and C. Key. Fire regimes of western larch-lodgepole pine forests in Glacier National Park, Montana. 1991.

Bollenbacher, B. and P. Kolb, J. Morrison. 2013. Review Draft: Vulnerability, exposure, and sensitivity in restoring and maintaining the adaptive capacity of forest landscapes in the northern region of the Northern Rocky Mountains.

Byler, James and Hagle, Susan. 2000. Succession functions of pathogens and insects. FHP Report No. 00-09.
Fins, Lauren, et al. "Return of the giants: restoring western white pine to the Inland Northwest." *Journal of forestry* 100.4 (2002): 20-26.

Fischer W., A. Bradley. Fire Ecology of Western Montana Forest Habitat Types. US Department of Agriculture. Forest Service. Intermountain Research Station. GTR-INT-223.

Flathead National Forest Plan. 2001. Appendix B.

Garrison-Johnston, R. Lewis, L. Johnson. 2007. Northern Idaho and Western Montana Nutrition Guidelines by Rock Type. Intermountain Forest Tree Nutrition Cooperative. Forest Resources Department, University of Idaho.

Green, P. J. Joy, D. Sirucek, W. Hann, A. Zack, and B. Naumann. April 1992. Old Growth Criteria. R-1 SES 4/92. Northern Region. USDA USFS.

Habeck, J.R. 1968. Forest succession in the Glacier Park cedar-hemlock forests. *Ecology* 49: 872-880.

Hagle S., USFS, Forest Health Protection and State Forestry Organizations. Management Guide for Armillaria Root Disease. February 2008. WEB July 2010.

Hagle, Susan K. "Succession functions of forest pathogens and insects." (2000).

Haig, Irvine, Davis, Kenneth and Weidman, Robert. Natural regeneration in the western white pine type. Technical Bulletin no. 767. May 1941. Northern Rocky mountain Forest and Range Experiment Station. USFS.

Harvey A., James Byler, Gerald McDonald, Leon Neuenschwander, Jonalea Tonn. Death of an Ecosystem: Perspectives on Western White Pine Ecosystems of North America at the End of the Twentieth Century. USDA Forest Service RMRS-GTR-208. 2008.

Hoff, Raymond J., GERAL I. McDonald, and Richard T. Bingham. Mass selection for blister rust resistance: a method for natural regeneration of western white pine. US Dept. of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, 1976.

Kimsey M., T. Shaw, M. Johnston, P. McDaniel. Intermountain Forest Tree Nutrition Cooperative. Ecological and physiological overview of volcanic soils and their influence on tree growth and vegetation.

Kimsey M. Intermountain Forest Tree Nutrition Cooperative. Geospatial tools for estimating and maintaining soil-site productivity. Northwest Forest Soils Council Meeting, February 28, 2012.

Lauren Fins, James Byler, Dennis Ferguson, Al Harvey, Mary Frances Mahalovich, Gerald McDonald, Dan Miller, John Schwandt, and Art Zack. Return of the Giants.

Little, E.L., Jr. Digital Representation of "Atlas of United States Trees"; U.S. Geological Survey Professional Paper 1650; U.S. Geological Survey: Reston, VA, USA, 1999. [Google Scholar]
Vulnerability, Exposure, and Sensitivity in Restoring and maintaining the adaptive capacity of forest landscapes in the Northern Region of the Northern Rocky Mountains. Review Draft.

Loehman, Rachel A., Jason A. Clark, and Robert E. Keane. "Modeling effects of climate change and fire management on western white pine (*Pinus monticola*) in the Northern Rocky Mountains, USA." *Forests* 2.4 (2011): 832-860.

Losensky, J. L. "Personal communication. Jack Losensky." *Ecologist*, Lolo National Forest, Missoula, MT (1992).

Mahalovich, Mary F. "The role of genetics in improving forest health." (1995).

McDonald, A. Harvey and J. Tonn. USDA U.S.F.S., Rocky Mountain Research Station. Fire, competition and forest pests: landscape treatment to sustain ecosystem function.

McKenzie, D. and D. Tinker. 2012. Fire-induced shifts in overstory tree species composition and associated understory plant composition in Glacier National Park, Montana. *Plant Ecology* 2012: 213:207-224.

N.P.S. Fire Ecology Annual Report, Calendar Year 2014.

NatureServe, 2007. U.S. National Vegetation Classification Standard: Terrestrial Ecological Classifications. Waterton-Glacier International Peace Park, Local and Global Association Descriptions.

Pfister, R., B. Kovalchik, S. Arno, R. Presby. Forest Habitat Types of Montana. USDA Forest Service General Technical Report INT-34. Intermountain Forest and Range Experiment Station, US Department of Agriculture. May 1977.

Rockwell, F.I. 1917. Western white pine bulletin.

Skovlin, J. Personal communication, 2015.

Soil Survey Staff. 2015. Illustrated guide to soil taxonomy. U.S. Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, Nebraska.

Tomback, Diana F., and Peter Achuff. "Blister rust and western forest biodiversity: ecology, values and outlook for white pines." *Forest Pathology* 40.3-4 (2010): 186-225.

Zack, A. Region One, Vegetation Classification, mapping, inventory and analysis report. U.S. Department of Agriculture, US Forest Service, Northern Region. Report 09-08 v1.0. 1997, revised 2005.

NPS Stand Age spatial layer

USFS Stand Health map

Approval

Kirt Walstad, 9/08/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	12/18/2020
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:**

- 2. Presence of water flow patterns:**

- 3. Number and height of erosional pedestals or terracettes:**

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

- 5. Number of gullies and erosion associated with gullies:**

- 6. Extent of wind scoured, blowouts and/or depositional areas:**

- 7. Amount of litter movement (describe size and distance expected to travel):**

- 8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial**

distribution on infiltration and runoff:

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-