

Ecological site EX043B23A122 Loamy (Ly) Absaroka Lower Foothills

Last updated: 3/27/2019
Accessed: 04/27/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Marji Patz, Ray Gullion
Contact for lead author	marji.patz@wy.usda.gov or 307-271-3130
Date	07/18/2014
Approved by	Marji Patz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Rare to nonexistent. Where present, short and widely spaced.

- 2. Presence of water flow patterns:** Barely observable.

- 3. Number and height of erosional pedestals or terracettes:** Rare to nonexistent.

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground can range from 10-30%.

- 5. Number of gullies and erosion associated with gullies:** Active gullies should not be present.

- 6. Extent of wind scoured, blowouts and/or depositional areas:** Rare to nonexistent.

- 7. Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter expected to move only in small amounts (to leeward side of shrubs). Large woody debris from sagebrush will show no movement.

-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 1 (interspaces) to 6 (under plant canopy), but average values should be 3.0 or greater.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil data is limited for this site. Described A-horizons vary from 1-12 inches (3-30 cm) with OM of 1 to 2%.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The plant community consists of 60-75% grasses, 10% forbs and 15-30% shrubs. Evenly distributed plant canopy (35-55%) and litter plus moderate to moderately rapid infiltration rates result in minimal runoff. Basal cover is typically less than 8% for this site and does very little to effect runoff on this site.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Mid-stature, cool-season bunchgrasses
- Sub-dominant: perennial shrubs = cool-season rhizomatous grasses
- Other: perennial forbs short-stature cool-season bunchgrasses
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal decadence, typically associated with shrub component of the canopy cover.
-
14. **Average percent litter cover (%) and depth (in):** Litter ranges from 20-45% of total canopy measurement with total litter (including beneath the plant canopy) from 35-85% expected. Herbaceous litter depth typical
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** English: 350 - 1125 lbs/ac (650 lbs/ac average); Metric: 392 - 1262 kg/ha (729 kg/ha average).
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site: The increase of bare ground above 35% is an indicator that a threshold is being crossed. Corresponding increase will be noted in one or more of the following species is common: Blue grama, Sandberg bluegrass, Wyoming big sagebrush, buckwheats, and Spiny phlox. Annual weeds such as kochia, mustards, Lambsquarter, Russian thistle, and pepperweeds are common invasive species in disturbed sites. Common noxious weeds that invade are: Cheatgrass (Downy brome), knapweeds, thistles (Bull, Canada), Houndstongue, Black henbane and Whitetop.

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in drought years.
-