

## Ecological site EX043B23A154 Shale (Sh) Absaroka Lower Foothills

Last updated: 10/04/2019  
Accessed: 05/02/2024

---

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Ray Gullion, E. Bainter
Contact for lead author	ray.gullion@wy.usda.gov or 307-347-2456
Date	05/01/2008
Approved by	E. Bainter
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:** Some rills to be expected on this site. Depending on slope, rills range from .5-2 inches (1-5 cm) wide and are found every 3-6 feet (1-2 m).

---
- 2. Presence of water flow patterns:** Barely observable.

---
- 3. Number and height of erosional pedestals or terracettes:** Slight pedestalling evident.

---
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground can range from 40-60%.

---
- 5. Number of gullies and erosion associated with gullies:** Active gullies, if present, should be rare.

---
- 6. Extent of wind scoured, blowouts and/or depositional areas:** Minimal to nonexistent.

---
- 7. Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter expected to move

only in small amounts. Woody debris will show only slight movement down slope.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 3 (interspaces) to 6 (under plant canopy), but average values should be 3.5 or greater.
- 

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Typically an A-horizon of 1-5 inches (3-12 cm) with weak granular or platy structure and brown to gray in color with OM of .5 to 1%.
- 

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 55-75% grasses, 10% forbs, and 15-35% shrubs. Minimal plant canopy (20-50%) and litter plus moderate to slow infiltration rates result in moderate runoff. Basal cover is typically less than 5% and does very little to effect runoff on this site.
- 

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer exists, but soil crusting in dry conditions is typical.
- 

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: perennial shrubs mid-size, cool season bunchgrasses

Sub-dominant: cool season rhizomatous grasses perennial forbs

Other: short, cool season bunchgrasses

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal decadence, typically associated with shrub component.
- 

14. **Average percent litter cover (%) and depth ( in):** Litter ranges from 5-25% of total canopy measurement with total litter (including beneath the plant canopy) from 15-50% expected. Herbaceous litter depth is typically shallow, ranging from 2-8mm. Woody litter can be up to an inch (3 cm).
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** English: 75 -300 lb/ac (188 lb/ac average); Metric: 84 -336 kg/ha (210 kg/ha average).
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if**

**their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Bare ground greater than 80% is the most common indicator of a threshold being crossed. Short warm season grasses, birdfoot sagebrush, Woodyaster and phlox are common increasers. Annual weeds such as halogeton, kochia, and Russian thistle are common invasive species in disturbed sites.

---

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in drought years.
-