

## Ecological site EX043B23A172 Stony Upland (StU) Absaroka Lower Foothills

Last updated: 3/04/2024  
Accessed: 04/29/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Marji Patz
Contact for lead author	marji.patz@usda.gov; 307-271-3130
Date	04/23/2020
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- Number and extent of rills:** Rare to nonexistent. Some increase in rill development may occur on steeper slopes or on areas located below exposed bedrock or other water shedding areas where increased runoff may occur. Where rills are present, they should be fairly short (2-5 feet), <1 inch deep and somewhat widely spaced (4-8 feet). Rills may increase in length (3-6 feet) and decrease in spacing (3-6 feet) on slopes greater than 30 percent. A minor increase in rill development may be observed on all slopes following major thunderstorm or spring runoff events but should heal during the next growing season.
- Presence of water flow patterns:** Barely observable. Some very minor evidence of water flow patterns may be found around perennial plant bases. They show little evidence of current erosion. They are expected to be short (3-6 feet), stable, sinuous, and not connected. There may be very minor evidence of deposition. Evidence of water flow may increase somewhat with slope.
- Number and height of erosional pedestals or terracettes:** Perennial vegetation shows little evidence of erosional pedestalling (1 to 2% of individual plants). Plant roots are covered and litter remains in place around plant crowns. Terracettes should be absent or, if present, stable. A slight increase in both pedestal and terracette development may occur with increasing slope.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground can range from 0-20%. Bare ground spaces should not be greater than 2 foot in diameter.

- 
5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present.
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** Rare to nonexistent. No evidence of wind generated soil movement is present. Wind caused blowouts and deposition are not present.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Herbaceous and large woody litter not expected to move. Most litter resides in place with some redistribution downslope caused by water movement. The majority of litter accumulates at the base of plants. Some grass leaves and stems may accumulate in soil depressions adjacent to plants. Woody stems are not likely to move. However, some litter movement is expected (up to 6 feet) with increases in slopes >25% and/or increased runoff resulting from heavy thunderstorms.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil Stability Index ratings range from 3 (interspaces) to 6 (under plant canopy), but average values should be 4.0 or greater.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil data is limited for this site. Soil OM of 2 to 5% is expected.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community consists of 50-80% grasses, 15% forbs, and 5-35% shrubs. Evenly distributed plant canopy (60-95%) and litter plus moderate infiltration rates result in minimal runoff. Basal cover is typically 5-15% for this site and does affect runoff on this site. Surface rock fragments of 5-20% provide stability to the site, but reduce infiltration
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Mid-stature, cool-season bunchgrasses >>
- Sub-dominant: perennial shrubs >> perennial forbs>>>
- Other: tall-stature cool-season bunchgrasses = cool-season rhizomatous grasses = short-stature cool-season bunchgrasses
- Additional: Following a disturbance such as fire, drought, rodents or insects that remove woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community for a period of time. If a disturbance has not occurred for an extended period of time, woody species may continue to increase. These conditions would reflect a functional community phase within the reference state.

- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** All age classes of perennial grasses should be present under average to above average growing conditions. There may be partial mortality on individual bunchgrasses and shrubs during drought periods, and complete mortality of individual plants during severe drought periods. Slight decadence in the principle shrubs could occur near the end of the fire cycle or during periods of extended drought, or insect infestations. In general, a mix of age classes should be expected with some dead and decadent plants present.
- 
14. **Average percent litter cover (%) and depth ( in):** Litter ranges from 5-30% of total canopy measurement with total litter (including beneath the plant canopy) from 30-50% expected. Herbaceous litter depth typically ranges from 5-15 mm. Woody litter can be up to a couple inches (4-6 cm).
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Average annual production is 600 lb/ac (673 kg/ha), and ranges from 450-800 lb/ac (504-897 kg/ha).
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Bare ground greater than 35% is the most common indicator of a threshold being crossed. Rhizomatous wheatgrasses, Sandberg bluegrass, juniper and big sagebrush are common increasers. Common dandelion, thistles, and annual weeds such as cheatgrass and mustards are common invasive species in disturbed sites.
- 
17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in extreme drought years.
-