

Ecological site R043BP820MT Upland Shrubland Group

Last updated: 3/01/2024
Accessed: 05/02/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043B—Central Rocky Mountains

The Central Rocky Mountains (MLRA 43B) of Montana occupy some 28,850 square miles and exist primarily in Central and SW portions of the state. The climate is extremely variable with precipitation lows of 9 to 100 inches per year and frost free days of less than 30 to over 110 days. The geology of the region is also highly variable. The combination of variable climate and geology create a complex relationship of plant communities. MLRA 43B elevations typically exist between 6000 and 12,799ft at Granite Peak (the highest point in Montana).

The Continental Divide runs through this MLRA effectively splitting its watershed to contribute to either the Missouri River to the East and the Columbia River to the West.

Ecological site concept

- Site does not receive any additional water
- Soils are
 - o Generally not saline or saline-sodic (limited extent)
 - o Moderately deep, deep, or very deep
 - o Typically less than 5% stone and boulder cover (<15% max)
- Soil surface texture ranges from sandy loam to clay loam in surface mineral 4"
- Area of rugged mountain, hills, plateaus, and valleys of the Central Rocky Mountains in Southwest Montana.
- Moisture Regime: ustic to udic (primarily udic)
- Temperature Regime: frigid to cryic (primarily cryic)
- Dominant Cover: rangeland (deciduous shrub dominated)
- Elevation Range: 3800-9700ft
- Slope: 0-60 (typically less than 25%)

Site Development and Testing Plan

This Provisional Ecological Site Description was developed to meet the criteria as defined in Soil Survey National Instruction part 306 (430-306-NI, April 2015) as interpreted by Regional Ecological Site Specialist. Information in this description are first approximations based on broad groupings of soil properties and vegetation characteristics associated with those groupings. Although this description has been through the quality control and quality assurance review process it has not been certified for use in conservation planning.

Associated sites

R043BP818MT	Upland Grassland Group The Upland Shrubland occupies the same landscape position and is often intermixed with the Upland Grassland. The two sites will have significant plant community overlap and have similar state and transition models.
-------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

R043BP819MT	Upland Sagebrush Shrubland Group The Upland Sagebrush Shrubland occupies the same landscape position and is often intermixed with the Upland Shrubland. The two sites will have significant plant community overlap and have similar state and transition models.
-------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Similar sites

R043BP818MT	Upland Grassland Group The Upland Shrubland occupies the same landscape position and is often intermixed with the Upland Grassland. The two sites will have significant plant community overlap and have similar state and transition models.
R043BP819MT	Upland Sagebrush Shrubland Group The Upland Sagebrush Shrubland occupies the same landscape position and is often intermixed with the Upland Shrubland. The two sites will have significant plant community overlap and have similar state and transition models.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Symphoricarpos albus</i> (2) <i>Dasiphora fruticosa</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Festuca campestris</i>

Physiographic features

This ecological site occurs on nearly level to very steep hills, escarpments, and plateaus. It often occurs in complex with other ecological sites. This site occurs on most slopes and exposures; however, typically less than 25 percent slope. Aspect sometimes becomes significant, especially on steep and very steep slopes. Variations in plant community composition and production can result due to aspect. Runoff and potential for water erosion are important features of this site.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Mountains > Hillslope (3) Mountains > Escarpment (4) Mountains > Plateau
Runoff class	Low to high
Elevation	3,800–10,000 ft
Slope	0–60%
Water table depth	100 in
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

Climate is quite variable on this site but will typically receive between 10 to 30 inches of precipitation with approximately 20 to 100 frost-free days. The climate of this site makes it ideal for a herbaceous dominated system with little or no shrub production.

Table 3. Representative climatic features

Frost-free period (characteristic range)	15-80 days
Freeze-free period (characteristic range)	47-113 days
Precipitation total (characteristic range)	13-23 in

Frost-free period (actual range)	4-102 days
Freeze-free period (actual range)	38-129 days
Precipitation total (actual range)	11-26 in
Frost-free period (average)	51 days
Freeze-free period (average)	90 days
Precipitation total (average)	19 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) PHILIPSBURG RS [USC00246472], Philipsburg, MT
- (2) WISE RIVER 3 WNW [USC00249082], Wise River, MT
- (3) BUTTE BERT MOONEY AP [USW00024135], Butte, MT
- (4) NORRIS MADISON PH [USC00246157], Ennis, MT
- (5) BIG SKY 2WNW [USC00240775], Gallatin Gateway, MT
- (6) PONY [USC00246655], Cardwell, MT
- (7) GARDINER [USC00243378], Gardiner, MT
- (8) HEBGEN DAM [USC00244038], West Yellowstone, MT
- (9) WEST YELLOWSTONE [USC00248857], West Yellowstone, MT
- (10) MYSTIC LAKE [USC00245961], Fishtail, MT
- (11) RED LODGE [USC00246918], Red Lodge, MT
- (12) WILLSALL 8 ENE [USC00249023], Willsall, MT

Influencing water features

n/a

Wetland description

n/a

Soil features

These soils are shallow, moderate to moderately rapid permeability, and are well to somewhat excessively drained. These soils formed from residuum from mixed origins primarily from non-calcareous geology. Typically soil surface textures consist of loam, clay loam, and silt loam textures. Soils are sometimes gravelly surface however will vary up depending on its association with to a neighboring sites.

Table 4. Representative soil features

Parent material	(1) Colluvium–igneous, metamorphic and sedimentary rock (2) Residuum–igneous, metamorphic and sedimentary rock (3) Slope alluvium–igneous, metamorphic and sedimentary rock
Surface texture	(1) Loam (2) Clay loam (3) Silt loam
Drainage class	Well drained
Permeability class	Moderately slow to rapid
Depth to restrictive layer	20 in
Soil depth	20 in
Surface fragment cover <=3"	0–20%
Surface fragment cover >3"	0–5%
Available water capacity (0-40in)	4.5–8.7 in
Soil reaction (1:1 water) (0-10in)	6.6–7.8
Subsurface fragment volume <=3" (10-20in)	0–40%
Subsurface fragment volume >3" (10-20in)	0–20%

Ecological dynamics

1 - Reference State

1.1 Mid-statured bunchgrasses dominant plant type. Bluebunch wheatgrass tends to be the most common; however, Richardson's needlegrass, or rough fescue and spike fescue (the two species rarely cohabitate) are possible. Minor component of forbs (lupine, balsamroot, arnica) growing with short-statured grasses. Shrubs are common and include grey horsebrush, skunkbush sumac, bearberry, russett buffaloberry, gooseberry, creeping juniper, and common juniper.

1.1a Plant community experiences long term drought, wildfire (low intensity), untimely grazing event

1.2 Mid-statured bunchgrasses share dominance with short bunchgrasses. Shrubs increases as well as forbs likely to increase. Limited tree cover may exist. Shrub canopy increases slightly

1.2a Plant community receives timely moisture and has an opportunity to rest from disturbance

T1A Catastrophic fire, multiple overgrazing events, long term drought, climate change

T1B Overgrazing, Catastrophic fire, introduction of invasive species

2 - Altered State -

2.1 Short-statured grasses take over dominance with shrubs and forbs as a subdominant plant groups. Mid-statured

bunchgrasses rare. Tree presence likely rare but may include Douglas fir, lodgepole pine, subalpine fir, or lesser extent ponderosa pine.

R2A Time and timely moisture, proper grazing management, brush management

T2B Overgrazing, Catastrophic fire, introduction of invasive species


3 - Invaded State

3.1 Site becomes invaded with invasive forbs and grasses. Tree encroachment also often occurs where fire has been suppressed long-term. Bare ground typically high

R3A Removal of invasive species (if possible), proper grazing management, time

T3B Removal of invasive species (if possible), proper grazing management, time

State and transition model



Legend

- **1.1** Mid-statured bunchgrasses dominant plant type. Bluebunch tends to be the most common however Richardson's needlegrass, or Rough fescue/Spike fescue (the two species rarely cohabitate) are possible. Minor component of forbs (Lupine, balsamroot, Arnica) growing with shortgrasses. Shrubs common; including Grey horsebrush, Skunkbush sumac, Bearberry, Russett Buffaloberry, Gooseberry, Creeping Juniper, and/or Common Juniper. 1.1a Plant community experiences long term drought, wildfire (low intensity), untimely grazing event
- **1.2** Mid-statured bunchgrasses share dominance with short bunchgrasses. Shrubs increases as well as forbs likely to increase. Limited tree cover may exist. Shrub canopy increases slightly
- **1.2a** Plant community receives timely moisture and has an opportunity to rest from disturbance
- **2.1** Shortgrasses take over dominance with shrubs and forbs as a subdominant plant groups. Mid-statured bunchgrasses rare. Tree presence likely rare but may include Douglas Fir, Lodgepole Pine, Subalpine fire, or lesser extent Ponderosa Pine.
- **T1A** Catastrophic fire, multiple overgrazing events, long term drought, climate change
- **R1A** Time and timely moisture, proper grazing management, brush management
- **3.1** Site becomes invaded with invasive forbs and grasses. Tree encroachment also often occurs where fire has been suppressed longterm. Bare ground typically high
- **T1B** Overgrazing, Catastrophic fire, introduction of invasive species
- **R3A** Removal of invasive species (if possible), proper grazing management, time
- **T2B** Overgrazing, Catastrophic fire, introduction of invasive species
- **R3B** Removal of invasive species (if possible), proper grazing management, time

Animal community

This site provides wildlife habitat for an array of species. Prior to the settlement of this area, large herds of antelope, elk and bison roamed. Though the bison have been replaced, mostly with domesticated livestock, elk and antelope still frequently utilize this largely intact landscape for winter habitat in areas adjacent to forest. Sites with large quantities of curleaf mountain mahogany are considered important winter range for mule deer, elk, and moose. In some areas it is considered critical habitat for dwindling wild ungulate populations.

Managed livestock grazing is suitable on this site due to the potential to produce an abundance of high quality forage. This is often a preferred site for grazing by livestock, and animals tend to congregate in these areas. In order to maintain the productivity of the site, grazing on adjoining sites with less production must be managed carefully to be sure utilization on this site is not excessive. Management objectives should include maintenance or improvement of the native plant community. Careful management of timing and duration of grazing is important. Shorter grazing periods and adequate deferment during the growing season are recommended for plant maintenance, health, and recovery. According to McLean et al, early season defoliation of bluebunch wheatgrass can result in high mortality and reduced vigor of plants. They also suggest, based on prior studies, that the opportunity for regrowth is necessary before dormancy to reduce injury to bluebunch wheatgrass.

Since needle and thread normally matures earlier than bluebunch wheatgrass and produces a sharp awn this species is usually avoided after seed set. Changing grazing season of use will help utilize needle and thread more efficiently.

Continual non-prescribed grazing of this site will be injurious, will alter the plant composition and production over time, and will result in transition to the Altered State. Transition to other states will depend on duration of poorly managed grazing as well as other circumstances such as weather conditions and fire frequency.

The Altered State is subject to further degradation to the Degraded Short-stature Grass State or Invaded State. Management should focus on grazing management strategies that will prevent further degradation, such as seasonal grazing deferment or winter grazing where feasible. Communities within this state are still stable and healthy under proper management. Forage quantity and quality may be substantially decreased from the Reference State.

Grazing is possible in the Invaded State. Invasive species are generally less palatable than native grasses. Forage production is typically greatly reduced in this state. Due to the aggressive nature of invasive species, sites in the

Invaded State face increased risk for further degradation to the Invasive Dominated Communities. Grazing has to be carefully managed to avoid further soil loss and degradation and possible livestock health issues.

Prescriptive grazing can be used to manage invasive species. In some instances, carefully targeted grazing (sometimes in combination with other treatments) can reduce or maintain species composition of invasive species. In the Degraded Short-statured Grass State, grazing may be possible but is generally not economically or environmentally sustainable.

Hydrological functions

The hydrologic cycle functions best in the Reference State (1) with good infiltration and deep percolation of rainfall; however, the cycle degrades as the vegetation community declines. Rapid rainfall infiltration, high soil organic matter, good soil structure, and good porosity accompany high bunchgrass canopy cover. High ground cover reduces rain drop impact on the soil surface, which keeps erosion and sedimentation transport low. Water leaving the site will have minimal sediment load, which allows for high water quality in associated streams. High rates of infiltration will allow water to move below the rooting zone during periods of heavy rainfall. The Bluebunch Wheatgrass Community (1.1) should have no rills or gullies present and drainage ways should be vegetated and stable. Water flow patterns, if present, will be barely observable. Plant pedestals are essentially non-existent. Plant litter remains in place and is not moved by wind or water.

Improper grazing management results in a community shift to the Mixed Bunchgrass Community (1.2). This plant community has a similar canopy cover, but bare ground will be less than 15 percent. Therefore, the hydrologic cycle is functioning at a level similar to the water cycle in the Bluebunch Wheatgrass Community/Needle and thread (1.1). Compared to the Reference Community (1.1) infiltration rates are slightly reduced and surface runoff is slightly higher.

In the Degraded State (3) and the Invaded State (4) canopy and ground cover are greatly reduced compared to the Reference State (1), which impedes the hydrologic cycle. Infiltration will decrease and runoff will increase due to reduced ground cover, presence of shallow-rooted species, rainfall splash, soil capping, reduced organic matter, and poor structure. Sparse ground cover and decreased infiltration can combine to increase frequency and severity of flooding within a watershed. Soil erosion is accelerated, quality of surface runoff is poor, and sedimentation increases.

Recreational uses

This site is often utilized for photography, hiking, hunting, bird watching, and flower collecting.

Wood products

n/a

Inventory data references

Information presented was derived from NRCS inventory data, literature, field observations, and personal contacts with range-trained personnel (i.e., used professional opinion of agency specialists, observations of land managers, and outside scientists).

Other references

- Barrett, H. 2007. Western Juniper Management: A Field Guide.
- Bestelmeyer, B., J.R. Brown, J.E. Herrick, D.A. Trujillo, and K.M. Havstad. 2004. Land Management in the American Southwest: a state-and-transition approach to ecosystem complexity. *Environmental Management* 34:38–51.
- Bestelmeyer, B. and J. Brown. 2005. State-and-Transition Models 101: A Fresh look at vegetation change.
- Blaisdell, J.P. 1958. Seasonal development and yield of native plants on the Upper Snake River Plains and their relation to certain climate factors.
- Colberg, T.J. and J.T. Romo. 2003. Clubmoss effects on plant water status and standing crop. *Journal of Range Management* 56:489–495.

- DiTomaso, J.M. 2000. Invasive weeds in Rangelands: Species, Impacts, and Management. *Weed Science* 48:255–265.
- Dormaar, J.F., B.W. Adams, and W.D. Willms. 1997. Impacts of rotational grazing on mixed prairie soils and vegetation. *Journal of Range Management* 50:647–651.
- Hobbs, J.R. and S.E. Humphries. 1995. An integrated approach to the ecology and management of plant invasions. *Conservation Biology* 9:761–770.
- Humphrey, L. David. 1984. Patterns and mechanisms of plant succession after fire on *Artemisia*-grass sites in southeastern Idaho *Vegetation*. 57: 91-101.
- Masters, R. and R. Sheley. 2001. Principles and practices for managing rangeland invasive plants. *Journal of Range Management* 38:21–26.
- McLean, A. and S. Wikeem. 1985. Influence of season and intensity of defoliation on bluebunch wheatgrass survival and vigor in southern British Columbia. *Journal of Range Management* 38:21–26.
- Miller, R.F., T.J. Svejcar, and J.A. Rose. 2000. Impacts of western juniper on plant community composition and structure. *Journal of Range Management* 53:574–585.
- Ross, R.L., E.P. Murray, and J.G. Haigh. July 1973. Soil and Vegetation of Near-pristine sites in Montana.
- Smoliak, S., R.L. Ditterlin, J.D. Scheetz, L.K. Holzworth, J.R. Sims, L.E. Wiesner, D.E. Baldrige, and G.L. Tibke. 2006. Montana Interagency Plant Materials Handbook.
- Stavi, I. 2012. The potential use of biochar in reclaiming degraded rangelands. *Journal of Environmental Planning and Management* 55:1–9.
- Stringham, T.K., W.C. Kreuger, and P.L. Shaver. 2003. State and Transition Modeling: an ecological process approach. *Journal of Range Management* 56:106–113.
- Stringham, T.K. and W.C. Krueger. 2001. States, Transitions, and Thresholds: Further refinement for rangeland applications.
- Tirmenstein, D. 1999. *Gutierrezia sarothrae*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). <https://www.fs.fed.us/database/feis/plants/shrub/gutsar/all.html> [2022, March 30].
- Walker, L.R. and S.D. Smith. 1997. Impacts of invasive plants on community and ecosystem properties. Pages 69–86 in *Assessment and management of plant invasions*. Springer, New York, NY.
- Whitford, W.G., E.F. Aldon, D.W. Freckman, Y. Steinberger, and L.W. Parker. 1989. Effects of Organic Amendments on Soil Biota on a Degraded Rangeland. *Journal of Range Management* 41:56–60.
- Wilson, A.M., G.A. Harris, and D.H. Gates. 1966. Cumulative Effects of Clipping on Yield of Bluebunch wheatgrass. *Journal of Range Management* 19:90–91.

Contributors

Petersen, Grant

Approval

Kirt Walstad, 3/01/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/02/2024
Approved by	Kirt Walstad
Approval date	

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-