

Ecological site R043BY366WY

Shallow Sandy (SwSy) 15-19" Foothills and Mountains East Precipitation Zone

Accessed: 05/06/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R043BY308WY	Coarse Upland (CU) 15-19" Foothills and Mountains East Precipitation Zone Coarse Upland
R043BY322WY	Loamy (Ly) 15-19" Foothills and Mountains East Precipitation Zone Loamy
R043BY350WY	Sandy (Sy) 15-19" Foothills and Mountains East Precipitation Zone Sandy
R043BY362WY	Shallow Loamy (SwLy) 15-19" Foothills and Mountains East Precipitation Zone Shallow Loamy

Similar sites

R032XY366WY	Shallow Sandy (SwSy) 10-14" East Precipitation Zone Shallow Sandy 10-14" Foothills and Basins East P.Z., 032XY366WY has lower production
-------------	--

Table 1. Dominant plant species

Tree	Not specified
------	---------------

Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on most slopes and along ridge tops.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Alluvial fan (3) Ridge
Flooding frequency	None
Ponding frequency	None
Elevation	1,829–2,743 m
Slope	20–45%
Ponding depth	0 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 15-19 inches per year. June is generally the wettest month. July, August, and September are somewhat less with daily amounts rarely exceeding one inch.

Snowfall is quite heavy in the area. Annual snowfall averages about 150 inches.

Because of the varied topography, the wind will vary considerably for different parts of the area. The wind is usually much lighter at the lower elevations and in the valleys as compared with the higher terrain. The average winter wind velocity is 8.5 mph while the summer wind velocity averages 7.5 mph. Winds during storms and on ridges may exceed 45 mph.

Growth of native cool-season plants begins about May 1 to May 15 and continues to about October 10.

The following information is from the “Crandall Creek” climate station, at the lower end of this precipitation zone:
 Minimum Maximum 5 yrs. out of 10 between
 Frost-free period (days): 16 80 July 8 – August 20
 Freeze-free period (days): 37 120 June 17 – September 5
 Mean Annual Precipitation (inches): 10.24 21.23

Mean annual precipitation: 14.90 inches

Mean annual air temperature: 38.16 F (21.88 F Avg. Min. to 54.66 F Avg. Max.)

For detailed information, visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. There are no other climate station(s) known to be representative of this precipitation zone.

Table 3. Representative climatic features

Frost-free period (average)	80 days
Freeze-free period (average)	120 days
Precipitation total (average)	483 mm

Influencing water features

Stream type: None

Soil features

The soils of this site are shallow (less than 20" to bedrock) well drained, rapidly permeable and may occur on all slopes. The bedrock may be of any kind except igneous or volcanic and is virtually impenetrable to plant roots. The soil textures are a fine sandy loam or coarser. Thin ineffectual layers of other soil textures are disregarded.

Table 4. Representative soil features

Surface texture	(1) Loamy fine sand (2) Fine sandy loam (3) Sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained to excessively drained
Permeability class	Rapid to very rapid
Soil depth	25–51 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	1.52–3.56 cm
Calcium carbonate equivalent (0-101.6cm)	0–5%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	6.6–7.8
Subsurface fragment volume <=3" (Depth not specified)	0–10%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

Potential vegetation on this site is dominated by mid cool-season perennial grasses. Other significant vegetation includes big sagebrush, antelope bitterbrush and a variety of forbs. On areas along the west slopes of the Big Horn Mountains, mountain mahogany is the significant shrub. The expected potential composition for this site is about 75% grasses, 15% forbs and 10% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates species such as bluegrasses, rhizomatous wheatgrasses and big sagebrush will increase. Cool season grasses such as Columbia needlegrass, spikefescue, bluebunch wheatgrass, and Idaho fescue will decrease in frequency and production. As conditions continue to deteriorate annuals such as cheatgrass will invade.

Big sagebrush and/or juniper and limber pine may become dominant on areas with an absence of fire. Wildfires are actively controlled in recent times and as a result old decadent stands of big sagebrush persist. On some sites, where limber pine and juniper naturally occur and become relatively numerous, an open forest land appearance is possible. This plant community should not be confused with a forest land site, as the site does not historically comprise a significant total production of limber pine and the atypical cover of limber pine is the result of the removal of fire as a natural interaction in plant community succession. Chemical and mechanical controls have replaced the

historic role of fire on this site. Recently, prescribed burning has regained some popularity.

The big sagebrush component may not be as resilient once it has been removed or severely reduced, if a vigorous stand of grass exists and is maintained. The exception to this is where the herbaceous component is severely degraded at the time of treatment, growing conditions are unfavorable after treatment, and/or recovery of herbaceous species are inadequate due to grazing management.


The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.

State and transition model

Site Type: Rangeland
 MLRA: 43BY – Central Rocky Mountains

Shallow Sandy (SwSy) 15"-19" East P.Z.
 RD43BY366WY



- BM** – Brush Management (fire, chemical, mechanical)
- Freq. & Severe Grazing** – Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season
- GLMT** – Grazing Land Mechanical Treatment
- LTPG** – Long-term Prescribed Grazing
- MCSLG** – Moderate, Continuous Season-long Grazing
- HCSLG** – Heavy, Continuous Season-long Grazing
- NU, NF** – No Use and No Fire
- PG** – Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)
- VLTPG** – Very Long-term Prescribed Grazing (could possibly take generations)
- Na** – Moderate Sodium in Soil
- WF** – Wildfire

State 1 Columbia Needlegrass/Idaho fescue Plant Community

Community 1.1 Columbia Needlegrass/Idaho fescue Plant Community

The interpretive plant community for this site is the Historic Climax Plant Community. This state evolved with grazing by large herbivores, shallow soils (less than 20 inches deep), and periodic fires. Potential vegetation is about 75% grasses or grass-like plants, 15% forbs, and 10% woody plants. The cyclical nature of the fire regime in this community prevents big sagebrush and/or juniper and limber pine from being the dominant landscape. This plant community can be found on areas that are properly managed with grazing and/or prescribed burning, and on areas receiving periods of rest. Cool season midgrasses dominate the state. The major grasses include Columbia needlegrass, Idaho fescue, and bluebunch wheatgrass. Big sagebrush and antelope bitterbrush are conspicuous elements of this state, occur in a mosaic pattern, and make up 5 to 10% of the annual production. A variety of forbs also occur in this state and plant diversity is high (see Plant Composition Table). Annual production on this state ranges from 500 to 1000 pounds depending on climatic conditions. This plant community is extremely stable and well adapted to the Central Rocky Mountains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity). Transitions or pathways leading to other plant communities are as follows: • Moderate, continuous season-long grazing will convert the plant community to the Bluebunch Wheatgrass/Mixed Shrub Plant Community.

Figure 4. Plant community growth curve (percent production by month).
WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

State 2 Bluebunch Wheatgrass/Mixed Shrub Plant Community

Community 2.1 Bluebunch Wheatgrass/Mixed Shrub Plant Community

Historically, this plant community evolved under grazing by large ungulates and a low fire frequency. Currently, this site is normally found under a moderate, season-long grazing regime and will be exacerbated by prolonged drought conditions. In addition, the fire regime for this site has been modified and extended periods without fire is now common. Shrubs are important components of this plant community. Cool-season grasses make up the majority of the understory with the balance made up of miscellaneous forbs. Dominant grasses include bluebunch wheatgrass, Idaho fescue, needleandthread, prairie junegrass, mountain brome, and of less frequency Columbia needlegrass. Grasses of secondary importance include rhizomatous wheatgrass, spikefescue, slender wheatgrass, bluegrasses, Indian ricegrass, and upland sedges. Forbs commonly found in this plant community include asters, phlox, hawksbeard, buckwheat, pussytoes, lupine, paintbrush, agoseris, and larkspurs. Sagebrushes and antelope bitterbrush make up from 15% to 20% of the total annual production and to a lesser extent juniper and limber pine will be included. When compared to the Historical Climax Plant Community, sagebrushes and/or juniper and limber pine, rhizomatous wheatgrasses, and bluegrasses have increased. Columbia needlegrass and Idaho fescue have decreased, often occurring only where protected from grazing by the sagebrush canopy. Some weedy species such as cheatgrass may have invaded the site but are in small patches. This state produces between 450 and 900 pounds annually, depending on the growing conditions. This plant community is resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. The herbaceous component is mostly intact and plant vigor and replacement capabilities are sufficient. Water flow patterns and litter movement may be occurring but only on steeper slopes. Incidence of pedestalling is minimal. Soils are mostly stable and the surface shows minimum soil loss. Wind scouring and deposition areas are few. The watershed is functioning and the biotic community is intact. Transitions or pathways leading to other plant communities are as follows: • Prescribed grazing will convert this plant community to the HCPC. The probability of this occurring is high especially if rotational grazing along with short deferred grazing is implemented as part of the prescribed method of use. In addition, the removal of fire suppression will allow a somewhat natural fire regime to reoccur to more easily transition between this plant community and the HCPC. A

prescribed fire treatment can be useful to hasten this transition if desired. • Heavy, continuous, season-long grazing plus no fires will convert the plant community to the Rhizomatous Wheatgrass/Mixed Shrub Plant Community. The probability of this occurring is high. This is especially evident on areas where drought or heavy browsing does not adversely impact the shrub stand. • Heavy, continuous, season-long grazing plus wildfire or brush management will convert the plant community to a Rhizomatous Wheatgrass/Needleandthread Plant Community. The probability for this is high, especially on areas where the shrubs have been heavily browsed or removed by natural or human causes. Drought can also exacerbate this transition.

Figure 5. Plant community growth curve (percent production by month). WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

State 3 Rhizomatous Wheatgrass/Mixed Shrub Plant Community

Community 3.1 Rhizomatous Wheatgrass/Mixed Shrub Plant Community

This plant community currently is found under heavy continuous season-long grazing by livestock and protection from fire. Shrubs are a significant component of this plant community. Cool-season grasses make up the majority of the understory, but some of the preferred grasses have been reduced or absent. Dominant grasses include rhizomatous wheatgrass, needleandthread, mountain brome, prairie junegrass, bluegrasses and of less frequency Columbia needlegrass, Idaho fescue, and bluebunch wheatgrass. Grasses and grasslikes of secondary importance include slender wheatgrass, Indian ricegrass, and upland sedges. Forbs commonly found in this plant community include asters, phlox, hawksbeard, buckwheat, pussytoes, lupine, paintbrush, agoseris, and larkspurs. Big sagebrush and antelope bitterbrush can make up to 30% of the total annual production. On areas along the west slope of the Big Horn Mountains, mountain mahogany replaces big sagebrush as the dominant shrub on this site. When compared to the Historic Climax Plant Community, big sagebrush or mountain mahogany, juniper, limber pine, bluegrasses, prairie junegrass, and rhizomatous wheatgrasses have increased. Most of the preferred grasses and antelope bitterbrush have been reduced and some are absent. Some annuals, such as cheatgrass, have invaded the site, but are not yet abundant. Annual production ranges from 400 to 700 pounds. This plant community is resistant to change as the shrubs become more abundant. These areas may actually be more resistant to fire as less fine fuels are available and the bare ground between the shrubs is increased. The herbaceous component is not as diverse and plant vigor and species regeneration capabilities of some cool-season perennials are deficient. The removal of grazing does not seem to affect the plant composition or structure of the plant community. Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling is more noticeable. Infiltration is reduced and runoff is increased. Rill channels may be noticeable in the interspaces on steeper areas and gullies may be establishing where rills have concentrated down slope. Transitions or pathways leading to other plant communities are as follows: • Prescribed grazing plus brush management or wildfire will convert this plant community to near HCPC. If prescribed fire is used as a means to reduce or remove the shrubs, sufficient fine fuels will need to be present. This may require deferment from grazing prior to treatment. Post management is critical to ensure success. This can range from two or more years of rest to partial growing season deferment, depending on the condition of the understory at the time of treatment and the growing conditions following treatment. Seeding will be required regardless of the brush treatment to reestablish the major cool-season grasses. • Frequent and Severe Grazing plus no fires will convert the plant community to the Mixed Shrub/Bluegrass Plant Community. The probability of this occurring is high and is especially evident on areas where drought or heavy browsing does not adversely impact the shrub stand. • Brush management or Wildfire with no change in grazing management will convert this plant community to the Rhizomatous Wheatgrass/Needleandthread Plant Community.

Figure 6. Plant community growth curve (percent production by month). WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

State 4 Mixed Shrub/Bluegrass Plant Community

Community 4.1 Mixed Shrub/Bluegrass Plant Community

This plant community is the result of frequent and severe grazing and protection from fire. Juniper is the dominant shrub of this plant community, but other shrubs and limber pine will occur and the total production will exceed 30%. The under story has been significantly altered and the preferred cool season grasses have been eliminated or greatly reduced. The dominant grasses are Sandberg, mutton, big, and Canby bluegrasses and upland sedges. Weedy annual species such as cheatgrass, povertyweed, gumweed, and stickseed occupy the site. Juniper is the dominant shrub. Noxious weeds such as Canada thistle and/or leafy spurge may invade the site if a seed source is available. When compared with the HCPC the annual production is less, as the major cool-season grasses are significantly reduced or absent, but the shrub production has increased significantly and compensates for some of the decline in the herbaceous production. The interspaces between plants have expanded leaving the amount of bare ground more prevalent and more soil surface exposed to erosive elements. Annual production ranges from 200 to 500 pounds. This plant community is resistant to change as the stand becomes more decadent. These areas may actually be more resistant to fire as less fine fuels are available and the bare ground between the shrubs is increased. The herbaceous component is not as diverse and plant vigor and species regeneration capabilities of cool-season perennials are deficient. The removal of grazing does not seem to affect the plant composition or structure of the plant community. Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling are obvious. Infiltration is reduced and runoff is increased. Rill channels may be noticeable in the interspaces and gullies may be establishing where rills have concentrated down slope. Transitions or pathways leading to other plant communities are as follows:

- Long-term prescribed grazing will convert this plant community to the Rhizomatous Wheatgrass/ Mixed Shrub Plant Community. This may take many years and the right conditions.
- Brush management or Wildfire with no change in grazing management will convert this plant community to the Bluegrass/Annual Plant Community.
- No fire plus frequent and severe season-long grazing will convert the plant community to the Juniper/Limber Pine Plant Community. This will occur where growing conditions are favorable and a seed source is available for limber pine.

Figure 7. Plant community growth curve (percent production by month).
WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

State 5 Juniper/Limber Pine Plant Community

Community 5.1 Juniper/Limber Pine Plant Community

This plant community is the result of frequent and severe grazing and protection from fire in the Mixed Shrub/Bluegrass plant community. This is likely only where conditions are more conducive and a seed source is available for limber pine. Juniper is the dominant shrub of this plant community, but limber pine can occur in sufficient numbers to give the site an open forestland appearance. The under story has been significantly altered and the preferred cool season grasses have been eliminated or greatly reduced. The dominant grasses are the bluegrasses such as Sandberg, mutton, big, and Canby. Weedy annual species such as cheatgrass, povertyweed, gumweed, and stickseed occupy the site. Juniper is the dominant shrub. Noxious weeds such as Canada thistle and/or leafy spurge may invade the site if a seed source is available. When compared with the HCPC the annual production is less, as the major cool-season grasses are reduced, but the shrub and tree production has increased significantly and compensates for some of the decline in the herbaceous production. The interspaces between plants have expanded leaving the amount of bare ground more prevalent and more soil surface exposed to erosive elements. Annual production ranges from 200 to 500 pounds. This plant community is resistant to change as the stand becomes more decadent. These areas may actually be more resistant to fire as less fine fuels are available and the bare ground between the shrubs is increased. The herbaceous component is not as diverse and plant vigor and species regeneration capabilities of cool-season perennials are deficient. The removal of grazing does not

seem to affect the plant composition or structure of the plant community. Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling are obvious. Infiltration is reduced and runoff is increased. Rill channels are noticeable in the interspaces and gullies may be establishing where rills have concentrated down slope. Transitions or pathways leading to other plant communities are as follows: • Prescribed grazing plus either wildfire or brush management will convert this plant community to near HCPC. If prescribed fire is used as a means to reduce or remove the shrubs, sufficient fine fuels will need to be present. This may require deferment from grazing prior to treatment. Post management is critical to ensure success. This can range from two or more years of rest to partial growing season deferment, depending on the condition of the understory at the time of treatment and the growing conditions following treatment. Seeding will be required regardless of the brush treatment to reestablish the major cool-season grasses. • Brush management or Wildfire with no change in grazing management will convert this plant community to the Bluegrass/ Annual Plant Community.

Figure 8. Plant community growth curve (percent production by month). WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

State 6

Rhizomatous Wheatgrass/Needleandthread Plant Community

Community 6.1

Rhizomatous Wheatgrass/Needleandthread Plant Community

This plant community currently is found under heavy continuous season-long grazing by livestock and is perpetuated by either brush management or a wildfire, which removes shrubs from this plant community. Some of the major cool-season bunchgrasses associated with this ecological site have been reduced and some may have been removed. Dominant grasses include rhizomatous wheatgrasses, needleandthread, bluegrasses, prairie junegrass, nodding brome, and slender wheatgrass, and of less frequency Columbia needlegrass, Idaho fescue, bluebunch wheatgrass, and mountain brome. Forbs commonly found in this plant community include asters, phlox, hawksbeard, buckwheat, pussytoes, lupine, paintbrush, agoseris, and larkspurs. When compared to the Historical Climax Plant Community, rhizomatous wheatgrass, prairie junegrass, Montana wheatgrass, and bluegrasses have increased. Columbia needlegrass, bluebunch wheatgrass, Idaho fescue, big sagebrush and antelope bitterbrush have decreased or been removed. Production of the preferred cool-season grasses has been reduced. Cheatgrass can be common and in large patches, but most of the invaded areas are relatively small. Annual production ranges from 400 to 850 pounds. This plant community is resistant to change as the herbaceous species present are well adapted to grazing. However, species composition can be altered through long-term overgrazing. The herbaceous component is mostly intact, but some cool-season bunchgrasses associated with the site have been reduced or removed. Plant vigor and replacement capabilities are sufficient for some species but not all. Water flow patterns and litter movement is occurring but only on steeper slopes. Incidence of pedestalling is moderate to slight. Soils are mostly stable and the surface shows minimum soil loss. The watershed is functioning and the biotic community is partially intact. Transitions or pathways leading to other plant communities are as follows: • Prescribed grazing will convert this plant community to near HCPC. Reestablishing the big sagebrush and antelope bitterbrush may be difficult and may take many years. Seeding may be required to reestablish any of the lost major bunchgrasses. • Frequent and severe grazing will convert this plant community to a Bluegrass/Annual Plant Community.

Figure 9. Plant community growth curve (percent production by month). WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

State 7

Bluegrass/Annual Plant Community

Community 7.1

Bluegrass/Annual Plant Community

This plant community evolved under frequent and severe heavy grazing and the shrub component has been removed by heavy browsing, wildfire or human means. Weedy annuals and bluegrasses are the most dominant plants and occupy any open bare ground area. The interspaces between plants have expanded leaving the amount of bare ground more prevalent and more soil surface exposed to erosive elements. Compared to the HCPC, weedy annual species and bluegrasses are widespread and virtually all of the major cool-season mid-grasses are absent or severely decreased. Big sagebrush has also been removed. The dominant grasses are the bluegrasses such as Sandberg, mutton, big, and Canby. Weedy annual species such as cheatgrass, povertyweed, gumweed, and stickseed occupy the site. Noxious weeds have usually invaded the site if a seed source is available. Annual production ranges from 100 to 500 pounds. This plant community is relatively stable and resistant to overgrazing. Annuals and bluegrasses are effectively competing against the establishment of perennial cool-season grasses. Plant diversity is greatly altered and the herbaceous component is not intact. Recruitment of the major perennial grasses is not occurring and the replacement potential is absent. The biotic integrity is missing. The soils are unstable and not protected from excessive erosion. Rill channels and maybe even gullies may be present on site and adjacent areas are impacted by excessive runoff. Water flow patterns and pedestalling are obvious. The watershed is not functioning. Transitions or pathways leading to other plant communities are as follows: • Long Term Prescribed Grazing will convert this plant community to the Rhizomatous Wheatgrass/Needleandthread Plant community.

Figure 10. Plant community growth curve (percent production by month). WY0601, 15-19E all upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			5	15	40	20	10	10			

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1				95–239	
	Columbia needlegrass	ACNE9	<i>Achnatherum nelsonii</i>	95–239	–
2				95–239	
	Idaho fescue	FEID	<i>Festuca idahoensis</i>	95–239	–
3				48–95	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	48–95	–
4				0–95	
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–95	–
5				0–95	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–95	–
6				0–95	
	mountain brome	BRMA4	<i>Bromus marginatus</i>	0–95	–
7				95–191	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–48	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	0–48	–
	nodding brome	BRAN	<i>Bromus anomalus</i>	0–48	–
	Pumpelly's brome	BRINP5	<i>Bromus inermis ssp. pumpellianus var. pumpellianus</i>	0–48	–
	sedge	CAREX	<i>Carex</i>	0–48	–
	thickspike	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	0–48	–

	wheatgrass				
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0–48	–
	spike fescue	LEKI2	<i>Leucopoa kingii</i>	0–48	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–48	–
Forb					
8				48–143	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–48	–
	yarrow	ACHIL	<i>Achillea</i>	0–48	–
	rosy pussytoes	ANRO2	<i>Antennaria rosea</i>	0–48	–
	Indian paintbrush	CASTI2	<i>Castilleja</i>	0–48	–
	field chickweed	CEAR4	<i>Cerastium arvense</i>	0–48	–
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	0–48	–
	larkspur	DELPH	<i>Delphinium</i>	0–48	–
	fleabane	ERIGE2	<i>Erigeron</i>	0–48	–
	sulphur-flower buckwheat	ERUM	<i>Eriogonum umbellatum</i>	0–48	–
	aster	EUCEP2	<i>Eucephalus</i>	0–48	–
	lupine	LUPIN	<i>Lupinus</i>	0–48	–
	beardtongue	PENST	<i>Penstemon</i>	0–48	–
	phlox	PHLOX	<i>Phlox</i>	0–48	–
Shrub/Vine					
9				0–48	
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	0–48	–
10				0–48	
	mountain mahogany	CERCO	<i>Cercocarpus</i>	0–48	–
11				0–48	
	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	0–48	–
12				0–48	
	Rocky Mountain juniper	JUSC2	<i>Juniperus scopulorum</i>	0–48	–
13				0–48	
	limber pine	PIFL2	<i>Pinus flexilis</i>	0–48	–
14				0–48	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–48	–

Animal community

Animal Community – Wildlife Interpretations

Columbia Needlegrass/Idaho fescue Plant Community (HCPC): The predominance of grasses in this plant community favors grazers and mixed-feeders, such as deer, bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. Due to the location of these sites on the foot slopes of mountains they are valuable for elk and deer winter ranges. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlark, lark bunting, sage thrasher, horned larks, red-tail and ferruginous hawks, and golden eagles. Many grassland obligate small mammals would occur here.

Bluebunch Wheatgrass/Mixed Shrub Plant Community: The combination of an overstory of big sagebrush and an

understory of grasses and forbs provides a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer, elk, and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants and hosts of other nesting birds utilize stands in the 20-30% cover range. Other birds that would frequent this plant community include western meadowlark, lark bunting, sage thrasher, horned larks, red-tail and ferruginous hawks, and golden eagles.

Rhizomatous Wheatgrass/Mixed Shrub Plant Community: The combination of an overstory of big sagebrush and an understory of grasses and forbs provides a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer, elk, and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants and hosts of other nesting birds utilize stands in the 20-30% cover range. Other birds that would frequent this plant community include western meadowlark, lark bunting, sage thrasher, horned larks, red-tail and ferruginous hawks, and golden eagles.

Mixed Shrub/Bluegrass Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15% protein and 40-60% digestibility during that time. This community provides escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for sage grouse. Other birds that would frequent this plant community include western meadowlark, lark bunting, sage thrasher, horned larks, red-tail and ferruginous hawks, and golden eagles. Due to the lack of herbaceous production and diversity of mid cool season grasses on this site, it is not as beneficial to grazers.

Juniper/Limber Pine Plant Community: This plant community can provide important winter and escape cover for elk, mule deer and antelope, as the juniper and limber pine can approach 70% cover. However, due to the lack of quality browsing and herbaceous species, this site provides only a minimal source of forage for most wildlife species. Specific bird species such as the nuthatches, western tanager, western kingbird, mountain bluebird, wood-warblers, and northern flicker frequent this site.

Rhizomatous Wheatgrass/Needleandthread Plant Community: The production of herbaceous species provided for good foraging for grazers. However, the lack of tall or mid growing shrubs does not benefit browsers nor provides cover for many wildlife species. As these site greens-up sooner in the spring, this site tends to provide early new growth for foraging large and small mammals. If located adjacent to shrub dominated sites, It provides good foraging habitat for sage grouse.

Bluegrass/Annual Plant Community: This community provides limited foraging for elk and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover. Generally, these are not target plant communities for wildlife habitat management.

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community Production Carrying Capacity*

(lb./ac) (AUM/ac)

Columbia Needlegrass/Idaho fescue 500-1000 .4

Bluebunch WG/Mixed Shrub 450-900 .35

Rhizomatous WG/ Mixed Shrub 400-700 .3

Mixed Shrub/Bluegrass 200-500 .15

Juniper/Limber Pine 200-500 .1

Rhizomatous WG/Needleandthread 400-850 .35

Bluegrass/Annual 100-500 .1

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Climate is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group D due to its shallow feature. Infiltration and runoff potential for this site varies from moderate to high depending on soil hydrologic group and water table. Runoff will be high on this site since the soil saturates easily and due to its shallow characteristic and water holding capacity. (Refer to Part 630, NRCS National Engineering Handbook for detailed hydraulic information.)

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides hunting opportunities for upland game species. The wide varieties of plants that bloom from spring until fall have an esthetic value that appeals to visitors. Other recreational uses may include hiking, camping, and mountain biking.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, James Haverkamp, Range Management Specialist, Steven Gullion, Range Management Specialist, James Mischke, District Conservationist, and Everet Bainter, State Range Management Specialist. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Contributors

J. Haverkamp

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Ray Gullion, Everet Bainter
--------------------------	-----------------------------

Contact for lead author	ray.gullion@wy.usda.gov 307-347-2456
Date	05/30/2008
Approved by	E. Bainter
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills should not be present

2. **Presence of water flow patterns:** Barely observable

3. **Number and height of erosional pedestals or terracettes:** Essentially non-existent

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 35-55% occurring in small areas throughout site

5. **Number of gullies and erosion associated with gullies:** Active gullies should be restricted to areas of concentrated water flow patterns on steeper slopes

6. **Extent of wind scoured, blowouts and/or depositional areas:** Small scoured sites may be observed

7. **Amount of litter movement (describe size and distance expected to travel):** Litter movement is little to none based on topography and water flow patterns

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Plant cover and litter is at 55% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 3 or greater.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use Soil Series description for depth and color of A-horizon

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur. Infiltration is rapid to very rapid

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be**

mistaken for compaction on this site): No compaction layer or soil surface crusting should be present.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Mid stature Cool Season Grasses > Forbs > Shrubs > Short Grasses/Grasslikes

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very Low
-

14. **Average percent litter cover (%) and depth (in):** Average litter cover is 20-30% with depths of 0.25 to 0.5 inches
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 850 lbs/acre
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Bare ground greater than 50% is the most common indicator of a threshold being crossed. Big sagebrush and bluegrasses are common increasers. Annual weeds such as cheatgrass and mustards are common invasive species in disturbed sites.
-

17. **Perennial plant reproductive capability:** All species are capable of reproducing, except in extreme drought years.
-