

Ecological site R043CY807OR Scabland (PSSPS-POSE-DAUN)

Last updated: 9/08/2023 Accessed: 05/06/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 043C-Blue and Seven Devils Mountains

This MLRA covers the Blue and Seven Devils Mountains of Oregon, Washington and Idaho. The area is characterized by thrust and block-faulted mountains and deep canyons composed of sedimentary, metasedimentary, and volcanic rocks. Elevations range from 1,300 to 9,800 feet (395 to 2,990 meters). The climate is characterized by cold, wet winters and cool, dry summers. Annual precipitation, mostly in the form of snow, averages 12 to 43 inches (305 to 1,090 millimeters) yet ranges as high as 82 inches (2,085 millimeters) at upper elevations. Soil temperature regimes are predominately Frigid to Cryic and soil moisture regimes are predominately Xeric to Udic. Mollisols and Andisols are the dominant soil orders. Ecologically, forests dominate but shrub and grass communities may occur on south aspects and lower elevations as well as in alpine meadow environments. Forest composition follows moisture, temperature and elevational gradients and typically ranges from ponderosa pine and Douglas-fir plant associations at lower elevations, grand fir at middle elevations and subalpine fir and Engelman spruce at upper elevations. Historical fire regimes associated with these forest types range from frequent surface fires in ponderosa pine - Douglas Fir forest types to mixed and stand replacing fire regimes in grand fir and subalpine fir types. A large percentage of the MLRA is federally owned and managed by the U.S. Forest Service for multiple uses.

Classification relationships

Plant Assoc. of Blue and Ochoco Mountains (R6 E TP-036-92) bluebunch wheatgrass-Sandberg bluegrass-onespike oatgrass - GB4911 Sandberg bluegrass-onespike oatgrass - GB9111

USDA Forest Service Ecological Sub-region M332 "Blue Mountains"

U.S. National Vegetation Classification Standard (NVCS) Group: G521. Vancouverian-Rocky Mountain Montane Wet Meadow & Marsh Alliance: A3814. Danthonia spp. - Camassia spp. Wet Meadow Alliance Association: Danthonia unispicata - *Poa secunda* Wet Meadow

Group: G273. Central Rocky Mountain Lower Montane, Foothill & Valley Grassland Alliance: A3987. Festuca idahoensis - Pseudoroegneria spicata - *Poa secunda* Dry Grassland Alliance Association(s): Pseudoroegneria spicata - *Poa secunda* Grassland Pseudoroegneria spicata - *Poa secunda* Lithosolic Grassland

Ecological site concept

This site occurs on upland plateaus with very shallow to shallow soils over hard igneous extrusive geologies

(basalt, andesite, rhyolite, tuff, etc.). In the reference state, the site is characterized by bluebunch wheatgrass (Pseudoroegneria spicata) and Sandberg bluegrass (*Poa secunda*). This is a low production site. Production is limited by the low water holding capacity of the soil. Precipitation comes in the form of snow and rain primarily in the winter and early spring. The soil profile is not able to store all the moisture that the site receives, and excess moves to adjacent sites. In areas where the entire soil profile stays wet into the late spring, one-spike oatgrass (Danthonia unispicata) is common. Historically, plant community dynamics were driven primarily by disturbances such as localized fire and drought.

This is a provisional ecological site that groups characteristics at a broad scale with little to no field verification and is subject to extensive review and revision before final approval. All data herein was developed using existing information and literature and should be considered provisional and contingent upon field validation prior to use in conservation planning.

Associated sites

R043CY809OR	Warm Foothills and Mountains (PSSPS-POSE)
	Deeper soils on adjacent landforms

Similar sites

R043CY806OR	Very Shallow Scabland (ARRI2/PSSPS-POSE) Influenced by a continental climate thus has a greater shrub (ARRI2) component	
	Claypan (ARAR8/FEID-PSSPS) Soils characterized by an abrupt textural change to a claypan within the upper horizo	

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) Pseudoroegneria spicata(2) Poa secunda

Physiographic features

This site typically occurs on benches, plateaus and gently sloping areas over hard igneous extrusive geologies (basalt, andesite, rhyolite, tuff, etc.). Slopes range from 0 to 20 %. Elevations are typically 4,000 - 5,000 feet (1,225 - 1,525 meters), but may range from 3,350 to 5,800 (1,025 to 1,775). This site occurs on all aspects. This site does not experience ponding or flooding and no water table is present within the upper two meters of soil.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Plateau(2) Mountains > Bench
Flooding frequency	None
Ponding frequency	None
Elevation	4,000–5,000 ft
Slope	0–20%
Ponding depth	0 in
Water table depth	100 in
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency Not specified

Ponding frequency	Not specified	
Elevation	3,350–5,800 ft	
Slope	Not specified	
Ponding depth	Not specified	
Water table depth	Not specified	

Climatic features

The annual precipitation ranges from 18 - 25 inches (650 to 1,950 mm). It occurs mainly between the months of October and June in the form of rain and snow. Localized convection storms occasionally occur during the summer but resulting precipitation does not generally have a significant effect on soil moisture. The soil temperature regime is frigid. Temperature extremes range from 90 to -20 degrees F (32 to -28 degrees C). The frost free period ranges from less than 30 up to 90 days. The optimum growth period for plant growth is late April through June. Climate graphs are populated from the closest available weather stations and are included to represent general trends rather than representative values.

Table 4. Representative climatic features

Frost-free period (characteristic range)	30-90 days
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	18-25 in
Frost-free period (average)	45 days
Freeze-free period (average)	
Precipitation total (average)	23 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) ENTERPRISE 20 NNE [USC00352678], Enterprise, OR
- (2) ELGIN [USC00352597], Elgin, OR
- (3) MEACHAM [USW00024152], Pendleton, OR

Influencing water features

This site is not influenced by water from a wetland or stream.

Soil features

The soils of this site are very shallow to shallow over basalt bedrock and are well drained. Areas of rubbleland may occur. Typically, the surface texture extremely cobbly silt loam but it may vary. The subsoil is typically clay loam or clay. The family particle size is typically loamy-skeletal but may also be clayey-skeletal. These soils formed in colluvium and residuum derived from hard igneous extrusive rocks and often have a minor influence of volcanic ash in the surface. Total coarse fragments throughout the profile range from 15 to 60 percent by volume, typically averaging over 35%. The potential for water or wind erosion is moderate. See Bocker and Anatone for modal series concepts.

Table 5. Representative soil features

Parent material	(1) Colluvium–volcanic rock(2) Residuum–volcanic rock
Surface texture	(1) Extremely cobbly silt loam(2) Very cobbly clay loam(3) Very stony silt loam
Family particle size	(1) Loamy-skeletal (2) Clayey-skeletal
Drainage class	Well drained
Permeability class	Moderate to moderately slow
Depth to restrictive layer	0–20 in
Soil depth	0–20 in
Surface fragment cover <=3"	10–45%
Surface fragment cover >3"	10–30%
Available water capacity (0-20in)	0.6–1.2 in
Soil reaction (1:1 water) (0-20in)	6.1–7.2
Subsurface fragment volume <=3" (4-20in)	10–45%
Subsurface fragment volume >3" (4-20in)	5–40%

Ecological dynamics

Range in Characteristics:

Variability in plant composition and yield is dependent on soil depth and bedrock fracture rather than on precipitation and elevation ranges that occur within the site. The highest yield and most bluebunch wheatgrass occurs over fractured bedrock and on soils 6 to 10+ inches deep. Sandberg bluegrass increases over unfractured bedrock and soils that are less than 4 inches deep.

Ecological Dynamics:

Ecological dynamics of this site are primarily driven by interactions between climatic patterns and disturbance regimes. Infrequent and typically small area fires were the historical disturbance that maintained the reference state and drove plant community shifts within the state. Intensity and frequency of these fires is strongly influenced by

drought cycles. Introduction of exotic annual grasses compromises the resistance and resiliency of the site, putting it at higher risk of crossing a threshold into another state.

Periodic drought regularly influences these ecosystems and drought duration and severity has increased throughout the 20th century in much of the Intermountain West. Major shifts away from historical precipitation patterns have the greatest potential to alter ecosystem function and productivity. Species composition and productivity can be altered by the timing of precipitation and water availability within the soil profile (Bates et al. 2006).

The invasibility of plant communities is often linked to resource availability. Disturbance can decrease resource uptake due to damage or mortality of the native species and depressed competition, or can increase resource pools by the decomposition of dead plant material following disturbance. The invasion of sagebrush communities by cheatgrass (*Bromus tectorum*) has been linked to disturbances (fire, abusive grazing) that have resulted in fluctuations in resources (Chambers et al. 2007).

The species most likely to invade these sites are cheatgrass, medusahead (*Taeniatherum caput-medusae*), and North Africa grass (*Ventenata dubia*); cool-season annual grasses that germinate in the fall, overwinter as seedlings, and initiate growth in the spring (Miller et al. 1999a). Expansion of these grasses creates seed reserves that can infest adjoining areas and cause changes to the fire regime. These grasses create fuel continuity in the understory, often resulting in increased fire frequency and more uniform burn patterns (less mosaic).


Fire Ecology:

Fire will remove aboveground biomass from bluebunch wheatgrass but plant mortality is generally low (Robberecht and Defossé 1995) because the buds are underground (Conrad and Poulton 1966) or protected by foliage. Uresk et al. (1976) reported burning increased vegetative and reproductive vigor of bluebunch wheatgrass. Thus, while bluebunch wheatgrass may experience slight damage to fire, it is more susceptible in drought years (Young 1983). Plant response will vary depending on season, fire severity, fire intensity and post-fire soil moisture availability.

Sandberg bluegrass (*Poa secunda*), has been found to increase following fire likely due to its low stature and productivity (Daubenmire 1975). Sandberg bluegrass competition may retard reestablishment of deeper rooted bunchgrass.

(Adapted from: Stringham, T.K. et al., 2016)

State and transition model



Reference State 1.0 Community Pathways

- 1.1a: Low severity fire creates grass/forb mosaic
- 1.2a: Time and lack of disturbance allows for perennial grasses to increase.

Transition T1A: Introduction of non-native species

Current Potential State 2.0 Community Pathways

2.1a: Low severity fire creates grass/forb mosaic

2.2a: Time and lack of disturbance allows for perennial grasses to increase.

Transition T2A: Severe or continuous disturbance in the presence of non-native annual species, may be combined with higher than normal spring precipitation. (e.g. Catastrophic fire and/or soil disturbing treatments and/or inappropriate grazing management resulting in soil compaction due to grazing while soils are wet and/or utilization of perennial grasses that impact the plant's ability to recover)

State 1 Historical reference state

The Reference State 1.0 is representative of the natural range of variability under pristine conditions. The reference state is bunchgrass dominated with a diverse forb component. State dynamics are maintained by interactions between climatic patterns and disturbance regimes. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These are maintained by elements of ecosystem structure and function such as the presence of all structural and functional groups, and retention of organic matter and nutrients. Plant community phase changes are primarily driven by fire and/or periodic drought.

Dominant plant species

- bluebunch wheatgrass (Pseudoroegneria spicata ssp. spicata), grass
- Sandberg bluegrass (Poa secunda), grass
- onespike danthonia (Danthonia unispicata), grass

State 2 Current Potential State

This state is similar to the Reference State 1.0. Ecological function has not changed, however the resiliency of the state has been reduced by the presence of invasive weeds. Non-native plant species may increase in abundance but will not become dominant within this State. These species can be highly flammable and can promote fire where

historically fire had been infrequent. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These are maintained by elements of ecosystem structure and function such as the presence of all structural and functional groups, and retention of organic matter and nutrients. Positive feedbacks decrease ecosystem resilience and stability of the state. Processes and characteristics that contribute to positive feedbacks include the non-natives species' high seed output, persistent seed bank, rapid growth rate, ability to cross pollinate, and adaptations for seed dispersal.

Dominant plant species

- bluebunch wheatgrass (Pseudoroegneria spicata ssp. spicata), grass
- Sandberg bluegrass (Poa secunda), grass
- onespike danthonia (Danthonia unispicata), grass

State 3 Exotic annual grass state

This community is characterized by the dominance of annual non-native species such as medusahead, cheatgrass, and ventenata.

Dominant plant species

- cheatgrass (Bromus tectorum), grass
- North Africa grass (Ventenata dubia), grass
- medusahead (Taeniatherum caput-medusae), grass

Transition T1A State 1 to 2

Introduction of non-native species

Transition T2A State 2 to 3

Severe or continuous disturbance in the presence of non-native annual species, may be combined with higher than normal spring precipitation. (e.g Catastrophic fire and/or soil disturbing treatments and/or inappropriate grazing management resulting in soil compaction due to grazing while soils are wet and/or utilization of perennial grasses that impact the plant's ability to recover).

References

Johnson Jr, C.G. and R.R. Clausnitzer. 1992. Plant Association of the Blue and Ochoco Mountains. Tech. Publ. R6-ERW-TP-036-92.. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallow-Whitman National Forest, Portland, OR.

Johnson, C.G. and S.A. Simon. 1987. Plant Association of the Walla-Snake Province, Wallowa-Whitman National Forest. R6-ECOL-TP-225A-86.. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallow-Whitman National Forest.

Other references

Bates, J.D., T. Svejcar, R.F. Miller and R.A. Angell. 2006. The effects of precipitation timing on sagebrush steppe vegetation. Journal of Arid Environments 64 (2006): 670-697.

Chambers, J., B. Bradley, C. Brown, C. D'Antonio, M. Germino, J. Grace, S. Hardegree, R. Miller, and D. Pyke. 2013. Resilience to Stress and Disturbance, and Resistance to *Bromus tectorum* L. Invasion in Cold Desert Shrublands of Western North America. Ecosystems:1-16.

Conrad, C. E. and C. E. Poulton. 1966. Effect of a wildfire on Idaho fescue and bluebunch wheatgrass. Journal of Range Management:138-141.

Daubenmire, R.F. 1975. Plant succession on abandoned fields, and fire influences, in a steppe area in southeastern Washington. Northwest Science 49(1):36-48.

Miller, H.C., D. Clausnitzer, and M.M Borman. 1999a. Medusahead. In: Sheley, R.L. and J.K. Petroff (eds.). Biology and Management of Noxious Rangeland Weeds. Oregon State University Press, Corvallis, OR.

Robberecht, R. and G. Defossé. 1995. The relative sensitivity of two bunchgrass species to fire. International Journal of Wildland Fire 5:127-134.

Stringham, T.K., D. Snyder, and A. Wartgow. 2016. State-and-Transition Models for USFS Crooked River National Grassland Major Land Resource Area B10 Oregon. DRAFT Report. University of Nevada Reno.

Uresk, D. W., J. F. Cline, and W. H. Rickard. 1976. Impact of wildfire on three perennial grasses in south- central Washington. Journal of Range Management 29:309-310.

USNVC [United States National Vegetation Classification]. 2020. United States National Vegetation Classification Database, V2.03. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. [http://usnvc.org/ accessed 9/25/2020]

Young, R.P. 1983. Fire as a vegetation management tool in rangelands of the Intermountain region. In: Monsen, S.B. and N. Shaw (Eds). Managing Intermountain rangelands—improvement of range and wildlife habitats: Proceedings of symposia; 1981 September 15-17; Twin Falls, ID; 1982 June 22-24; Elko, NV. Gen. Tech. Rep. INT-157. Ogden, UT. U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. Pp. 18-31.

Contributors

Jennifer Moffitt - Original concept developed for 2020 PES initiative Andrew Neary - Further concept development for 2020 PES initiative

Approval

Kirt Walstad, 9/08/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/06/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: