

Ecological site R046XN589MT

Shallow Clay (SwC) RRU 46-N 13-16 PZ

Last updated: 7/19/2023
Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R046XN247MT	Clayey (Cy) RRU 46-N 13-19 PZ
R046XN250MT	Shallow (Sw) RRU 46-N 13-19 PZ
R046XN594MT	Silty Steep (SiStp) RRU 46-N 13-19 PZ

Similar sites

R046XN250MT	Shallow (Sw) RRU 46-N 13-19 PZ The Shallow site differs by having a different texture, and generally being over different parent materials/bedrock.
R046XN247MT	Clayey (Cy) RRU 46-N 13-19 PZ Clayey sites have similar textures, but differ mainly by being over 20 inches to rock (shale), and having significantly more production. The plant community can be similar because of the clayey textures.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified

Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Festuca campestris</i>
------------	---

Physiographic features

This ecological site can occur on nearly level to very steep uplands. It often occurs in complex with other ecological sites, particularly in rougher terrain. This site occurs on all slopes and exposures and aspect sometimes becomes significant. Variations in plant community composition and production can result due to aspect. The amount of exposed rock outcrop tends to increase as slopes increase. Runoff and the potential for water erosion can be important features of this site.

Table 2. Representative physiographic features

Landforms	(1) Plain (2) Hill (3) Escarpment
Flooding frequency	None
Ponding frequency	None
Slope	0–70%
Water table depth	152 cm
Aspect	Aspect is not a significant factor

Climatic features

See Climatic Data Sheet for more details (Section II of the Field Office Technical Guide) or reference the following climatic web site: <http://www.wrcc.sage.dri.edu/>.

Table 3. Representative climatic features

Frost-free period (characteristic range)	57-84 days
Freeze-free period (characteristic range)	109-120 days
Precipitation total (characteristic range)	381-432 mm
Frost-free period (actual range)	37-90 days
Freeze-free period (actual range)	101-122 days
Precipitation total (actual range)	356-432 mm
Frost-free period (average)	69 days
Freeze-free period (average)	114 days
Precipitation total (average)	406 mm

Climate stations used

- (1) BABB 6 NE [USC00240392], Babb, MT
- (2) AUGUSTA [USC00240364], Augusta, MT
- (3) CASCADE 5 S [USC00241552], Cascade, MT
- (4) ROGERS PASS 9 NNE [USC00247159], Wolf Creek, MT

Influencing water features

No influencing water features.

Soil features

These soils form on alluvium and colluvium over residuum of semiconsolidated or consolidated shale. They are clayey soils that are 10 to 20 inches deep to underlying shale or nearly impervious clays. Few roots penetrate deeper than 20 inches.

Table 4. Representative soil features

Surface texture	(1) Clay loam (2) Silty clay loam (3) Silty clay
Drainage class	Well drained to somewhat excessively drained
Permeability class	Slow to very slow
Soil depth	25–51 cm
Available water capacity (0–101.6cm)	5.08–10.16 cm
Electrical conductivity (0–101.6cm)	0 mmhos/cm
Subsurface fragment volume ≤3" (Depth not specified)	0–5%
Subsurface fragment volume >3" (Depth not specified)	0–5%

Ecological dynamics


This site developed under Northern Rocky Mountain foothills climatic conditions, which included the natural influence of large herbivores and occasional fire. The plant community upon which interpretations are primarily based is the Historic Climax Plant Community (HCPC). The Historic Climax Plant Community is described as a reference to understand the original potential of this site, and is not considered to be the management goal for every acre of rangeland. The following descriptions should enable the landowner/ manager to better understand which plant communities occupy their land, and assist with setting goals for vegetation management. It can also be useful to understand the environmental and economic values of each plant community.

This site is considered moderately resilient to disturbance as it has only moderate soil limitations for plant growth. Changes may occur to the Historic Climax Plant Community due to management actions and/or climatic conditions. Under continued adverse impacts, a moderate decline in vegetative vigor and composition will occur. Under favorable vegetative management treatments, this site can more readily return to the Historic Climax Plant Community (HCPC).

Continual adverse impacts to the site over a period of years results in a departure from the HCPC, with a decrease of the taller, more palatable species such as rough fescue, bluebunch wheatgrass, plains muhly, and the taller needlegrasses. These plants will be replaced by Idaho fescue, western or thickspike wheatgrass, Sandberg bluegrass, plains reedgrass, threadleaf sedge, various increaser forbs, and mountain big sagebrush. Continued deterioration results in increased amounts of fringed sagewort, threeawns, and weedy forbs. Rubber rabbitbrush may also increase at some locations.

Plants that are not a part of the reference community that are most likely to invade are annuals, broom snakeweed, and thistles. There are several noxious weeds that are also likely to invade this site including spotted knapweed, leafy spurge, dalmation toadflax, and sulphur cinquefoil.

State and transition model



Smaller boxes within a larger box indicate that these communities will normally shift among themselves with slight variations in precipitation and other disturbances. Moving outside the larger box indicates the community has crossed a threshold (heavier line) and will require intensive treatment to return to Community 1 or 2. Dotted lines indicate a reduced probability for success. Yellow boxes indicate caution that the community may be in danger of crossing a threshold. Orange boxes represent communities that have crossed over thresholds from the HCPC and may be difficult to restore with grazing management alone. Red boxes represent communities that have severely shifted away from the HCPC and probably cannot be restored without mechanical inputs.

NOTE: Not all species present in the community are listed in this table. Species listed are representative of the plant functional groups that occur in the community.

PG = Prescribed Grazing: Use of a planned grazing strategy to balance animal forage demand with available forage resources. Timing, duration, and frequency of grazing are controlled and some type of grazing rotation is applied to allow for plant recovery following grazing.

NPG = Non-Prescribed Grazing: Grazing which has taken place that does not control the factors as listed above, or animal forage demand is higher than the available forage supply.

Figure 8. State and Transition Model

State 1 Tall and Medium Grasses, Forbs, Half-Shrub

Community 1.1 Tall and Medium Grasses, Forbs, Half-Shrub

This is the interpretive plant community and is considered to be the Historic Climax Plant Community (HCPC) for this site. This plant community contains a high diversity of tall and medium height, cool and warm season grasses

(bluebunch wheatgrass, rough fescue, green or Columbia needlegrasses, thickspike or western wheatgrass and, plains muhly), and short grasses and sedges (Idaho fescue, Cusick bluegrass, Sandberg bluegrass, plains reedgrass, threadleaf and needleleaf sedge). There are abundant forbs (prairie clovers, dotted gayfeather) which occur in smaller percentages. Half shrubs such as winterfat should also be common. Shrubs such as mountain big sagebrush can also be present. This plant community is well adapted to the Northern Rocky Mountain foothills climatic conditions. The diversity in plant species allows for drought tolerance. Individual species can vary greatly in production depending on growing conditions (timing and amount of precipitation, and temperature). This plant community is well suited to managed livestock grazing and provides diverse habitat for many wildlife species. Plants on this site have strong, healthy root systems that allow production to increase significantly with favorable moisture conditions. This plant community provides for soil stability and a properly functioning hydrologic cycle. Abundant plant litter is available for soil building and moisture retention. Plant litter is properly distributed with very little movement off-site and natural plant mortality is very low. The soils associated with this site provide a limited soil-water-plant relationship.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1048	1191	1334
Forb	123	140	157
Shrub/Vine	—	37	78
Total	1171	1368	1569

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	5-10%
Grass/grasslike foliar cover	20-40%
Forb foliar cover	1-5%
Non-vascular plants	0-1%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Soil surface cover

Tree basal cover	0%
Shrub/vine/liana basal cover	0-2%
Grass/grasslike basal cover	14-21%
Forb basal cover	1-2%
Non-vascular plants	0-1%
Biological crusts	0%
Litter	50-60%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	5-15%
Bedrock	0%

Water	0%
Bare ground	10-20%

State 2

Medium and Short Grasses, Sedge, and Increaser Forbs

Community 2.1

Medium and Short Grasses, Sedge, and Increaser Forbs

Early stages of degradation, including non-prescribed grazing, will tend to change the HCPC to a community dominated by medium and short grasses and sedges such as Idaho fescue, thickspike/western wheatgrass, Sandberg bluegrass, threadleaf sedge, Cusick bluegrass, and prairie junegrass. Bluebunch wheatgrass can still be relatively abundant. Most of the other taller and more palatable plants (rough fescue, tall needlegrasses, winterfat) will still be present but in smaller amounts. There may be an increase in the amount of some shrubs, particularly mountain big sagebrush. Palatable and nutritious forbs will be replaced by less desirable and more aggressive species such as fringed sagewort. This plant community will readily respond to improved grazing management, but a significant amount of time can be necessary to move it toward a higher successional stage and a more productive plant community similar to community 1. Biomass production and litter become slightly reduced on the site with Community 2 as the taller grasses become replaced by shorter ones. Evapotranspiration tends to increase, moisture retention is reduced, and soil surface temperatures increase. Some natural ecological processes will be altered. These plant communities provide for moderate soil stability. Increased amounts of bare ground can result in undesirable species invading. Common invaders can include spotted knapweed, leafy spurge, dalmatian toadflax, and sulphur cinquefoil.

State 3

Shrub, Short and Mid Increaser Grasses, , Fringed Sagewort, Increaser Forbs Cactu

Community 3.1

Shrub, Short and Mid Increaser Grasses, , Fringed Sagewort, Increaser Forbs Cactu

With continued heavy disturbance, the site will become dominated by species such as mountain big sagebrush, short and medium increaser grasses such as Sandberg bluegrass, plains reedgrass, prairie junegrass, western or thickspike wheatgrass, and Idaho fescue, fringed sagewort, and increaser forbs such as scarlet globemallow and Hood's phlox. There may still be remnant amounts of some of the late-seral species such as bluebunch wheatgrass, rough fescue, and green/Columbia needlegrass present. The taller grasses will occur only occasionally, often under sagebrush plants. Palatable forbs will be mostly absent. Plains pricklypear cactus and rubber rabbitbrush may become common. Plant community 3 is often less productive than 1 or 2. The lack of litter and short plant heights result in higher soil temperatures, poor water infiltration rates, and higher evapotranspiration rates, thus eventually favoring species that are more adapted to drier conditions. These communities have lost many of the attributes of a healthy rangeland, including good infiltration, minimal erosion and runoff, nutrient cycling and energy flow. This plant community is the result of long-term, heavy, continuous grazing and/or annual, early spring seasonal grazing. Repeated spring grazing depletes stored carbohydrates, resulting in weakening and eventual death of the cool season tall and medium grasses. This plant community can occur throughout the pasture, on spot grazed areas, and around water sources where season-long grazing patterns occur. This community will respond positively to improved grazing management, but significant economic inputs along with a significant amount of time are usually required to move it toward a higher successional stage and a more productive plant community. Practices such as prescribed burning and brush management are generally needed, along with extended rest and prescribed grazing once the community has degraded to this point. There are limitations to using mechanical treatment on this site due to the shallow soils. Periodic wild fire will result in a community similar to number 3, but with a smaller component of big sagebrush for a few years. There may also be a slight decrease in the amount of fringed sagewort, depending on frequency, timing, and severity of the fire. However, these will soon return if there's no change in grazing management.

State 4

Shrub, Half Shrubs, Weedy Forbs, Annuals, and Short Grasses

Community 4.1

Shrub, Half Shrubs, Weedy Forbs, Annuals, and Short Grasses

Further deterioration of community 3 results in a plant community dominated by mountain big sagebrush and undesirable plants such as fringed sagewort, broom snakeweed, plains pricklypear, weedy forbs (e.g., pussytoes and thistles), annuals such as cheatgrass and Japanese bromes and sixweeks fescue. Many increaser short grasses such as Sandberg bluegrass and plains reedgrass will be abundant. Frequently, a remnant population of climax species such as bluebunch wheatgrass will occur under sagebrush plants. Rubber rabbitbrush can be abundant in some locations. Plant community 4 produces less usable forage than the others described. The continuation of the downward trend and degradation of this site has resulted in higher soil surface temperatures, reduced water infiltration, and higher evapotranspiration. This has resulted in plant species that are more adapted to drier conditions, such as cactus. A thick canopy cover (e.g., 20% or more) of big sagebrush often results in precipitation being intercepted, thus not reaching the soil. Most of the attributes of a healthy rangeland, including good infiltration, minimal erosion and runoff, nutrient cycling and energy flow, have been lost. This community can respond positively to improved grazing management but it will take additional inputs to move it towards communities similar in production and composition to others that have been described. Once plants such as mountain big sagebrush become established, they are very difficult to remove and replace by grazing management alone. Additionally, the chances for success are significantly reduced. Practices such as prescribed burning or brush management can reduce the amount of sagebrush, as well as some of the other susceptible plant species. The potential for success depends on the composition of the rest of the plant community. Because of the shallow soils (and sometimes, steeper slopes) associated with this ecological site, other practices such as mechanical treatment or seeding are generally not feasible nor recommended.

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Shrub/Vine					
0	Shrubs and Half-shrubs			0–78	
	Shrub, broadleaf	2SB	<i>Shrub, broadleaf</i>	0–78	–
	silver sagebrush	ARCAV2	<i>Artemisia cana ssp. viscidula</i>	0–78	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0–78	–
	mountain big sagebrush	ARTRV	<i>Artemisia tridentata ssp. vaseyana</i>	0–78	–
	rubber rabbitbrush	ERNAN5	<i>Ericameria nauseosa ssp. nauseosa var. nauseosa</i>	0–78	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0–78	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–1	–
Grass/Grasslike					
0	Grasses and Sedges			1048–1334	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	493–1098	–
	rough fescue	FECA4	<i>Festuca campestris</i>	370–785	–
	Idaho fescue	FEID	<i>Festuca idahoensis</i>	62–235	–
	plains muhly	MUCU3	<i>Muhlenbergia cuspidata</i>	62–157	–
	green needlegrass	NAVI4	<i>Nassella viridula</i>	31–78	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	31–78	–
	Cusick's bluegrass	POCU3	<i>Poa cusickii</i>	0–78	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–78	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–78	–
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–78	–

	Forb, perennial	2FP	Forb, perennial	0–62	–
	Columbia needlegrass	ACNEN2	<i>Achnatherum nelsonii</i> ssp. <i>nelsonii</i>	31–78	–
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	0–78	–
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	0–78	–
	plains reedgrass	CAMO	<i>Calamagrostis montanensis</i>	0–78	–
	tufted wheatgrass	ELMA7	<i>Elymus macrourus</i>	31–78	–
	purple threeawn	ARPU9	<i>Aristida purpurea</i>	0–1	–
	Fendler's threeawn	ARPUF	<i>Aristida purpurea</i> var. <i>fendleriana</i>	0–1	–
Forb					
0	Forbs			123–157	
	common starlily	LEMO4	<i>Leucocrinum montanum</i>	0–62	–
	dotted blazing star	LIPU	<i>Liatris punctata</i>	12–62	–
	desertparsley	LOMAT	<i>Lomatium</i>	0–62	–
	lupine	LUPIN	<i>Lupinus</i>	0–62	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	0–62	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0–62	–
	American vetch	VIAM	<i>Vicia americana</i>	0–62	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–62	–
	prairie clover	DALEA	<i>Dalea</i>	12–62	–
	larkspur	DELPH	<i>Delphinium</i>	0–1	–
	deathcamas	ZIGAD	<i>Zigadenus</i>	0–1	–

Animal community

Livestock Grazing Interpretations: Managed livestock grazing is suitable on this site as it has the potential to produce a limited amount of high quality forage. Grazing must be managed carefully on this site to be sure livestock drift onto the better, more productive sites is not excessive. Management objectives should include maintenance or improvement of the plant community.

Using shorter grazing periods and providing for adequate re-growth after grazing are recommended for plant maintenance, health, and recovery. Continual over stocking and season-long use of this site can be detrimental and will alter the plant composition and production over time. The result will be plant communities that resemble numbers 3 and 4, depending on how long this grazing management is used as well as other circumstances such as weather conditions and fire frequency.

Whenever Plant Community 2 (medium and short grasses) occurs, grazing management strategies that will prevent further degradation need to be implemented. This community is still stable, productive, and healthy provided it receives proper management. It will respond fairly quickly to improved grazing management, including increased growing season rest of key forage plants. Grazing management alone can usually move this back towards the potential / historic climax community.

Plant community 3 is the result of long-term, heavy, continuous grazing and/or annual, early spring seasonal grazing. Repeated heavy early spring grazing, especially during stem elongation (generally mid May through mid June), can also have detrimental affects on the taller, key forage species. Repeated spring grazing depletes stored carbohydrates, resulting in weakening and eventual death of the cool season tall and medium grasses. This plant community can occur throughout the pasture, on spot grazed areas, and around water sources where season-long grazing patterns occur.

Plant Communities 3 and 4 have a high percentage of aggressive, less-desirable species. Once these have become established, it is significantly more difficult using grazing management alone to restore the site to one that resembles the HCPC. It becomes critical at this point to implement a grazing strategy that will restore the stability

and health of the site. Additional rest, sometimes for the growing season, or more probable for a full year or more, is often necessary for re-establishment of the desired species. There are limitations to using mechanical treatment on this site due to the shallow soils.

Calculating Safe Stocking Rates: Proper stocking rates should be incorporated into a grazing management strategy that protects the resource, maintains or improves rangeland health, and is consistent with management objectives. Safe stocking rates will be based on useable forage production, and should consider ecological condition and trend of the site, and past grazing use history.

Calculations used to determine a safe stocking rate are based on the amount of useable forage available, taking into account the harvest efficiency of the animal and the grazing strategy to be implemented. Average annual production must be measured or estimated to properly assess useable forage production and stocking rates.

Stocking rates are calculated from average forage production values using a 25% Harvest Efficiency factor for preferred and desirable plants, and 10% Harvest Efficiency for less desirable species. AUM calculations are based on 915 pounds (air-dry) per animal unit month (AUM) for a 1,000-pound cow with calf up to 4 months. No adjustments have been made for site grazability factors, such as steep slopes, site inaccessibility, or distance to drinking water.

The following is an example of how to calculate the recommended stocking rate. This example does not use production estimates from this specific ecological site. You will need to adjust the annual production values and run the calculations using total annual production values from the ecological sites encountered on each individual ranch/pasture. Before making specific recommendations, an on-site evaluation must be made.

Example of total annual production amounts by type of year:

Favorable years = 2200 lbs/acre

Normal years = 1480 lbs/acre

Unfavorable years = 1200 lbs/acre

It is recommended that on slopes of 30% or less, stocking rate should be derived from the total annual production pounds minus 500 pounds for residual dry matter and 25% harvest efficiency. On slopes over 30%, stocking rate is derived from total annual production pounds minus 800 pounds for residual dry matter and 25% harvest efficiency. Refer to the NRCS National Range and Pasture Handbook for a list of Animal Unit Equivalents.

Sample Calculations using Favorable Year production amounts:

< 30% slopes: $AUM/AC = [(2200-500)(0.25)]/915 \text{ lbs/month for one AU} = 0.46 \text{ AUM/AC}$
 $AC/AUM = (1.0 \text{ AU})/(0.46 \text{ AUM/AC}) = 2.2 \text{ AC/AUM}$

> 30% slopes: $AUM/AC = [(2200-800)(0.25)]/915 \text{ lbs/month for one AU} = 0.38 \text{ AUM/AC}$
 $AC/AUM = (1.0 \text{ AU})/(0.38 \text{ AUM/AC}) = 2.6 \text{ AC/AUM}$

NOTE: 915 lbs/month for one Animal Unit is used as the baseline for maintenance requirements. This equates to 30 lbs/day of air-dry forage (1200 lb cow at 2.5% of body weight).

Hydrological functions

The soils associated with this ecological site are generally in Hydrologic Soil Group D. The infiltration rates for these soils will normally be slow to very slow. The runoff potential for this site is moderate to high, depending on slope and ground cover/health. Runoff curve numbers generally range from 79 to 94.

Good hydrologic conditions exist on rangelands if plant cover (grass, litter, and brush canopy) is greater than 70%. Fair conditions exist when cover is between 30 and 70%, and poor conditions exist when cover is less than 30%. Sites in high similarity to HCPC (Plant Communities 1 and 2) generally have enough plant cover and litter to optimize infiltration, minimize runoff and erosion, and have a good hydrologic condition. The deep root systems of

the potential vegetation help maintain or increase infiltration rates and reduce runoff.

Sites in low similarity (Plant Communities 3 and 4) are generally considered to be in poor hydrologic condition as the majority of plant cover is from shallow-rooted species such as Sandberg bluegrass.

Erosion is minor for sites in high similarity. Rills and gullies should not be present. Water flow patterns, if present, will be barely observable. Plant pedestals are essentially non-existent. Plant litter remains in place and is not moved by erosion. Soil surfaces should not be compacted or crusted. Plant cover and litter helps retain soil moisture for use by the plants. Maintaining a healthy stand of perennial vegetation will optimize the amount of precipitation that is received. (Reference: Engineering Field Manual, Chapter 2 and Montana Supplement 4).

Recreational uses

This site provides some recreational opportunities for hiking, horseback riding, big game and upland bird hunting. The forbs have flowers that appeal to photographers. This site provides valuable open space and visual aesthetics. Caution should be used during wet weather periods.

Wood products

None.

Contributors

Robert Leinard; Barbara Gibbons; Loretta Metz; Peter Husby

Approval

Kirt Walstad, 7/19/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	J. Siddoway, R. Bandy, G. Petersen
Contact for lead author	grant.petersen@usda.gov
Date	04/19/2005
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Slopes most common on this site are between 15–35% and with at least 85% of the soil surface well-covered, rills, if evident will be rare, but may occur in bare areas after extreme convection storms – rills in this case would be narrow and less than 10 feet in length.
-

2. **Presence of water flow patterns:** Will be evident on this site with the steeper slopes, and with areas of bare ground, there may be areas which show accumulations of litter due to water movement, even after minor storm events.
-
3. **Number and height of erosional pedestals or terracettes:** Wind erosion will be rare on this site, but water erosion on the steeper slopes may have plants that could have pedestals and terracettes which could be 0.5 inch in height at the top of the slope and 1.0 inch towards the bottom of the slope.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground should be 15% or less on this site.
-
5. **Number of gullies and erosion associated with gullies:** Current gully erosion will not be evident on this site, but there may be gullies which have "healed" from the distant past.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** Appearance or evidence of these erosional features on the landscape would not be present on this site.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Litter movement will be minimal on the gradual slopes, however on the steeper slopes there will be evidence of litter movement (i.e. debris dams) which may travel up to 10 feet.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Resistance to erosion will be high with soil stability values of 5 or 6; areas of bare soil on this site may have values between 1 and 4 under plant canopy.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface structure is blocky; A horizon depth is 2 – 4 ".
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Dominance of taller, deep rooted bunchgrasses will maximize infiltration and minimize runoff throughout the site.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Will not be present generally, but there may be areas that have "healed" from former bison trails and wallows as well as more current livestock trails which could have a compaction layer below the soil surface.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Cool season, taller bunchgrasses (bluebunch wheatgrass, rough fescue)

Sub-dominant: cool season mid grasses (Idaho fescue) > cool season rhizomatous grasses (western wheatgrass) = perennial forbs > cool season short grasses (Sandberg bluegrass) = perennial forbs > shrubs = warm season shortgrass (blue grama) = sedges

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Will be low for all functional groups in a given year. Prolonged droughts which last more than 3 years may show increases in mortality and decadence for all plant groups.
-

14. **Average percent litter cover (%) and depth (in):** Moderate litter cover that tends to be thin in depth.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1100 - 1400 #/acre. This would be the expected production for the reference state during adequate moisture years. 1300 pounds would be the expected production in a 15 inch precipitation zone.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Blue grama, field brome, a variety of annual or biennial weedy forbs, fringed sagewort, broom snakeweed, curlycup gumweed, prickly pear cactus, cheatgrass.
-

17. **Perennial plant reproductive capability:** During adequate moisture years bunchgrasses will generally produce seeds, however the cool season rhizomatous grasses may not necessarily produce seed even with adequate moisture.
-