

Ecological site R048AY238CO Brushy Loam

Last updated: 3/05/2024
Accessed: 05/17/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	J. Murray, C. Holcomb, L. Santana, F. Cummings, A. Jones, P. Billig, S. Jaouen
Contact for lead author	
Date	12/08/2004
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None

2. **Presence of water flow patterns:** None

3. **Number and height of erosional pedestals or terracettes:** None

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Expect <10% bareground. Extended drought can cause bareground to increase.

5. **Number of gullies and erosion associated with gullies:** None

6. **Extent of wind scoured, blowouts and/or depositional areas:** None

7. **Amount of litter movement (describe size and distance expected to travel):** Movement expected to be short and

minimal.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability class rating anticipated to be 5-6 in the interspaces at soil surface.
-

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soils are typically deep with a very dark color. Soil surface texture is fine clay loam, well drained.
-

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Diverse grass, forb, shrub canopy and root structure reduces raindrop impact and slows overland flow providing increased time for infiltration to occur.
-

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Shrubs >

Sub-dominant: cool season bunchgrass > forbs > sedges >

Other: cool season rhizomatous grass

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Typically minimal, expect for weather related (late spring freezes on oak brush)
-

14. **Average percent litter cover (%) and depth (in):** 70-80% litter cover at 1-2 inch depth
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 2000 lbs./ac. low precip years; 2500 lbs./ac. average precip years; 4000 lbs./ac. above average precip years. After extended drought or the first growing season following wildfire, production may be significantly reduced by 600 - 800 lbs./ac. or more.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site: Kentucky bluegrass and noxious weeds

17. **Perennial plant reproductive capability:** The only limitations are weather-related, wildfire, natural disease, inter-species competition, wildlife, and insects that may temporarily reduce reproductive capability.
-