

## **Ecological site R048AY248CO Mountain Clay Loam**

Last updated: 3/05/2024  
Accessed: 05/03/2024

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 048A–Southern Rocky Mountains

MLRA 48A makes up about 45,920 square miles (119,000 square kilometers) and is the southern part of the Rocky Mountains. The Southern Rocky Mountains lies east of the Colorado Plateau, south of the Wyoming Basin, west of the Great Plains, and north of the Rio Grande Rift. It is in western and central Colorado, southeastern Wyoming, eastern Utah, and northern New Mexico. The headwaters of major rivers such as the Colorado, Yampa, Arkansas, Rio Grande, North Platte and South Plate rivers are located here. This MLRA has numerous national forests, including the Medicine Bow National Forest in Wyoming; the Routt, Arapaho, Roosevelt, Pike, San Isabel, White River, Gunnison, Grand Mesa, Uncompahgre, Rio Grande, and San Juan National Forests in Colorado; the Carson National Forest and part of the Santa Fe National Forest in New Mexico. Rocky Mountain National Park also is in this MLRA.

MLRA 48A is the southern Rocky Mountains physiographic region. The Southern Rocky Mountains consist primarily of two belts of strongly sloping to precipitous mountain ranges trending north to south. Several basins, or parks, are between the belts. Some high mesas and plateaus are included. It is characterized by mountain ranges that were uplifted during the Laramide Orogeny and then had periods of glaciation. The ranges include the Sangre de Cristo Mountains, the Laramie Mountains, and the Front Range in the east and the San Juan Mountains and the Sawatch and Park Ranges in the west. The ranges are dissected by many narrow stream valleys having steep gradients. In some areas the upper mountain slopes and broad crests are covered by snowfields and glaciers. Elevation typically ranges from 6,500 to 14,400 feet (1,980 to 4,390 meters) in this area. The part of this MLRA in central Colorado includes the highest point in the Rockies, Mount Elbert, which reaches an elevation of 14,433 feet (4,400 meters). More than 50 peaks in the part of the MLRA in Colorado are at an elevation of more than 14,000 feet (4,270 meters). Many small glacial lakes are in the high mountains.

The mountains in this area were formed mainly by crustal uplifts during the late Cretaceous and early Tertiary periods. This large MLRA can be subdivided into at least 4 large general divisions. First is the Rockies on the east side of this area are called the "Front Range," which is a fault block that has been tilted up on edge and uplifted and is largely igneous and metamorphic geology. It was tilted up on the east edge, so there is a steep front on the east and the west side is more gently sloping and in the south east there are rocks exposed in the mountains are mostly Precambrian igneous and metamorphic rocks. Second is the tertiary rocks, primarily basalt and andesitic lava flows, tuffs, breccias, and conglomerates, are throughout this area (San Juan Mountains Area). The third division is Northwest part of the MLRA is dominantly sedimentary rock from the cretaceous/tertiary and Permian/Pennsylvanian periods. The fourth subset is the long and narrow Sangre de Cristos mountains uplifted in the Cenozoic are between the Rio Grande rift and the great plains. Many of the highest mountain ranges were reshaped by glaciation during the Pleistocene. Alluvial fans at the base of the mountains are recharge zones for local basin and valley fill aquifers. They also are important sources of sand and gravel.

The average annual precipitation ranges predominantly from 12 to 63 inches. Summer rainfall commonly occurs as high-intensity, convective thunderstorms. About half of the annual precipitation occurs as snow in winter; this proportion increases with elevation. In the mountains, deep snowpacks accumulate throughout the winter and

generally persist into spring or early summer, depending on elevation. Some permanent snowfields and small glaciers are on the highest mountain peaks. In the valleys at the lower elevations, snowfall is lighter and snowpacks can be intermittent. The average annual temperature is 26 to 54 degrees F (-3 to 12 degrees C). The freeze-free period averages 135 days and ranges from 45 to 230 days, decreasing in length with elevation. The climate of this area is strongly dependent upon elevation; precipitation is greater, and temperatures are cooler at the higher elevations. The plant communities vary with elevation, aspect and change in latitudes due to changing in precipitation kind and timing and temperature.

The dominant soil orders in this MLRA are Mollisols, Alfisols, Inceptisols, and Entisols. The soils in the area dominantly have a frigid or cryic soil temperature regime and an ustic or udic soil moisture regime. Mineralogy is typically mixed, smectitic, or paramicaceous. In areas with granite, gneiss, and schist bedrock, Glossocryalfs (Seitz, Granile, and Leadville series) and Haplocryolls (Rogert series) formed in colluvium on mountain slopes. Dystrocryepts (Leighcan and Mummy series) formed on mountain slopes and summits at the higher elevations. In areas of andesite and rhyolite bedrock, Dystrocryepts (Endlich and Whitecross series) formed in colluvium on mountain slopes. In areas of sedimentary bedrock, Haplustolls (Towave series) formed on mountain slopes at low elevations and with low precipitation. Haplocryolls (Lamphier and Razorba series), Argicryolls (Cochetopa series), and Haplocryalfs (Needleton series) formed in colluvium on mountain slopes at high elevations.

## **Classification relationships**

NRCS:

Major Land Resource Area 48A, Southern Rocky Mountains (United States Department of Agriculture, Natural Resources Conservation Service, 2006).

USFS:

M331F- Southern Parks and Rocky Mountain Range Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M331G – South Central Highlands Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M331H – North Central Highlands and Rocky Mountains Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M331I – North Parks and Ranges Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M341B – Tavaputs Plateau Section M341 Nevada-Utah Mountains Semi-Desert - Coniferous Forest - Alpine Meadow (Cleland, et al., 2007).

EPA:

21a – Alpine Zone, 21b – Crystalline Subalpine Forests, 21c – Crystalline Mid-Elevations Forests, 21d -Foothill Shrublands, 21e – Sedimentary Subalpine Forests, 21f – Sedimentary Mid-Elevation Forests, 21g – Volcanic Subalpine Forests, and 21h – Volcanic Mid-Elevation Forests < 21 Southern Rockies < 6.2 Western Cordillera < 6 Northwestern Forested Mountains North American Deserts (Griffith, 2006).

20c – Semiarid Benchlands and Canyonlands and 20e - Escarpements < 20 Colorado Plateau < 10.1 Cold Deserts < 10 North American Deserts (Griffith, 2006).

USGS: Southern Rocky Mountain Province and the southern part of Unita Basin Section Colorado Plateaus Province

## **Ecological site concept**

R048AY248CO Mountain Clay Loam occurs on alluvial fans, mesas, hills, dip slopes, and mountain slopes. Slopes is between 1 to 40 percent. Soils are deep to very deep (40 to 60+ inches). Soils are derived from alluvium and slope alluvium from shale; or alluvium, slope alluvium, colluvium and/or residuum from sandstone and shale. Soil

surface texture is loam, silty clay loam, gravelly loam or a clay loam with fine-textured subsurface. It is Arizona Fescue – western wheatgrass – Gambel's Oak community. It has a typical ustic moisture regime and frigid temperature regime. The effective precipitation ranges from 16 to 20 inches.

### Associated sites

R048AY228CO	<b>Mountain Loam</b> R048AY228CO Mountain Loam occurs mainly alluvial fans, mountain slopes, benches, terraces, or hills. Slopes average between 5 and 10% but can range from 0 to 30%. Soils are moderately deep to deep (20-60 inches) loamy soils derived from residuum from igneous and metamorphic rocks or sandstone and shale; slope alluvium from sandstone and shale, or igneous and metamorphic rocks; colluvium from igneous and metamorphic rocks or sandstone and shale, and/or alluvium from igneous and metamorphic rocks. Soil surface texture are loam, sandy loam or silt loam with loamy subsurface. It is a Mountain Big Sagebrush -Arizona Fescue community. It has a typical ustic moisture regime. The effective precipitation ranges from 16 to 20 inches.
R048AY238CO	<b>Brushy Loam</b> R048AY238CO Brushy Loam occurs on hills, mountains, complex landslides, and benches. Slopes is between 3 to 35%. Soils are moderately deep to deep (20 to 60+ inches), soils derived from colluvium, residuum, slope alluvium and alluvium from sandstone and shale. Soil surface texture is loam or clay loam with fine-textured subsurface. It is a Gambel's oak – slender wheatgrass community. It has a typical ustic moisture regime. The effective precipitation ranges from 16 to 20 inches.
R048AY247CO	<b>Deep Clay Loam</b> R048AY247CO Deep Clay Loam occurs on hills, hillsides, mountain-slope, complex landslides, alluvial fans, and structural benches. Slopes is between 0 to 35%. Soils are deep (60+ inches). Soils are derived from colluvium and slide deposits from igneous, metamorphic and sedimentary rocks, and/or alluvium, residuum or complex landslide deposits from shale. Soil surface texture is loam, clay loam or silty clay loam with fine-textured subsurface. It is a mountain big sagebrush – western wheatgrass community. It has a typical ustic moisture regime. The effective precipitation ranges from 16 to 20 inches.
R048AY255CO	<b>Pine Grasslands</b> R048AY255CO Pine Grassland occurs on structural benches, dip slopes, hills, mesas and canyon benches. Slopes is between 0 to 30%. This site has more than one soil concept correlated to it. The concepts are shallow soils (<20

### Similar sites

F048AY925CO	<b>Ponderosa Pine Forest</b> F048AY925CO occurs on hillsides, mountain-slopes, mesas, structural benches and cuestas. Slopes are 3 to 30%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived from slope alluvium from sandstone and/or shale, colluvium from sandstone and/or shale, or residuum from sandstone and shale. Soil surface texture is a loam, clay loam, sandy loam, fine sandy loam, very stony loam, cobbly sandy loam, or very boulder sandy loam with fine textured subsurface. It is a Ponderosa Pine - Muttongrass – squirreltail community. It has a typical ustic moisture regime. The effective precipitation ranges from 16 to 20 inches.
R048AY240CO	<b>Shallow Pine</b> R048AY240CO Shallow Pine occurs on mountains and mountainsides. Slopes are 5 to 40%. Soils are shallow (10 to 20 inches). Soils are derived from slope alluvium from volcanic breccia, gneiss, granite, or sandstone and/or residuum from granite, granodiorite and/or gneiss. Soil surface texture is a gravelly to very gravelly sandy loam or very gravelly loam with loamy-skeletal subsurface. It is a Ponderosa Pine - Arizona Fescue – Mountain Muhly community. It has a typical ustic moisture regime. The effective precipitation ranges from 16 to 20 inches.
R048AY257CO	<b>Clayey Valley</b> R048AY257CO Clayey Valley occurs on hillslopes, and old high terraces. Slopes is between 1 to 12%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived from slope alluvium from shale or alluvium from sandstone and shale. Soil surface texture is loam or clay loam with fine textured subsurface. It is Western wheatgrass – Arizona Fescue with scattered Ponderosa Pine. It has a Typical ustic moisture regime and frigid temperature regime. The effective precipitation ranges from 16 to 20 inches.

R048AY255CO	<b>Pine Grasslands</b> R048AY255CO Pine Grassland occurs on structural benches, dip slopes, hills, mesas and canyon benches. Slopes is between 0 to 30%. This site has more than one soil concept correlated to it. The concepts are shallow soils (<20
R048AY247CO	<b>Deep Clay Loam</b> R048AY247CO Deep Clay Loam occurs on hills, hillsides, mountain-slope, complex landslides, alluvial fans, and structural benches. Slopes is between 0 to 35%. Soils are deep (60+ inches). Soils are derived from colluvium and slide deposits from igneous, metamorphic and sedimentary rocks, and/or alluvium, residuum or complex landslide deposits from shale. Soil surface texture is loam, clay loam or silty clay loam with fine-textured subsurface. It is a mountain big sagebrush – western wheatgrass community. It has a typic ustic moisture regime. The effective precipitation ranges from 16 to 20 inches.

**Table 1. Dominant plant species**

Tree	(1) <i>Pinus ponderosa</i>
Shrub	(1) <i>Quercus gambelii</i>
Herbaceous	(1) <i>Festuca arizonica</i> (2) <i>Muhlenbergia montana</i>

## Physiographic features

Although it is in the ponderosa pine zone, it is sparsely timbered, if at all. Slopes are about 1 to 40 percent. North and east-facing slopes are generally the most productive and tend to grow more tall shrubs and trees, but in many cases, are not significantly higher in potential for range forage. In some cases this range site is on the drier slopes, northerly facing slopes being forested. Elevations are about 7000 to 9500 feet.

**Table 2. Representative physiographic features**

Landforms	(1) Hill (2) Mountain slope (3) Mesa (4) Dip slope (5) Alluvial fan (6) Mountain (7) Bench
Runoff class	High to very high
Flooding frequency	None
Ponding frequency	None
Elevation	7,000–9,500 ft
Slope	1–40%
Aspect	Aspect is not a significant factor

## Climatic features

Average annual precipitation is about 16 to 20 inches. Of this, approximately 45-55% falls as snow, and 45-55% falls as rain between middle of May to and the end of September. Summer moisture is mostly from thundershowers in July, August and September. December to February is the driest period of the year with the driest month being January. July thru September is the wettest period and the wettest month is usually August. The average annual total snowfall is 84.9 inches. The snow depth usually ranges from 1 to 5 inches during November thru March. The highest winter snowfall record in this area is 127 inches which occurred in 2007-2008. The lowest snowfall record is 46.5 inches during the 2017-2018 winter. The frost-free period typically ranges from 80 to 120 days. The last spring frost is typically the middle of June to the end of June. The first fall frost is usually the end of August to the middle of September. Mean daily annual air temperature ranges from about 25.5°F to 60.3°F, averaging about 24°F for the winter and 61.8°F in the summer. Summer high temperatures of mid-70°F to low 80°F are not unusual. The coldest winter temperature recorded was -36°F on February 2, 1985 and the warmest winter temperature recorded was 65°F on December 5, 1995. The coldest summer temperature recorded was 19°F on June 2, 1990 and the warmest

was 98°F on July 31, 2002. Wide yearly and seasonal fluctuations are common for this climatic zone. Data taken from Western Regional Climate Center (2018) for Ridgway, Colorado Climate Station.

This zone in MLRA 48 will need to be broken up into at multiple land resources zones in future projects based on current knowledge of precipitation and temperature patterns.

West Central Zone Stations: Alterbern, Aspen, Avon, Glenwood Springs #2, Shoshone, Placerville and Ridgway. This LRU zone is use in write up above. Driest month is usually January, February and June and wettest months are July, August and September.

Northwest Zone Climate Stations: Meeker and Yampa are at the low end of this LRU zone. Driest months usually are January and February. Wettest months usually are April and August.

Southwest Zone Climate Stations (Precambrian sedimentary and igneous): There are no climate stations in this LRU zone.

Southwest Volcanics: There are no climate stations in this LRU zone.

Northeast (Front Range Igneous and Metamorphic): Cabin Creek, Caribou Ranch, Dillion 1 R, Fraser, Georgetown, Grand lake 1 NW, Hourglass Reservoir, Nederland 2 NNE, Red Feathers Lakes, Red Feather Lakes 2 SE and Victor. April, May, July and August are the wettest months. February, December, November and October are the driest. The climate stations is this zone are cryic. These areas have shorter growing seasons by 20 to 40 days over the frigid stations.

Southeast (Sangre de Cristo Mtns): There are no climate stations in this zone in MLRA 48A. Closest ones are in MLRA 49. The growing season appears to be longer on the Sangre de Cristos. Driest months are December to February and the wettest are July & August.

Crylic High elevation valleys: Pitkin, Taylor River and Meredith. These areas have shorter growing seasons by 20 to 40 days over the frigid stations.

**Table 3. Representative climatic features**

Frost-free period (characteristic range)	23-78 days
Freeze-free period (characteristic range)	75-111 days
Precipitation total (characteristic range)	17-18 in
Frost-free period (actual range)	5-101 days
Freeze-free period (actual range)	43-134 days
Precipitation total (actual range)	17-19 in
Frost-free period (average)	54 days
Freeze-free period (average)	92 days
Precipitation total (average)	18 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

### Climate stations used

- (1) ASPEN PITKIN CO AP [USW00093073], Aspen, CO
- (2) GEORGETOWN [USC00053261], Idaho Springs, CO
- (3) GLENWOOD SPGS #2 [USC00053359], Glenwood Springs, CO
- (4) HOURGLASS RSVR [USC00054135], Bellvue, CO
- (5) RIDGWAY [USC00057020], Ridgway, CO
- (6) YAMPA [USC00059265], Toponas, CO
- (7) FRASER [USC00053116], Fraser, CO

### Influencing water features

None

### Soil features

Shale parent material has had a strong influence on these soils. Surface soils are high in clay or, if loam-textured, are thin over clayey material. They are dark to gray colored, friable, and usually non-calcareous. the subsoil is typically a compact, slowly permeable clay which retards root penetration of many plants. Depth to shale usually 20 inches or greater Deeper slopes are found commonly on footslopes. Outcrops of raw shale are commonly associated with the site. On the steeper slopes, these soils are subject to geologic slides when completely saturated. Water-holding capacity is generally moderate, but the high wilting point is a factor in dry weather. As much as 2.5 inches of water may be required to wet one foot of dry soil. Light showers therefore have little effect and runoff from heavier storms is often high. The soils are highly erodible.

Soils correlated to this site are:

Fine: Bradfield, Campspass, Gladlow, Herm, Holderness, Kinesava, Sagedale, and Zoltay

Table 4. Representative soil features

Parent material	(1) Alluvium–shale (2) Slope alluvium–shale (3) Alluvium–sandstone and shale (4) Slope alluvium–sandstone and shale (5) Colluvium–sandstone and shale (6) Residuum–sandstone and shale (7) Alluvium–shale and siltstone (8) Colluvium–shale and siltstone
Surface texture	(1) Clay loam (2) Loam (3) Gravelly loam (4) Silty clay loam
Family particle size	(1) Fine
Drainage class	Well drained
Permeability class	Slow to moderate
Soil depth	40–100 in
Surface fragment cover ≤3"	0–15%
Surface fragment cover >3"	0–5%
Available water capacity (Depth not specified)	5–7.5 in
Calcium carbonate equivalent (Depth not specified)	0–5%
Soil reaction (1:1 water) (Depth not specified)	6.6–7.8
Subsurface fragment volume ≤3" (Depth not specified)	0–15%
Subsurface fragment volume >3" (Depth not specified)	0–20%

## Ecological dynamics

Gambel oak with an understory of grasses and scattered forbs and low shrubs is a typical feature of this site. A few trees may occur, but are not always present. Major grasses are Arizona fescue, western wheatgrass, and mountain muhly.

Others commonly present are Letterman and Columbia needlegrass, needle-and-thread, nodding brome, slender wheatgrass, muttongrass, junegrass, squirreltail, spike muhly, and elk sedge. Thurber fescue shows up at upper elevations. Herbaceous sage and members of the milkvetch-poison vetch group are among the most prevalent forbs. Others are buckwheats, mountain goldenweed, lupine, ligusticum, trailing fleabane, penstemon, yarrow, sweetvetch, and mules-ear wyethia. Minor amounts of black sagebrush are common, and there may be some big or finged sagebrush. Associated with gambel oak is a scattering of shrubs such as snowberry, serviceberry, mountain mahogany, and bitterbrush. Dwarf rabbitbrush is usually noticeable. Differences in micro-climate cause a great deal of variation in the vegetation.

There may be scattered trees or patches of ponderosa pine, but these are not consistently present. An occasional pinyon pine or rocky mountain juniper may occur on drier portions of the site.


The state and transition model was added to fill the provisional ecological site instruction. It is a very general model.

## State and transition model




## R048AY248CO Mountain Clay Loam

### State 1: Reference State



T1A

### 2: Seeded State



## Legend

1.1A, 2.1A – lack of fire/disturbance, time without disturbance, lack of insect/pathogen outbreaks, and/or possible tree encroachment (pinyon, or Doug fir)

1.1B, 1.3B, 2.1B, 2.3B – fire/natural disturbance, insect/pathogens outbreaks, and/or tree encroachment removal

1.2A, 2.2A – lack of fire/disturbance, time without disturbance, lack of insect/pathogen outbreaks

1.3A, 2.3A – natural disturbance, insect/pathogens outbreaks and/or tree encroachment removal on a smaller scale

T1A – Seeding, and/or vegetative treatments (fire, mechanical, chemical) of shrubs and/trees

**State 1**  
**Reference State**

**Community 1.1**

Reference State

Approximate ground cover is 35%. Species most likely to invade or increase from trace amounts are kentucky bluegrass, sleepygrass, snakeweed, colorado rubberweed and pussytoes. As the ecological condition deteriorates, grasses weaken and thin out. Arizona fescue and mountain mully may eventually disappear. Western wheatgrass, needlegrasses, and elk sedge han on longer than other important grasses. Either gambel oak or black sagebrush often assumes strong dominance. Forbs likely to become much more prominent than originally are poisonvetches, lupine, hairy goldaster, and in more moist locations, mulesear wyethia. These changes are likely to be accompanied by serious erosion. Approximate ground cover is 35%.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	410	550	825
Shrub/Vine	190	250	375
Tree	75	100	150
Forb	75	100	150
Total	750	1000	1500

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1				400–700	
	Arizona fescue	FEAR2	<i>Festuca arizonica</i>	150–250	–
	Thurber's fescue	FETH	<i>Festuca thurberi</i>	100–150	–
	Columbia needlegrass	ACNE9	<i>Achnatherum nelsonii</i>	100–150	–
	mountain muhly	MUMO	<i>Muhlenbergia montana</i>	100–150	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	100–150	–
	muttongrass	POFE	<i>Poa fendleriana</i>	10–50	–
	Letterman's needlegrass	ACLE9	<i>Achnatherum lettermanii</i>	10–50	–
	nodding brome	BRAN	<i>Bromus anomalus</i>	10–50	–
	Geyer's sedge	CAGE2	<i>Carex geyeri</i>	10–50	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	10–50	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	10–50	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	10–50	–
<b>Forb</b>					
2				50–150	
	milkvetch	ASTRA	<i>Astragalus</i>	1–3	–
	trailing fleabane	ERFL	<i>Erigeron flagellaris</i>	1–2	–
	buckwheat	ERIOG	<i>Eriogonum</i>	1–2	–
	Porter's licorice-root	LIPO	<i>Ligusticum porteri</i>	1–2	–
	silvery lupine	LUAR3	<i>Lupinus argenteus</i>	1–2	–
	locoweed	OXYTR	<i>Oxytropis</i>	1–2	–
	beardtongue	PENST	<i>Penstemon</i>	1–2	–
	American vetch	VIAM	<i>Vicia americana</i>	1–2	–
	mule-ears	WYETH	<i>Wyethia</i>	1–2	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	1–2	–
<b>Shrub/Vine</b>					
3				100–350	
	Gambel oak	QUGA	<i>Quercus gambelii</i>	100–150	–
	common snowberry	SYAL	<i>Symphoricarpos albus</i>	10–50	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	10–50	–
	Saskatoon serviceberry	AMAL2	<i>Amelanchier alnifolia</i>	10–50	–
	black sagebrush	ARNO4	<i>Artemisia nova</i>	10–50	–
	alderleaf mountain mahogany	CEMO2	<i>Cercocarpus montanus</i>	10–20	–
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	10–20	–
<b>Tree</b>					
4				50–150	
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	50–150	–

## Animal community

### WILDLIFE INTERPRETATIONS:

The site has a high value rating for deer, cottontail, and wild turkey. There is a medium value for elk, upland game

birds. There is a low rating for jackrabbit.

#### LIVESTOCK INTERPRETATIONS:

This site has a high value for goats, a medium value for cattle, sheep, and horses.

#### Recreational uses

This site has a medium value for recreation and natural beauty.

#### Wood products

This site is low for wood products.

#### Inventory data references

Field offices in Colorado where the site occurs are Cortez, Durango, Norwood, and Pagosa Springs.

#### Type locality

Location 1: Dolores County, CO	
General legal description	Location of typical example of the site: Hills near Thompson Lane and on upper drainages of Beaver and Plateau Creeks, north of Dolores in Dolores County.

#### Other references

##### References

Chapman, S.S., G.E. Griffith, J.M. Omernik, A.B. Price, J. Freeouf, and D.L. Schrupp. 2006. Ecoregions of Colorado. (2-sided color poster with map, descriptive text, summary tables, and photographs). U.S. Geological Survey, Reston, VA. Scale 1:1,200,000.

Cleland, D.T.; Freeouf, J.A.; Keys, J.E.; Nowacki, G.J.; Carpenter, C.A.; and McNab, W.H. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Gen. Tech. Report WO-76D [Map on CD-ROM] (A.M. Sloan, cartographer). Washington, DC: U.S. Department of Agriculture, Forest Service, presentation scale 1:3,500,000; colored.

Soil Conservation Service (SCS). August 1975. Range Site Description for Mountain Clay Loam #248. : USDA, Denver Colorado.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Western Regional Climate Center. Retrieved from <http://www.wrcc.dri.edu/summary/Climsmco.html> on December 10, 2018

#### Contributors

Suzanne Mayne-Kinney

#### Approval

Kirt Walstad, 3/05/2024

#### Acknowledgments

Project Staff:

Suzanne Mayne-Kinney, Ecological Site Specialist, NRCS MLRA, Grand Junction SSO  
Chuck Peacock, MLRA Soil Survey Leader, NRCS MLRA Grand Junction SSO

**Program Support:**

Rachel Murph, NRCS CO State Rangeland Management Specialist, Denver  
Scott Woodhall, NRCS MLRA Ecological Site Specialist-QA Phoenix, AZ  
Eva Muller, Regional Director, Rocky Mountain Regional Soil Survey Office, Bozeman, MT  
B.J. Shoup, CO State Soil Scientist, Denver  
Eugene Backhaus, CO State Resource Conservationist, Denver

Those involved in developing earlier versions of this site description include: Bob Rayer, retired NRCS Soil Scientist; Herman Garcia, retired CO State RMS and NRCS MLRA Ecological Site Specialist-QA Phoenix, AZ.

**--Site Development and Testing Plan--:**

Future work to validate and further refine the information in this Provisional Ecological Site Description is necessary. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data.

Additional information and data is required to refine the Plant Production and Annual Production tables for this ecological site. The extent of MLRA 48A must be further investigated.

Field testing of the information contained in this Provisional ESD is required. As this ESD is moved to the Approved ESD level, reviews from the technical team, quality control, quality assurance, and peers will be conducted.

## **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/03/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## **Indicators**

**1. Number and extent of rills:**

---

**2. Presence of water flow patterns:**

---

**3. Number and height of erosional pedestals or terracettes:**

---

**4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not**

bare ground):

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-