

Ecological site R048AY308CO Shallow Alpine

Last updated: 3/05/2024
Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 048A–Southern Rocky Mountains

MLRA 48A makes up about 45,920 square miles (119,000 square kilometers) and is the southern part of the Rocky Mountains. The Southern Rocky Mountains lies east of the Colorado Plateau, south of the Wyoming Basin, west of the Great Plains, and north of the Rio Grande Rift. It is in western and central Colorado, southeastern Wyoming, eastern Utah, and northern New Mexico. The headwaters of major rivers such as the Colorado, Yampa, Arkansas, Rio Grande, North Platte and South Plate rivers are located here. This MLRA has numerous national forests, including the Medicine Bow National Forest in Wyoming; the Routt, Arapaho, Roosevelt, Pike, San Isabel, White River, Gunnison, Grand Mesa, Uncompahgre, Rio Grande, and San Juan National Forests in Colorado; the Carson National Forest and part of the Santa Fe National Forest in New Mexico. Rocky Mountain National Park also is in this MLRA.

MLRA 48A is the southern Rocky Mountains physiographic region. The Southern Rocky Mountains consist primarily of two belts of strongly sloping to precipitous mountain ranges trending north to south. Several basins, or parks, are between the belts. Some high mesas and plateaus are included. It is characterized by mountain ranges that were uplifted during the Laramide Orogeny and then had periods of glaciation. The ranges include the Sangre de Cristo Mountains, the Laramie Mountains, and the Front Range in the east and the San Juan Mountains and the Sawatch and Park Ranges in the west. The ranges are dissected by many narrow stream valleys having steep gradients. In some areas the upper mountain slopes and broad crests are covered by snowfields and glaciers. Elevation typically ranges from 6,500 to 14,400 feet (1,980 to 4,390 meters) in this area. The part of this MLRA in central Colorado includes the highest point in the Rockies, Mount Elbert, which reaches an elevation of 14,433 feet (4,400 meters). More than 50 peaks in the part of the MLRA in Colorado are at an elevation of more than 14,000 feet (4,270 meters). Many small glacial lakes are in the high mountains.

The mountains in this area were formed mainly by crustal uplifts during the late Cretaceous and early Tertiary periods. This large MLRA can be subdivided into at least 4 large general divisions. First is the Rockies on the east side of this area are called the "Front Range," which is a fault block that has been tilted up on edge and uplifted and is largely igneous and metamorphic geology. It was tilted up on the east edge, so there is a steep front on the east and the west side is more gently sloping and in the south east there are rocks exposed in the mountains are mostly Precambrian igneous and metamorphic rocks. Second is the tertiary rocks, primarily basalt and andesitic lava flows, tuffs, breccias, and conglomerates, are throughout this area (San Juan Mountains Area). The third division is Northwest part of the MLRA is dominantly sedimentary rock from the cretaceous/tertiary and Permian/Pennsylvanian periods. The fourth subset is the long and narrow Sangre de Cristos mountains uplifted in the Cenozoic are between the Rio Grande rift and the great plains. Many of the highest mountain ranges were reshaped by glaciation during the Pleistocene. Alluvial fans at the base of the mountains are recharge zones for local basin and valley fill aquifers. They also are important sources of sand and gravel.

The average annual precipitation ranges predominantly from 12 to 63 inches. Summer rainfall commonly occurs as high-intensity, convective thunderstorms. About half of the annual precipitation occurs as snow in winter; this proportion increases with elevation. In the mountains, deep snowpacks accumulate throughout the winter and

generally persist into spring or early summer, depending on elevation. Some permanent snowfields and small glaciers are on the highest mountain peaks. In the valleys at the lower elevations, snowfall is lighter and snowpacks can be intermittent. The average annual temperature is 26 to 54 degrees F (-3 to 12 degrees C). The freeze-free period averages 135 days and ranges from 45 to 230 days, decreasing in length with elevation. The climate of this area is strongly dependent upon elevation; precipitation is greater, and temperatures are cooler at the higher elevations. The plant communities vary with elevation, aspect and change in latitudes due to changing in precipitation kind and timing and temperature.

The dominant soil orders in this MLRA are Mollisols, Alfisols, Inceptisols, and Entisols. The soils in the area dominantly have a frigid or cryic soil temperature regime and an ustic or udic soil moisture regime. Mineralogy is typically mixed, smectitic, or paramicaceous. In areas with granite, gneiss, and schist bedrock, Glossocryalfs (Seitz, Granile, and Leadville series) and Haplocryolls (Rogert series) formed in colluvium on mountain slopes. Dystrocryepts (Leighcan and Mummy series) formed on mountain slopes and summits at the higher elevations. In areas of andesite and rhyolite bedrock, Dystrocryepts (Endlich and Whitecross series) formed in colluvium on mountain slopes. In areas of sedimentary bedrock, Haplustolls (Towave series) formed on mountain slopes at low elevations and with low precipitation. Haplocryolls (Lamphier and Razorba series), Argicryolls (Cochetopa series), and Haplocryalfs (Needleton series) formed in colluvium on mountain slopes at high elevations.

Classification relationships

NRCS:

Major Land Resource Area 48A, Southern Rocky Mountains (United States Department of Agriculture, Natural Resources Conservation Service, 2006).

USFS:

M331F- Southern Parks and Rocky Mountain Range Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M331G – South Central Highlands Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M331H – North Central Highlands and Rocky Mountains Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M331I – North Parks and Ranges Section Southern Rocky Mountain Steppe - Open Woodland - Coniferous Forest - Alpine Meadow

M341B – Tavaputs Plateau Section M341 Nevada-Utah Mountains Semi-Desert - Coniferous Forest - Alpine Meadow (Cleland, et al., 2007).

EPA:

21a – Alpine Zone, 21b – Crystalline Subalpine Forests, 21c – Crystalline Mid-Elevations Forests, 21d -Foothill Shrublands, 21e – Sedimentary Subalpine Forests, 21f – Sedimentary Mid-Elevation Forests, 21g – Volcanic Subalpine Forests, and 21h – Volcanic Mid-Elevation Forests < 21 Southern Rockies < 6.2 Western Cordillera < 6 Northwestern Forested Mountains North American Deserts (Griffith, 2006).

20c – Semiarid Benchlands and Canyonlands and 20e - Escarpements < 20 Colorado Plateau < 10.1 Cold Deserts < 10 North American Deserts (Griffith, 2006).

USGS: Southern Rocky Mountain Province and the southern part of Unita Basin Section Colorado Plateaus Province

Ecological site concept

R048AY308CO Shallow Alpine occurs mountain-slopes, ridges, mountains, basin floor, cirque floors, and valleys floors. Slopes is between 5 to 60 percent. Soils are shallow (7 to 20 inches). Soils are derived mostly from slope alluvium from tuff, rhyolite, andesite, basalt volcanic breccia, and other volcanic rocks; residuum from sandstone,

tuff, rhyolite, volcanic breccia, basalt, andesite, and limestone; till from andesite, conglomerate or sedimentary rock; or colluvium from rhyolite, tuff, andesite, basalt, and other volcanic rocks. Soil surface texture is very gravelly, very stony, or very cobby loam; loam; very stony or very cobby sandy loam, or very cobby silt loam with loamy—skeletal or loamy subsurface. It is Kobresia – Tufted Hairgrass community. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.

Associated sites

R048AY250CO	<p>Subalpine Loam R048AY250CO Subalpine Loam occurs on hills, mountain-slopes, and mountains. Slopes is between 1 to 30%. Soils are deep to very deep (20 to 60+ inches). Soils are derived from colluvium and alluvium from volcanic rock; complex landslide deposits from igneous, metamorphic, and sedimentary rock; and slope alluvium, colluvium, residuum, alluvium or complex landslide deposits from sandstone and shale or shale. Soil surface texture is loam with loamy textured subsurface. It is a mountain big sagebrush – Thurber's Fescue community. It has an ustic udic/typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 20 to 30 inches.</p>
R048AY253CO	<p>Wet Subalpine R048AY253CO Wet Subalpine occurs on mesa tops and swales, drainageways, glacial moraines, and alluvial fans formed by glacial action. Slopes is between 0 to 30%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived from till from basalt, andesite, or tuff; slope alluvium from andesite or tuff, alluvium from andesite or tuff; eolian deposits from basalt; or residuum from basalt. Soil surface texture is loam, or gravelly loam with fine-loamy, loamy-skeletal or clayey-skeletal textured subsurface. It is slender wheatgrass – tufted hairgrass. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 37 to 45 inches.</p>
R048AY304CO	<p>Alpine Slopes R048AY304CO Alpine Slopes occurs on mountain-slopes, ridges, mountains and valleys. Slopes is between 30 to 60%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived mostly from colluvium and alluvium from volcanic, metamorphic and igneous rocks. Soil surface texture is gravelly, very gravelly, extremely gravelly, very cobby, extremely cobby, or very stony loam or extremely stony sandy loam with loamy--skeletal subsurface. It is Kobresia – Tufted Hairgrass community. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.</p>
R048AY305CO	<p>Alpine Meadow R048AY305CO Alpine Meadow occurs on depressions, drainageways, cirques, flood plains and steams. Slopes is between 0 to 15%. Soils are greater than 60 inches in depth. Soils are derived mostly from colluvium from Monzonite, volcanic rock; alluvium from volcanic rocks; or slope alluvium from Andesite or latite. Soil surface texture is loam, cobby loam or silt loam. The underlying material to a depth of 60 inches or more is variable material ranging from sandy loam to clay loam with variable amounts and sizes of rock fragments. It is Kobresia – Tufted Hairgrass community. This site has a water table at 0 to 36 inches in depth. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.</p>

Similar sites

R048AY309CO	<p>Warm Alpine R048AY309CO Warm Alpine occurs mountain-slopes, alluvial fan and valleys. Slopes is between 10 to 60%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived mostly from alluvium from rhyolite, slope alluvium from rhyolite, tuff, sandstone and volcanic rocks or colluvium from rhyolite, tuff, and volcanic rocks. Soil surface texture very gravelly loam, loam or very stony sandy loam with loamy--skeletal subsurface. It is Thurber's Fescue community. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.</p>
R048AY253CO	<p>Wet Subalpine R048AY253CO Wet Subalpine occurs on mesa tops and swales, drainageways, glacial moraines, and alluvial fans formed by glacial action. Slopes is between 0 to 30%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived from till from basalt, andesite, or tuff; slope alluvium from andesite or tuff, alluvium from andesite or tuff; eolian deposits from basalt; or residuum from basalt. Soil surface texture is loam, or gravelly loam with fine-loamy, loamy-skeletal or clayey-skeletal textured subsurface. It is slender wheatgrass – tufted hairgrass. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 37 to 45 inches.</p>

R048AY250CO	<p>Subalpine Loam</p> <p>R048AY250CO Subalpine Loam occurs on hills, mountain-slopes, and mountains. Slopes is between 1 to 30%. Soils are deep to very deep (20 to 60+ inches). Soils are derived from colluvium and alluvium from volcanic rock; complex landslide deposits from igneous, metamorphic, and sedimentary rock; and slope alluvium, colluvium, residuum, alluvium or complex landslide deposits from sandstone and shale or shale. Soil surface texture is loam with loamy textured subsurface. It is a mountain big sagebrush – Thurber’s Fescue community. It has an ustic udic/typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 20 to 30 inches.</p>
R048AY304CO	<p>Alpine Slopes</p> <p>R048AY304CO Alpine Slopes occurs on mountain-slopes, ridges, mountains and valleys. Slopes is between 30 to 60%. Soils are moderately deep to very deep (20 to 60+ inches). Soils are derived mostly from colluvium and alluvium from volcanic, metamorphic and igneous rocks. Soil surface texture is gravelly, very gravelly, extremely gravelly, very cobbly, extremely cobbly, or very stony loam or extremely stony sandy loam with loamy--skeletal subsurface. It is Kobresia – Tufted Hairgrass community. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.</p>
R048AY305CO	<p>Alpine Meadow</p> <p>R048AY305CO Alpine Meadow occurs on depressions, drainageways, cirques, flood plains and steams. Slopes is between 0 to 15%. Soils are greater than 60 inches in depth. Soils are derived mostly from colluvium from Monzonite, volcanic rock; alluvium from volcanic rocks; or slope alluvium from Andesite or latite. Soil surface texture is loam, cobbly loam or silt loam. The underlying material to a depth of 60 inches or more is variable material ranging from sandy loam to clay loam with variable amounts and sizes of rock fragments. It is Kobresia – Tufted Hairgrass community. This site has a water table at 0 to 36 inches in depth. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.</p>
R048AY306CO	<p>Shrubby Alpine</p> <p>R048AY306CO Shrubby Alpine occurs on cirque floors, mountains, mountain slopes and basin floor. Slopes is between 0 to 30%. Soils are greater than 60 inches in depth. Soils are derived mostly from till from conglomerate, sandstone or slope alluvium from basalt, rhyolite, andesite, tuff or volcanic breccia; or colluvium from andesite. Soil surface texture is very gravelly, cobbly, , or stony loam; or gravelly silt loam with loamy--skeletal subsurface. It is Kobresia – Tufted Hairgrass community. It has a Typic udic moisture regime and cryic temperature regime. The effective precipitation ranges from 30 to 50 inches.</p>

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Salix planifolia ssp. planifolia</i> (2) <i>Dryas octopetala ssp. hookeriana</i>
Herbaceous	(1) <i>Kobresia myosuroides</i> (2) <i>Deschampsia cespitosa</i>

Physiographic features

This site is in high mountainous country where physical and climatological conditions are extremely severe. Various phases of geological erosion are pronounced. Blowing soil particles and snow-blast injure plants; very high intensity sunlight prevails. This site is usually on slopes ranging from 5 to 60 percent. It occurs on all exposures. Elevation ranges from about 10,600 to 14,000 feet.

Table 2. Representative physiographic features

Landforms	(1) Mountain slope (2) Ridge (3) Basin floor (4) Mountain (5) Valley floor (6) Cirque floor
Runoff class	Medium to very high
Flooding frequency	None

Ponding frequency	None
Elevation	3,231–4,267 m
Slope	5–60%
Aspect	Aspect is not a significant factor

Climatic features

The climate is harsh and cold. Due to the elevation, summers are wet and short. The plant growth period is less than 60 days. The average annual precipitation ranges from 30 to 50 inches. Of this, approximately 80-90% falls as snow, and 10-20% falls as rain between middle of June to and the middle of September 1. Summer moisture is mostly from thundershowers in July, and August. The driest period is usually from May to June; and June is normally the driest month. The average annual total snowfall is 435 inches. The snow depth usually ranges from 16 to 67 inches during November thru May. The highest winter snowfall record in this area is 807 inches which occurred in 1978-1979. The lowest snowfall record is 205 inches during the 1995-1996 winter. The frost-free period typically ranges from 54 to 96 days. The last spring frost is the middle of June to the end of June. The first fall frost is the last week of August to the middle of September. Mean daily annual air temperature is about 21.5°F to 45.9°F, averaging about 21.5°F for the winter and 54.4°F in the summer. Summer high temperatures of mid-70°F to 80°F are not unusual. The coldest winter temperature recorded was -40°F on February 5, 1982 and the warmest winter temperature recorded was 63°F on February 20, 1958. The coldest summer temperature recorded was 9°F on June 26, 1969 and the warmest was 80 °F on July 6, 1989. The hottest day on record is 81°F on September 14, 1990. Wide yearly and seasonal fluctuations are common for this climatic zone. Data taken from Western Regional Climate Center (2018) for Wolf Creek Pass 1 E, Colorado Climate Station. There are only 2 climate stations in the zone in the state of Colorado. Wolf Creek Pass 1 E and Wolf Creek Pass 4 W; they are at the lower precipitation end of this zone.

The grasses generally start growth in late June. Generally forbs start their growth about June 15, while the shrub species start growth at the end of June. Plant growth usually stops in September. The optimum growth period is generally between July and September for most of these plants. c. The average annual temperature is 26 to 34°F. Frost can occur any day of the year during the growing season. Plant growth is seldom hampered by moisture stress but strong winds can induce drought-like conditions. Wind velocities for the area average 5 to 25 miles per hour. Strong winds during the winter and spring create snow cornices.

Table 3. Representative climatic features

Frost-free period (characteristic range)	51 days
Freeze-free period (characteristic range)	81 days
Precipitation total (characteristic range)	1,168 mm
Frost-free period (actual range)	51 days
Freeze-free period (actual range)	81 days
Precipitation total (actual range)	1,168 mm
Frost-free period (average)	51 days
Freeze-free period (average)	81 days
Precipitation total (average)	1,168 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) WOLF CREEK PASS 1 E [USC00059181], Creede, CO

Influencing water features

None

Soil features

The soils in this site are shallow, well-drained soils that formed in residuum and colluvium from volcanic and sedimentary rocks. Typically the surface is extremely stony loam. Available water capacity is very low. Effective rooting depth is 7 to 20 inches. Runoff is rapid, and the hazard of water erosion is high. Rock fragments make up 35 to 80 percent of the profile.

Soils Series associated with this site:

Granturk, Kite, Teewinot, Teewinot Family, Telluride, and Whitecross.

Table 4. Representative soil features

Parent material	(1) Slope alluvium–tuff (2) Slope alluvium–volcanic rock (3) Slope alluvium–rhyolite (4) Residuum–sandstone (5) Colluvium–rhyolite (6) Colluvium–tuff (7) Colluvium–volcanic rock (8) Colluvium–andesite (9) Slope alluvium–andesite (10) Colluvium–basalt (11) Slope alluvium–basalt (12) Colluvium–volcanic breccia (13) Slope alluvium–volcanic breccia (14) Till–andesite (15) Residuum–tuff (16) Residuum–rhyolite (17) Residuum–volcanic breccia (18) Residuum–basalt (19) Residuum–andesite (20) Residuum–volcanic rock (21) Residuum–limestone and sandstone (22) Residuum–metamorphic and sedimentary rock (23) Till–conglomerate (24) Till–sedimentary rock
Surface texture	(1) Very gravelly, very stony, very cobbly loam (2) Very stony, very cobbly sandy loam (3) Loam (4) Very cobbly silt loam
Family particle size	(1) Loamy-skeletal (2) Loamy
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderate to moderately rapid
Soil depth	18–51 cm
Surface fragment cover <=3"	0–30%
Surface fragment cover >3"	0–40%
Available water capacity (Depth not specified)	1.27–7.62 cm
Soil reaction (1:1 water) (Depth not specified)	4.5–6
Subsurface fragment volume <=3" (Depth not specified)	10–35%
Subsurface fragment volume >3" (Depth not specified)	5–40%

Ecological dynamics


The Shallow Alpine site is represented by an association of sedges, grasses, and forbs in which kobresia is dominant. This sedge, in the highest development of the climax, forms a sod which virtually excludes all other plants. Pure stands of kobresia seldom occur over large areas because of interruption by other plant communities which are determined by variations in exposure, slope, rock outcrops, and other physical and climatic features.

The state and transition model was added to fill the provisional ecological site instruction. It is a very general model. More field work and research is needed to make this model specific for each LRU (Land Resource Unit)

State and transition model

R048AY308CO Shallow Alpine

State 1: Reference State



Legend

1.1A – lack of fire, improper grazing, prolonged drought, time without disturbance

1.2A – disturbance, fire, insect herbivory of shrubs, proper grazing, wetter climate cycles

State 1 Reference State

Community 1.1 Reference State

The plant community is about 35 to 45 percent sedges, 25 to 30 percent grasses, 25 to 30 percent forbs, and 5 to 10 percent shrubs, air-dry weight. Kobresia, alpine fescue, bluegrass, and tufted hairgrass dominate the plant species, making up from 50 to 75 percent depending on the disturbance to the soil and kobresia community. The Elymus species appear on this site and aid the recognition of the Shallow Alpine Range Site. With a lack of disturbance, kobresia will tend to dominate areas within this site to the exclusion of all other species. The variety of grasses, forbs and shrubs adds color to the site, enriches the diet of grazing animals and enhances the wildlife habitat. If retrogression is caused by sheep, desirable grasses, forbs, and shrubs will be reduced. Deterioration will decrease kobresia, alpine fescue, arctiv and alpine bluegrass, desirable shrubs, and sedges. Cushion plants, sandwort phlox, dwarf or alpine clover, mountain avens, and sibbaldia will increase and occupy this site. Plant species likely to invade this site and become part of the plant community when the range is in a degenerated condition are mountain avens, sibbaldia (false-strawberry), yarrow, fleabane, low seedy cinquefoil, and pussytoes; or the aspect may be a uniform tufted hairgrass cover and other plants ecologically undesirable to the site. Basal

area (the are of ground surface covered by the perennial vegetation measured one inch above the soil) is approximately 30-40 percent. Annual Production: If the range is in excellent condition, the approximate annual production (air-dry) ranges are: Favorable years 1000 lb/ac Normal years 800 lb/ac Unfavorable years 600 lb/ac Of this production, 10-15 percent will likely be unpalatable or out of reach of grazing animals.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	437	583	729
Forb	168	224	280
Shrub/Vine	67	90	112
Total	672	897	1121

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Grasses and Grasslikes			493–673	
	Bellardi bog sedge	KOMY	<i>Kobresia myosuroides</i>	359–493	–
	tufted hairgrass	DECE	<i>Deschampsia cespitosa</i>	45–90	–
	alpine fescue	FEBR	<i>Festuca brachyphylla</i>	45–90	–
	spreading wheatgrass	ELSC4	<i>Elymus scribneri</i>	17–45	–
	slender wheatgrass	ELTRS	<i>Elymus trachycaulus ssp. subsecundus</i>	17–45	–
	alpine bluegrass	POAL2	<i>Poa alpina</i>	28–45	–
	arctic bluegrass	POAR2	<i>Poa arctica</i>	28–45	–
	alpine bentgrass	AGHU	<i>Agrostis humilis</i>	17–45	–
	Letterman's bluegrass	POLE3	<i>Poa lettermanii</i>	0–28	–
	spike trisetum	TRSP2	<i>Trisetum spicatum</i>	11–28	–
	timber oatgrass	DAIN	<i>Danthonia intermedia</i>	11–28	–
	alpine timothy	PHAL2	<i>Phleum alpinum</i>	11–28	–
	blackandwhite sedge	CAAL6	<i>Carex albonigra</i>	11–17	–
	southwestern showy sedge	CABE3	<i>Carex bella</i>	11–17	–
	sheep sedge	CAIL	<i>Carex illota</i>	11–17	–
	black sedge	CANO3	<i>Carex nova</i>	11–17	–
	Drummond's sedge	CARUD	<i>Carex rupestris var. drummondiana</i>	11–17	–
Forb					
2	Forbs			224–269	
	alpine clover	TRDA2	<i>Trifolium dasyphyllum</i>	17–90	–
	American bistort	POBI6	<i>Polygonum bistortoides</i>	11–45	–
	sticky polemonium	POVI	<i>Polemonium viscosum</i>	17–45	–
	Colorado ragwort	SESO	<i>Senecio soldanella</i>	11–45	–
	graylocks four-nerve daisy	TEGR3	<i>Tetraneuris grandiflora</i>	17–45	–
	alpine sagebrush	ARSC	<i>Artemisia scopulorum</i>	11–45	–

	Ross' avens	GERO2	<i>Geum rossii</i>	17–45	–
	snow cinquefoil	PONI2	<i>Potentilla nivea</i>	11–34	–
	alpine bluebells	MEAL7	<i>Mertensia alpina</i>	11–28	–
	Whipple's penstemon	PEWH	<i>Penstemon whippleanus</i>	11–28	–
	alpine springbeauty	CLME	<i>Claytonia megarhiza</i>	11–28	–
	showy draba	DRSP	<i>Draba spectabilis</i>	11–28	–
	western Indian paintbrush	CAOC4	<i>Castilleja occidentalis</i>	11–28	–
	Austrian draba	DRFL	<i>Draba fladnizensis</i>	11–28	–
	alpine dustymaiden	CHDOA2	<i>Chaenactis douglasii</i> var. <i>alpina</i>	0–17	–
	Rocky Mountain snowlover	CHJA	<i>Chionophila jamesii</i>	0–17	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0–17	–
	littleleaf pussytoes	ANMI3	<i>Antennaria microphylla</i>	0–17	–
	hairy arnica	ARMO4	<i>Arnica mollis</i>	0–17	–
	sulphur-flower buckwheat	ERUMM	<i>Eriogonum umbellatum</i> var. <i>majus</i>	0–17	–
	Parry's geranium	GECAP2	<i>Geranium caespitosum</i> var. <i>parryi</i>	0–17	–
	Colorado tansyaster	MACO13	<i>Machaeranthera coloradoensis</i>	0–17	–
	arctic alpine forget-me-not	ERNAE	<i>Eritrichium nanum</i> var. <i>elongatum</i>	0–11	–
	bluebell bellflower	CARO2	<i>Campanula rotundifolia</i>	0–11	–
	splitleaf Indian paintbrush	CARH4	<i>Castilleja rhexiifolia</i>	0–11	–
Shrub/Vine					
3	Shrubs			45–135	
	Hooker's mountain-avens	DROCH	<i>Dryas octopetala</i> ssp. <i>hookeriana</i>	11–45	–
	alpine willow	SAPE18	<i>Salix petrophila</i>	11–45	–

Animal community

This site provides excellent summer and fall forage for domestic sheep, bighorn sheep, and elk. The animal forage preference generally changes as the seasons progress into summer and fall. The dominant grasses or grasslike species, kobresia, alpine fescue, alpine bluegrass, and arctic bluegrass, and tufted hairgrass are desirable during the entire grazing season along with the excellent variety of forbs which this site provides.

Since sheep grazing occurs during the time of flowering and seed setting, a heavily grazed range may be marked by a scarcity of alpine forbs, especially the upright and relatively succulent species. Grasses, because of their higher palatability, also may be thinned out. On such a range, various alpine sedges and less palatable grasses are frequently the dominant vegetation. However, the natural succession is toward sedges and kobresia.

The date when alpine ranges are ready to graze varies with elevation, exposure, and seasonal climatic conditions. Generally speaking, most of these ranges should not be grazed before July 15 except for minor local areas in the subalpine area. In some places where snowbanks are prevalent, grazing may need to be deferred until as late as August 15. Grazing should not begin until the major feed areas are firm and free from excessive snow melt and water. Flower heads on alpine bluegrass and tufted hairgrass should be well developed.

Grazing after September 15 is hazardous because of the suddenness of fall storms. Because of these conditions, sheep allotments should be so designed that they will have one or more camps in suitable areas below timberline for grazing before July 15 and after September 1. If this is not feasible, the grazing season should be adjusted

accordingly.

Because of the short season above timberline, it is unavoidable that some areas will still have some saturated ground when sheep enter. It is important, therefore, that a rotation grazing plan be in effect. A simple but effective system is to reverse the band movement over the allotment during alternate years.

To the fullest extent possible, sheep should be kept off snowbanks and especially off the wet ground around snowbanks and off the saturated ground below. Certain local north and east slopes where snow may remain very late might well be closed to grazing.

Grazing should be once over, light, progressive, and without prolonged stops in favored areas. The practice of allowing sheep to browse their way back into willow fields for days or weeks in one location should not be tolerated. The practice of trailing sheep back into central bedgrounds is very destructive to the alpine sod. Progressive feeding and one night bedding should be the prevailing method of use.

Guide to Initial Stocking Rates:

The stocking rates given below are based on continuous use for the entire growing season and are intended only as an initial guide. Forage needs are calculated on the basis of 900 pounds of air-dry forage per animal unit month (AUM). To maintain proper use and allow for forage that disappears through trampling, small herbivore use, weathering, etc., 35 percent of the palatable forage produced is considered available for grazing by large herbivores.

Condition	Percent Climax	Class Vegetation	AUM/Ac	Ac/AUM
Excellent	76-100	.33	3	
Good	51-75	.25	4	
Fair	26-50	.16	6.25	
Poor	0-25	.08	12.55	

Adjustments to the initial stocking rates should be made as needed to obtain proper use. With specialized grazing systems, large livestock breeds, uncontrolled big game herbivores, inaccessibility, dormant season use, presence of introduced species, etc., stocking rate adjustments will be required.

If retrogression is caused by sheep, desirable grasses, forbs, and shrubs will be reduced. Deterioration will decrease kobresia, alpine fescue, arctic and alpine bluegrass, desirable shrubs, and sedges. Cushion plants, sandwort phlox, dwarf or alpine clover, mountain avens, and sibbaldia will increase and occupy this site.

Plant species likely to invade this site and become part of the plant community when the range is in a degenerated condition are mountain avens, sibbaldia (falsestrawberry), yarrow, fleabane, low seedy cinquefoil, and pussytoes; or the aspect may be a uniform tufted hairgrass cover and other plants ecologically undesirable to the site.

Basal area (the area of ground surface covered by the perennial vegetation measured one inch above the soil) is approximately 30-40 percent.

Annual Production

If the range is in excellent condition, the approximate total annual production (air-dry) ranges are:

Favorable years 1000 pounds/Ac
Normal years 800 pounds/Ac
Unfavorable years 600 pounds/Ac

Of this production, 10-15 percent will likely be unpalatable or out of reach of grazing animals.

This site is part of the normal habitat for Rocky Mountain goats and ptarmigan. Elk and bighorn sheep remain year long in some areas. More frequently these animals, along with deer, remain in this area only during the summer. Snowshoe rabbits, blue grouse, and pica, along with songbirds and other animals, depend on this site for their needs at least a part of the year.

As forage quality declines from overuse and excessive trailing, big game animals such as elk, deer, bighorn sheep,

and Rocky Mountain goats are the most affected species. Poor nutrition will result in declines of Rocky Mountain goats and big horn sheep. Elk and deer will move to other accessible forage if available in the vicinity. Ptarmigan, blue grouse, and snowshoe rabbits will also decline in numbers as range condition deteriorates. Pica, songbirds, and small mammals are the least affected by a reduction of the potential plant community.

Hydrological functions

Soils in this site are grouped into the "D" hydrologic group, as outlined in the Soils of Colorado Loss Factors and Erodibility Hydrologic Groupings 1979 Handbook. Field investigations are needed to determine hydrologic cover conditions and hydrologic curve numbers. Refer to NRCS National Engineering Handbook, Section 4, and Peak Flows in Colorado Handbook for more information.

Recreational uses

Recreation and Natural Beauty:

In the field of recreation, the alpine type has appeal and is a source of inspiration to many people through its attraction as an area for mountain climbing, photography, pack trips, and fishing in high elevation lakes. Over grazing and other uses that destroy the vegetation will impinge on recreational values, including use of alpine flower fields, strategic pastures for horse feed, and denudation of camp sites and lake shores. Hunting for ptarmigan and snowshoe rabbits is generally good to excellent on this site.

Wood products

No potential production on this range site.

Other information

Endangered Plants and Animals:

No endangered or threatened species have been identified. Species names will be included as reliable information becomes available.

The site occurs in Alamosa, Archuleta, Boulder, Chaffee, Clear Creek, Conejos, Costilla, Custer, Delta, Delores, Eagle, El Paso, Fremont, Garfield, Gilpin, Grand, Gunnison, Hinsdale, Huerfano, Jackson, Lake, La Plata, Larimer, Mesa, Mineral, Montezuma, Ouray, Park, Pitkin, Pueblo, Rio Blanco, Rio Grande, Routt, Saguache, San Juan, San Miguel, Summit, and Teller counties.

Major Poisonous Plants to Livestock:

There appear to be no major poisonous plants to livestock on this site.

Type locality

Location 1: La Plata County, CO	
General legal description	West side of Engineer Pass on Kindell Gulch Rd.

Other references

References

Chapman, S.S., G.E. Griffith, J.M. Omernik, A.B. Price, J. Freeouf, and D.L. Schrupp. 2006. Ecoregions of Colorado. (2-sided color poster with map, descriptive text, summary tables, and photographs). U.S. Geological Survey, Reston, VA. Scale 1:1,200,000.

Cleland, D.T.; Freeouf, J.A.; Keys, J.E.; Nowacki, G.J.; Carpenter, C.A.; and McNab, W.H. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Gen. Tech. Report WO-76D [Map on CD-ROM] (A.M. Sloan, cartographer). Washington, DC: U.S. Department of Agriculture, Forest Service, presentation scale 1:3,500,000; colored.

Soil Conservation Service (SCS). May 1987. Range Site Description for Shallow Alpine #308. : USDA, Denver Colorado.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Western Regional Climate Center. Retrieved from <http://www.wrcc.dri.edu/summary/Climsmco.html> on December 10, 2018

Contributors

Suzanne Mayne-Kinney

Approval

Kirt Walstad, 3/05/2024

Acknowledgments

Project Staff:

Suzanne Mayne-Kinney, Ecological Site Specialist, NRCS MLRA, Grand Junction SSO

Chuck Peacock, MLRA Soil Survey Leader, NRCS MLRA Grand Junction SSO

Program Support:

Rachel Murph, NRCS CO State Rangeland Management Specialist, Denver

Scott Woodhall, NRCS MLRA Ecological Site Specialist-QA Phoenix, AZ

Eva Muller, Regional Director, Rocky Mountain Regional Soil Survey Office, Bozeman, MT

B.J. Shoup, CO State Soil Scientist, Denver

Eugene Backhaus, CO State Resource Conservationist, Denver

Those involved in developing earlier versions of this site description include: Bob Rayer, retired NRCS Soil Scientist; Herman Garcia, retired CO State RMS and NRCS MLRA Ecological Site Specialist-QA Phoenix, AZ.

--Site Development and Testing Plan--:

Future work to validate and further refine the information in this Provisional Ecological Site Description is necessary. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data.

Additional information and data is required to refine the Plant Production and Annual Production tables for this ecological site. The extent of MLRA 48A must be further investigated.

Field testing of the information contained in this Provisional ESD is required. As this ESD is moved to the Approved ESD level, reviews from the technical team, quality control, quality assurance, and peers will be conducted.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	

Date	05/18/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-