

Ecological site R051XY264CO Chico Land

Last updated: 9/07/2023
Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 051X–High Intermountain Valleys

This MLRA encompasses the San Luis Valley in south central Colorado and the Taos Plateau and Taos alluvial fans of north central New Mexico. As part of the northern portion of the Rio Grande Rift, the MLRA consists of large, alluvium filled basins washed down from adjacent mountain ranges. The Rio Grande River flows through this MLRA, continuing its long function of carrying mountain sediment down to the basin. Cenozoic volcanism is an extensive characteristic of the MLRA where large basalt flows with volcanic hills and domes are abundant. Ancient Lake Alamosa is a large feature within the MLRA.

Classification relationships

NRCS:

Major Land Resource Area 51, High Intermountain Valleys (United States Department of Agriculture, Natural Resources Conservation Service, 2006).

USFS:

331J – Northern Rio Grande Basin M3311c > 331Ja - San Luis Valley, 331Jb - San Luis Hills and 331C - Mogotes

EPA:

22 - Arizona/New Mexico Plateau > 22a - San Luis Shrublands and Hills ; 22b -San Luis Alluvial Flats and Wetlands ; 22c - Salt Flats; 22e - Sand Dunes and Sand Sheets and 22f -Taos Plateau (Griffith, 2006).

USGS:

Southern Rocky Mountain Province

Ecological site concept

This site occurs on the broad central floor of the San Luis Valley. Elevation ranges from 7500 to 7750 feet. This site occurs in association with Salt Flats and Salt Meadow sites and is very similar to the Salt Flats site. Chico land exhibits extensive areas of "slick spots" and typically supports large stands of greasewood. Alkali sacaton, and inland saltgrass are dominant grasses. Soils are highly alkali with clayey subsoil which inhibits water infiltration and productivity.

Associated sites

R051XY312CO	Sand Hummocks Chico Land sites are found adjacent to and interspersed with Sand Hummocks along the eastern side of the San Luis Valley and throughout Alamosa County. The Chico Land exists on the alluvial flat while sand hummocks is part of the playa complex.
-------------	--

R051XY314CO	Alkali Overflow Chico Land sites are found adjacent to and on slightly higher ground than Alkali Overflow sites in Alamosa and Saguache Counties. In some areas around the Blanca Wetlands Chico Land sites intergrade into Alkali Overflow sites. Chico land resides on the alluvial flat while alkali overflow is associated with the playa complex.
R051XY263CO	Salt Flats Chico land is higher in clay content with greater slick spots, higher in alkalinity and is less vegetatively productive than salt flats.
R051XY294CO	Valley Sand Valley sand occurs throughout the alluvial flat of the basin floor where soils are coarse textured.
R051XY264CO	Chico Land
R051XY267CO	Salt Meadow Chico Land sites can be found adjacent to and on slightly higher ground than Salt Meadow sites. Salt Meadows have access to extra water due to higher flooding frequency and higher water table. Salt Meadows are grass dominated while chico land has a strong shrub component.

Similar sites

R051XY263CO	Salt Flats
R051XE260CO	Chico Fan 8-12 PZ
R051XY314CO	Alkali Overflow




Figure 1.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Sarcobatus vermiculatus</i>
Herbaceous	(1) <i>Sporobolus airoides</i> (2) <i>Distichlis spicata</i>

Physiographic features

This site occurs within the basin floor of the San Luis Valley. Elevation ranges from 7500 to 7800 feet. Landforms include flood plains on valley floors and alluvial flats.

Table 2. Representative physiographic features

Landforms	(1) Flood plain (2) Valley floor (3) Alluvial flat (4) Deflation basin
-----------	---

Runoff class	Medium
Flooding frequency	None
Ponding frequency	None
Elevation	2,286–2,377 m
Slope	0–2%
Water table depth	122–183 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate that typifies the High Intermountain Valley, ranges from arid to semi-arid, and is characterized by cold winters, moderate summers, and much sunshine. Average annual precipitation ranges from 6 to 10 inches along the valley floor and throughout most of the resource area. Approximately 55-60 percent of the annual precipitation falls between May 1 and September 1. May and June are normally dry. Precipitation comes mostly from short duration high intensity thundershowers in July and August. Wide seasonal and yearly variations are common. The San Juan mountain range to the west and the Sangre de Cristo Mountains to the east intercept much of the precipitation causing a two-way rain shadow effect.

Cold air from the encompassing mountain ranges drain into the valley and settle. This phenomena results in long cold winters and moderate summer temperatures. Mean average annual temperature ranges between 42 to 44 degrees F. July is the hottest month and January is the coldest. Summer temperatures range from highs in the upper 70's and low 80's and occasionally reach to the mid 90 degrees F. Summer nights are cool. Average frost-free period is 90-115 days, from late May or early June to September. There is a 50% probability that the first frost in the fall will occur near September 16, and the last frost in the spring on about June 9. Most major plant species initiate growth between mid May and late July, but growth may extend into September. Some cool season plants begin growth earlier and complete growth by mid June. There may be late re-growth on some of the plants.

Wind speeds average 7 miles per hour annually. Wind that often reaches high velocities are common, especially in the spring. Relative humidity is usually low. Even so, evaporation rates average lower than those of many dry regions because of the cooler climate. Snow cover is often light and is sometimes lacking through much of the winter. There is usually some snow, though, during the coldest weather.

Table 3. Representative climatic features

Frost-free period (characteristic range)	70-83 days
Freeze-free period (characteristic range)	97-107 days
Precipitation total (characteristic range)	152-254 mm
Frost-free period (actual range)	69-83 days
Freeze-free period (actual range)	94-108 days
Precipitation total (actual range)	152-254 mm
Frost-free period (average)	76 days
Freeze-free period (average)	102 days
Precipitation total (average)	203 mm




Figure 2. Monthly precipitation range




Figure 3. Monthly minimum temperature range




Figure 4. Monthly maximum temperature range




Figure 5. Monthly average minimum and maximum temperature




Figure 6. Annual precipitation pattern




Figure 7. Annual average temperature pattern

Climate stations used

- (1) ALAMOSA SAN LUIS AP [USW00023061], Alamosa, CO
- (2) WAVERLY 1W [USC00058860], Alamosa, CO
- (3) MANASSA [USC00055322], La Jara, CO
- (4) MONTE VISTA 2W [USC00055706], Monte Vista, CO
- (5) CENTER 4 SSW [USC00051458], Center, CO

Influencing water features

Seasonal high water table exists from 4 to 6 feet below the surface. The water table can be found usually June thru September. Lower areas may receive low amounts of run-on from adjacent higher areas.

Soil features

Soil surface textures are fine-loamy. Clay content in the surface ranges from 25 to 35%. They are strongly alkaline. They are usually underlain by sand and gravel and this shows are a restrictive feature in the soil at 20 to 40 inches (50 to 100 cm) . Large areas of "slick spots" are mottled throughout the site where particles are highly dispersed causing impermeable soils with virtually no penetration of water. Runoff from rain or snow melt sit in these lower slick spot areas until evaporation occurs.

Typical soil correlated to this site is Hooper.

Table 4. Representative soil features

Parent material	(1) Alluvium–igneous and metamorphic rock (2) Alluvium–volcanic rock
Surface texture	(1) Clay loam
Family particle size	(1) Fine-loamy over sandy or sandy-skeletal
Drainage class	Well drained


Permeability class	Very slow to slow
Depth to restrictive layer	51–102 cm
Soil depth	152–305 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	9.65–13.21 cm
Calcium carbonate equivalent (Depth not specified)	0–5%
Electrical conductivity (Depth not specified)	4–8 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0–1
Soil reaction (1:1 water) (Depth not specified)	8.5–9.6
Subsurface fragment volume <=3" (Depth not specified)	0–10%
Subsurface fragment volume >3" (Depth not specified)	0–2%

Ecological dynamics


The structural-functional groups are mostly a mix of shrubs and grasses with a few forbs. The plant community in the sodium dispersed clay areas or slick spots is almost exclusively black greasewood with occasional patches of inland saltgrass. Alkali sacaton is the dominant grass with patches of inland saltgrass. Grasses which occur in minor amounts include alkali cordgrass, creeping (alkali) wildrye, mat muhly, and an occasional plant of blue grama. Baltic rush occurs in the wetter areas which receive runoff from rain or snowmelt. Rubber rabbitbrush occurs with black greasewood where the loamy sand surface is deeper. Green rabbitbrush and an occasional prickly pear plant may occur on higher knolls.

State and transition model


Ecosystem states



State 1 submodel, plant communities



State 2 submodel, plant communities



State 1

Reference: Shrub-Grass mix



Figure 8. Reference: shrub-grass mix

The reference state provides the greatest amount of ecological resilience. The plant community mix slightly favors shrubs over grasses with air-dry weight. Vegetation is patchy with open slick spot areas present throughout the site. Black greasewood is the dominant shrub while alkali sacaton is the dominant grass. The soil is stable with the A horizon intact.

Dominant plant species

- greasewood (*Sarcobatus vermiculatus*), shrub
- alkali sacaton (*Sporobolus airoides*), grass

Community 1.1

Reference: Black Greasewood/Alkali Sacaton



Figure 9. Reference



Figure 10. Bare ground or slick spots forming continuous patches.

Alkali Sacaton is the dominant grass, with associated inland saltgrass. Grasses which occur in minor amounts include: alkali cordgrass, alkali wildrye, western wheatgrass, mat muhly, and an occasional blue grama plant. Bare ground or slick spots are mottled through the site. Greasewood is the dominant shrub.

Dominant plant species

- greasewood (*Sarcobatus vermiculatus*), shrub
- rubber rabbitbrush (*Ericameria nauseosa*), shrub
- alkali wildrye (*Leymus simplex*), shrub
- alkali sacaton (*Sporobolus airoides*), grass
- saltgrass (*Distichlis spicata*), grass

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	157	224	291
Grass/Grasslike	112	168	224
Forb	11	28	45
Total	280	420	560




Figure 12. Plant community growth curve (percent production by month). CO5104, Warm season dominant, cool season sub-dominant MLRA-51; valley floor- fine textured soils.


Community 1.2 Shrub Dominant



Figure 13. Shrub dominant

This community phase is dominated by shrubs, especially greasewood. Rubber rabbitbrush has also increased on the site. Grasses such as alkali sacaton have decreased. Overall annual production is less.

Pathway P1.1A Community 1.1 to 1.2



Reference: Black Greasewood/Alkali Sacaton

Shrub Dominant

Time and colonization of shrub species plus repetitive, high utilization of palatable species that does not allow for the regrowth and recovery during the growing season. These drivers slowly decrease grass cover and density while shrubs gain a competitive advantage.

Pathway P1.2A Community 1.2 to 1.1



Shrub Dominant



Reference: Black Greasewood/Alkali Sacaton

An event to set the shrubs back, such as disease, drought, fire, mechanical, or chemical. This coupled with managed grazing that takes into account critical growth periods for warm and cool season grasses plus the monitoring of intensity and frequency of grazing events will help in the building of soil and colonization of plants.

State 2 Degraded



Figure 14. Degraded Chico Land

Palatable species such as alkali sacaton, western wheatgrass, and alkali cordgrass have become a remnant or are no longer present. The degraded state is sparsely dominated by shrubs such as greasewood and rubber rabbitbrush. There may be a remnant of inland saltgrass, and annuals. The degraded state is greater than 70 percent bare ground. Erosion and deposition have a significant negative impact, creating feedback loops that substantially decreases hydrologic function.

Community 2.1 Eroded



Figure 15. Unstable soil

This community phase has a loss of topsoil due to erosion and absence of plant cover.

Community 2.2 Early Recovery



Figure 16. A wet spring showing shoots of alkali wildrye and alkali chardgrass, along with Russian knapweed.

This community has much bare ground and unstable soil but has started the restoration process with a small percentage of perennial, native plant species.

Resilience management. This is a fragile time where additional disturbance may put an end to restoration efforts.

Pathway P2.1 Community 2.1 to 2.2



Eroded



Early Recovery

Timely precipitation events in the spring and early summer, coupled with rest from grazing may allow perennial, native species to begin colonization.

Pathway P2.2 Community 2.2 to 2.1



Early Recovery




Eroded

After early signs of restoration further disturbance dries up the young shoots and roots of reference species and the site reverts back to the eroded state.

State 3 Irrigation tail water influenced state

Sometimes this site gets regular saturation from stream overflow or irrigation. If this happens for multiple years both black greasewood and alkali sacaton will decrease and inland saltgrass and baltic rush will increase. Greasewood does not mind being flooded unless the water sits for over 40 days, then it will begin die-off. Alkali sacaton will die off if water sits for over 3 weeks, continuously. Noxious weeds such as tall whitetop and perennial pepperweed may invade and dominate areas.

Transition T1A State 1 to 2




A major long-term driver is repetitive defoliation at high utilization for plants throughout multiple growing seasons and especially during drought. Plant species such as alkali sacaton, alkali cordgrass, and western wheatgrass are greatly decreased and may be lost. Greasewood also gets grazed and starts to lose vitality over time. This in turn increases both erosion and deposition, causing a decrease in soil stability and hydrologic function. A trigger event, such as drought may snap the site across the degraded threshold by causing excessive mortality to plants with low vigor.

**Transition T1B
State 1 to 3**

A regular accumulation of water over multiple years causes a species shift to more water tolerant plants such as inland saltgrass and baltic rush.

**Restoration pathway R2A
State 2 to 1**



Restoration will take a long time as the site itself is harsh on plants and with an decrease in soil stability due to erosion hydrologic function is very low. A slow process of plant and soil building is necessary to restore ecological processes. In theory, renewed grazing management can be part of the restoration process. It would require planning and monitoring to allow for moderate frequency and intensity along with adequate recovery periods during the growing season. An abundance of litter and cover must be left during the dormant season to aid in moisture retention and slow erosion. Monitoring for plant and soil ecological processes is important for restoration success.

**Restoration pathway R3A
State 3 to 1**

An end to water saturation followed by years of grazing management and monitoring that allows for the colonization of reference species. This state may take many years to restore depending on the amount of soil erosion and accumulated salts form years of irrigation.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1				140–196	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	90–135	–
	saltgrass	DISP	<i>Distichlis spicata</i>	45–56	–
	alkali wildrye	LESI5	<i>Leymus simplex</i>	6–17	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–9	–
	alkali cordgrass	SPGR	<i>Spartina gracilis</i>	0–9	–
	mat muhly	MURI	<i>Muhlenbergia richardsonis</i>	0–9	–
	mountain rush	JUARL	<i>Juncus arcticus ssp. littoralis</i>	0–9	–
	needleleaf sedge	CADU6	<i>Carex duriuscula</i>	0–4	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0–4	–
	wildrye	ELYMU	<i>Elymus</i>	0–4	–
Forb					
2				11–45	
	western tansymustard	DEPI	<i>Descurainia pinnata</i>	0–4	–
	seepweed	SUAED	<i>Suaeda</i>	0–2	–
	tanseyleaf tansyaster	MATA2	<i>Machaeranthera tanacetifolia</i>	0–2	–
	Pursh seepweed	SUCA2	<i>Suaeda calceoliformis</i>	0–2	–
	red swampfire	SARU	<i>Salicornia rubra</i>	0–2	–
Shrub/Vine					
3				168–280	
	greasewood	SAVE4	<i>Sarcobatus vermiculatus</i>	179–269	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	0–22	–
	Greene's rabbitbrush	CHGR6	<i>Chrysothamnus Greenei</i>	0–9	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–4	–

Animal community

This site offers only limited value for livestock grazing because of the low production, high percentage of unpalatable species, and because of the vast areas of "slick spots" which have little or no vegetation. These grasses and shrubs can provide fair nutrition for grazing animals, however, if properly managed. Care should be exercised when grazing pastures which are dominated by this site, to avoid nutritional stress.

A planned grazing system should be used which benefits both early and later season grasses. Grazing periods should be kept relatively short, and adequate rest provided prior to re-grazing. This type of rotational grazing, if carefully monitored, will provide the highest level of production and nutrition for grazing animals. Spring rest for May 1 to June 20 benefits cool-season plants such as western wheatgrass and creeping wildrye. While rest from June through July would be most beneficial to alkali sacaton and alkali cordgrass. Dormant season grazing greatly favors the plants of this site and allows for recovery during the growing season. Chemical brush control would effectively reduce black greasewood, however, adequate grass cover of alkali sacaton should exist prior to spraying. If rubber rabbitbrush is part of the community, the brush control must be conducted just as the rabbitbrush is beginning to flower, otherwise it will respond to the disturbance with an increase in population. A least greasewood provides good browse. Most of this site, consequently, is not suited for brush management. Any attempts at reseeding would result in total failure.

b. Guide to Initial Stocking Rates:

(1) Stocking rates given below are based on continuous use for the entire growing season, and are intended only as

an initial guide. Forage needs are calculated on the basis of 900 lbs of air-dry forage per animal unit month (AUM). To maintain proper use and allow for forage that disappears through trampling, small herbivore use, weathering, etc., 35% of the palatable forage produced is considered available for grazing by large herbivores.

Condition Class/ % Climax Vegetation/ Ac/AUM 1/ AUM/Ac 1
excellent/ 76-100/ 15/ .07
good/ 51-75/ 25/ .04
fair/ 26-50/ 40/ .03
poor/ 0-25/ 55/ .02

Due to the high amount of bare ground in the mapping units, the stocking rates have been adjusted accordingly.

Adjustments to the initial stocking rates should be made as needed to obtain proper use. With specialized grazing systems, large livestock breeds, uncontrolled big game herbivores, inaccessibility, dormant season use, presence of introduced species, etc., stocking rate adjustments will be required.

Major Poisonous Plants to Livestock:

Plant Common Name/ Livestock Affected/ Type of Poisoning/ Season Serious
Black greasewood/ sheep, cattle/ acute/ spring

The relative value of this site for wildlife is low. As ecological degradation occurs, however, food availability for jackrabbits and other small herbivores would decrease. Value for habitat would not be decreased.

Hydrological functions

Soils in this site are grouped into "D" hydrologic group, as outlined in the "Soils of Colorado Loss Factors and Erodibility Hydrologic Groupings" handbook. Field investigations are needed to determine hydrologic cover conditions and hydrologic curve numbers. Refer to "Peak Flows in Colorado" handbook, and SCS "National Engineering Handbook", Section 4, for hydrologic curve numbers and determining runoff quantities.

Recreational uses

Recreation values are limited to hunting of rabbit species and the incidental use of this site in hunting or observing waterfowl near artesian wells, and other wetland areas associated with this site.

Inventory data references

Counties where this ecological site occurs include:
Alamosa, Conejos; Costilla. Rio Grande and Saguache

Field Offices in Colorado where the site occurs:
Alamosa, San Luis, and Center

References

. 2021 (Date accessed). USDA PLANTS Database. <http://plants.usda.gov>.

Other references

Chapman, S.S., G.E. Griffith, J.M. Omernik, A.B. Price, J. Freeouf, and D.L. Schrupp. 2006. Ecoregions of Colorado. (2-sided color poster with map, descriptive text, summary tables, and photographs). U.S. Geological Survey, Reston, VA. Scale 1:1,200,000.

Cleland, D.T.; Freeouf, J.A.; Keys, J.E.; Nowacki, G.J.; Carpenter, C.A.; and McNab, W.H. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Gen. Tech. Report WO-76D [Map on

CD-ROM] (A.M. Sloan, cartographer). Washington,DC: U.S. Department of Agriculture, Forest Service, presentation scale 1:3,500,000; colored.

Natural Resources Conservation Service (NRCS). 2017. Ecological Site Description for Chico Land #264(ESIS) : USDA, Denver Colorado.

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Contributors

C. Villa, H. Garcia, S. Woodall
Suzanne Mayne-Kinney

Approval

Kirt Walstad, 9/07/2023

Acknowledgments

Project Staff:

Suzanne Mayne-Kinney, Ecological Site Specialist, NRCS MLRA, Grand Junction SSO
Alan Stuebe, MLRA Soil Survey Leader, NRCS MLRA Alamosa SSO

Program Support:

Rachel Murph, NRCS CO State Rangeland Management Specialist, Denver
Eva Muller, Regional Director, Rocky Mountain Regional Soil Survey Office, Bozeman, MT
B.J. Shoup, CO State Soil Scientist, Denver
Eugene Backhaus, CO State Resource Conservationist, Denver

--Site Development and Testing Plan--:

Future work to validate and further refine the information in this Provisional Ecological Site Description is necessary. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data.

Additional information and data are required to refine the Plant Production and Annual Production tables for this ecological site. The extent of MLRA 51 must be further investigated.

Field testing of the information contained in this Provisional ESD is required. As this ESD is moved to the Approved ESD level, reviews from the technical team, quality control, quality assurance, and peers will be conducted.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/19/2024
Approved by	Kirt Walstad
Approval date	

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-