

Ecological site R051XY273CO Sandy Bench

Last updated: 7/19/2021
 Accessed: 04/24/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	S. Woodall, C. Villa, D. Sparks, H. Garcia
Contact for lead author	
Date	12/16/2004
Approved by	Curtis Talbot
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None

2. **Presence of water flow patterns:** None

3. **Number and height of erosional pedestals or terracettes:** Minor pedestalling may be evident near wind scoured areas.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** A fair amount of gravel and cobble stones are scattered about the soil surface. Approximately 15% or less bare ground, with bare patches ranging from 2-3 inches in diameter. Prolonged drought or wildfire events will cause bare ground to increase upwards to 15-20% with bare patches ranging from 3-5 inches in diameter.

5. **Number of gullies and erosion associated with gullies:** None

6. **Extent of wind scoured, blowouts and/or depositional areas:** Some minor wind scouring is inherent to this site. Surface gravels provide surface roughness. Soil movement can intensify with disturbances such as wildfire, wildlife, or extended drought.

-
7. **Amount of litter movement (describe size and distance expected to travel):** Litter movement on this site is minor.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability class rating anticipated to be 3-4 in interspace at soil surface.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface soils are most commonly loamy sand, fine sandy loam or sandy loam. The A-horizon ranges from 2-15 inches in depth. Surface horizon structure is usually weak fine granular structure.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Plant community composition has a minor effect on infiltration and runoff on this site. Infiltration rates are high due to soil texture.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: cool season bunchgrass >
- Sub-dominant: warm season bunchgrass > shrubs > cool season rhizomatous grass >
- Other: forbs
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Minimal. Decadence will exist on areas excluded from grazing animals.
-
14. **Average percent litter cover (%) and depth (in):** 25-35% litter cover at 0.25 inch depth. Litter cover during and following drought can range from 10-15% and 5-10% following wildfire.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 800 lbs./ac. low precipitation years; 1000 lbs./ac. average precipitation years; 1200 lbs./ac. high precipitation years. After extended drought, production will be reduced to 400 – 600 lbs./ac. or more.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize**

degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: Invasive plants should not occur in reference plant community.

17. **Perennial plant reproductive capability:** The only limitations are weather-related, wildfire, natural disease, and insects that may temporarily reduce reproductive capability.
-