

Ecological site R053BY010ND Shallow Gravel

Last updated: 1/11/2024 Accessed: 05/20/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Printz, Stan Boltz, Lee Voigt, Jody Forman
Contact for lead author	Jeff.printz@nd.usda.gov 701-530-2080
Date	03/09/2012
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1.	Number and extent of rills: None.
2.	Presence of water flow patterns: None.
3.	Number and height of erosional pedestals or terracettes: None.
4.	Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 5 to 15%.
5.	Number of gullies and erosion associated with gullies: None.
6.	Extent of wind scoured, blowouts and/or depositional areas: None.

7. Amount of litter movement (describe size and distance expected to travel): Plant litter remains in place. Slight movement may be visible following intense thunderstorm events particularly after extended periods of below normal

8.	Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Soil aggregate stability ratings should typically be 5 or greater. Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.
9.	Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use soil series description for depth, color and structure of A horizon/surface layer.
10.	Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Combination of shallow and deep rooted species (mid rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
11.	Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to): Dominant: Mid and short cool-season bunchgrasses >
	Sub-dominant: Short warm-season grasses
	Other: Mid warm-season bunchgrasses = grass-likes > forbs = shrubs > cryptogam
	Additional: Due to differing root structure and distribution, Kentucky bluegrass and smooth bromegrass do not fit into reference plant community F/S groups.
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): None.
14.	Average percent litter cover (%) and depth (in): In contact with soil surface.
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): Representative value = 1600 lbs./ac with a range of 800 to 2100 lbs./ac air dry depending upon growing conditions.
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not

precipitation.

	for the ecological site: State and local noxious, Kentucky bluegrass, smooth bromegrass
7.	Perennial plant reproductive capability: All species exhibit high vigor relative to climatic conditions. Do not rate base solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.