

## Ecological site R053BY014ND Choppy Sands

Last updated: 1/11/2024  
Accessed: 05/14/2024

---

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|   |   |
|---|---|
| Author(s)/participant(s)                    | Jeff Printz, Stan Boltz, Lee Voigt, Jody Forman |
| Contact for lead author                     | Jeff.printz@nd.usda.gov 701-530-2080            |
| Date  | 03/02/2012                                      |
| Approved by                                 | Suzanne Mayne-Kinney                            |
| Approval date                               |   |
| Composition (Indicators 10 and 12) based on | Annual Production                               |

### Indicators

- Number and extent of rills:** None.  

---
- Presence of water flow patterns:** None.  

---
- Number and height of erosional pedestals or terracettes:** Some pedestalling of bunchgrasses may be observable but roots will not be exposed.  

---
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 15 to 20%. Patch size is 6 inches or less and not connected.  

---
- Number of gullies and erosion associated with gullies:** None.  

---
- Extent of wind scoured, blowouts and/or depositional areas:** Occasional small (< a few feet across) areas of blowouts/wind scoured areas with associated depositional areas associated with increased animal activity (i.e. rodent burrows, animal trailing). The areas may increase in number and size following prolonged drought.  

---

7. **Amount of litter movement (describe size and distance expected to travel):** Slight movement of small sized litter may be observable on slopes greater than 20%.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability averages 5 or greater.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use soil series description for depth, color and structure of A horizon/surface layer.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Buried A horizon may be present but does not constitute a compaction layer.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Tall, warm-season rhizomatous grasses >
- Sub-dominant: Mid, cool-season bunchgrasses >
- Other: Grass-likes = forbs > short, warm-season grasses = mid, cool-season rhizomatous grasses = shrubs > short, cool-season grasses
- Additional: Due to differing root structure and distribution, Kentucky bluegrass and smooth brome grass do not fit into reference plant community F/S groups.
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** None.
- 
14. **Average percent litter cover (%) and depth ( in):** Plant litter is in contact with soil surface.
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Representative value = 1700 lbs./ac air dry with a range of 1100 to 2300 lbs./ac air dry depending upon growing conditions.
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if**

**their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious weeds, Kentucky bluegrass, smooth brome grass

---

17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.
-