

Ecological site R053CY019SD Closed Depression

Last updated: 1/22/2024
Accessed: 05/06/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 053C–Southern Dark Brown Glaciated Plains

The Southern Dark Brown Glaciated Plains (53C) is located within the Northern Great Plains Region. It is entirely in South Dakota encompassing about 3,990 square miles (Figure 1). The elevation ranges from 1,300 to 2,300 feet. The MLRA is level to gently rolling till plains including many areas of potholes. A terminal moraine occurs in the southern end of the MLRA. Moderately steep and steep slopes are adjacent to the major valleys. The headwaters of many creeks in central South Dakota occur in the high-lying MLRA. (USDA-NRCS 2006).

The dominant soil orders in this MLRA are Mollisols and Inceptisols. The soils in the area dominantly have a mesic soil temperature regime, an ustic soil moisture regime, and mixed or smectitic mineralogy. They generally are very deep, well drained or moderately well drained, and are loamy or clayey. This area supports natural prairie vegetation characterized by western wheatgrass (*Pascopyrum smithii*), big bluestem (*Andropogon gerardii*), needleandthread (*Hesperostipa comata*), and green needlegrass (*Nassella viridula*). Little bluestem (*Schizachyrium scoparium*), sideoats grama (*Bouteloua curtipendula*), and prairie sandreed (*Calamovilfa longifolia*) are important species on steeper sites. Western snowberry (*Symphoricarpos occidentalis*) and prairie rose (*Rosa arkansana*) are commonly dispersed throughout the area. (USDA-NRCS 2006).

Classification relationships

Major Land Resource Area (MLRA): Southern Dark Brown Glaciated Plains (53C) (USDA-NRCS 2006)

USFS Subregions: Northeastern Glaciated Plains Section (331E); Missouri Coteau Subsection (331Ea); Western Great Plains Section (331F); Missouri Breaks Subsection (331Fe); Western Glaciated Plains Section (332B); Southern Missouri Coteau Slope Subsection (332Bd, 332Be); North Central Great Plains Section (332D); Southern Missouri Coteau Slope Subsection (332Dd); Southern Missouri Coteau Subsection (332De) - (Cleland et al. 2007).

US EPA Level IV Ecoregion: Missouri Coteau (42a); Southern Missouri Coteau (42e); Southern Missouri Coteau Slope (42f) - (USEPA 2013)

Ecological site concept

The Closed Depression ecological site typically occurs in slight depressions on nearly level slopes in the upland areas. Soils are poorly drained and may have a claypan (columnar structure) within 6 inches of the soil surface or an abrupt texture change within 12 inches of the soil surface. Permeability is very slow due to the claypan (columnar structure) or the clayey subsoil and the site may pond water 4 to 8 weeks in the spring of the year. Pondered water conditions and very slow permeability or a natric horizon separately or the combination of both together strongly influences the soil-water-plant relationship. The natric horizon in the subsoil typically has a Sodium Absorption Ratio (SAR) greater than 13 or an Exchangeable Sodium Percentage (ESP) greater than 15 separately or the combination of both together. Vegetation in the Reference State includes western wheatgrass and common spikerush. Common forbs on this site include American licorice, smartweeds, and knotweeds. The site may become degraded due to change in disturbance regime, and vegetation may shift to community dominated by foxtail barley, inland saltgrass, and bareground.

Associated sites

R053CY010SD	Loamy These sites occur on upland areas. The soils are well drained and have less than 40 percent clay in the surface and subsoil. The central concept soil series are Agar, Glenham, and Highmore, but other series are included.
R053CY011SD	Clayey These sites occur on upland areas. The soils are well drained and have greater than 40 percent clay in the surface and subsoil. The central concept soil series are Demky, Oko, and Raber, but other series are included.
R053CY013SD	Claypan These sites occur on uplands. Soils are moderately well drained and have a claypan (columnar structure) between 6 and 16 inches from the soil surface. The central concept soil series are Cavo and DeGrey, but other series are included.

Similar sites

R053CY004SD	Wet Meadow The Wet Meadow site occurs in a similar landscape position and does not have a claypan (columnar structure) within 6 inches of the soil surface or an abrupt texture change within 12 inches of the soil surface.
R053CY015SD	Thin Claypan The Thin Claypan site occurs in a similar landscape position, is moderately well drained, and has a claypan (columnar structure) within 6 inches of the soil surface.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Pascopyrum smithii</i> (2) <i>Hordeum jubatum</i>

Physiographic features

This site occurs on nearly level depressions.




Figure 2. Distribution map

Table 2. Representative physiographic features

Landforms	(1) Pothole (2) Depression
Flooding frequency	None
Ponding duration	Long (7 to 30 days)
Ponding frequency	Frequent
Elevation	396–701 m
Slope	0–1%
Ponding depth	0–30 cm
Water table depth	0–203 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 53C is considered to have a continental climate – cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the Northern Great Plains and air masses move freely across the plains and account for rapid changes in temperature.

Annual precipitation typically ranges from 15 to 25 inches per year. The average annual temperature is about 45°F. January is the coldest month with average temperatures ranging from about 15°F (Stephan, South Dakota (SD)), to about 16°F (Onida 4 NW, SD). July is the warmest month with temperatures averaging from about 72°F (Stephan, SD), to about 74°F (Onida 4 NW, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 58°F. This large annual range attests to the continental nature of this area's climate. Hourly winds are estimated to average about 12 miles per hour (mph) annually, ranging from about 13 mph during the spring to about 11 mph during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 mph.

Growth of cool-season plants begins in early to mid-March, slowing or ceasing in late June. Warm-season plants begin growth about mid-May and continue to early or mid-September. Greenup of cool-season plants may occur in September and October when adequate soil moisture is present.

Table 3. Representative climatic features

Frost-free period (characteristic range)	107-127 days
Freeze-free period (characteristic range)	128-150 days
Precipitation total (characteristic range)	508-533 mm
Frost-free period (actual range)	104-129 days
Freeze-free period (actual range)	127-159 days
Precipitation total (actual range)	483-610 mm
Frost-free period (average)	117 days
Freeze-free period (average)	139 days
Precipitation total (average)	533 mm

Climate stations used

- (1) GETTYSBURG 13W [USC00393302], Gettysburg, SD
- (2) GETTYSBURG [USC00393294], Gettysburg, SD
- (3) HIGHMORE 23 N [USC00393838], Highmore, SD
- (4) ONIDA 4 NW [USC00396292], Onida, SD
- (5) PIERRE RGNL AP [USW00024025], Pierre, SD
- (6) HARROLD 12 SSW [USC00393608], Pierre, SD
- (7) STEPHAN 2 NW [USC00397992], Highmore, SD
- (8) WESSINGTON SPRINGS [USC00399070], Wessington Springs, SD

Influencing water features

No riparian areas or wetland features are directly associated with this site.

Soil features

The common features of soils in this site are the clay to clay loam textured subsoil and slopes of zero to one percent. The soils in this site are poorly drained and formed in alluvium or alluvium over till. The silt loam surface layer is three to nine inches thick. Some soils exhibit an extremely hard clayey Btn horizon that has round-topped or bun shaped columnar structure. These Btn horizons are high in sodium. The soils have a very slow infiltration rate. Available water capacity is five to six inches. The soils crack when dry and heavy traffic can cause surface compaction when wet. Subsurface soil layers are restrictive to water movement and root penetration. This site should show slight to no evidence of rills, wind scoured areas, or pedestalled plants. Water flow paths are broken, irregular in appearance, or discontinuous. The soil surface is stable and intact. The central concept soil series are Hoven and Plankinton, but other series are included.

High accumulations of sodium and slow permeability strongly influence the soil-water-plant relationship on this site. Access Web Soil Survey (<http://websoilsurvey.nrcs.usda.gov/app/>) for specific local soils information.

Table 4. Representative soil features

Surface texture	(1) Silt loam (2) Silty clay loam (3) Clay
Family particle size	(1) Clayey
Drainage class	Poorly drained
Permeability class	Very slow
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%

Available water capacity (0-101.6cm)	12.7–17.78 cm
Calcium carbonate equivalent (0-101.6cm)	1–15%
Electrical conductivity (0-101.6cm)	0–16 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–20
Soil reaction (1:1 water) (0-101.6cm)	5.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–5%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

State and Community Phases

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

The site which is located in the Southern Dark Brown Glaciated Plains Region developed under Northern Great Plains climatic conditions and included natural influence of large herding herbivores and occasional fire. Changes will occur in the plant communities due to weather fluctuations or management actions separately or the combination of both together. Under adverse impacts, a relatively rapid decline in vegetative vigor and composition can occur. Under favorable conditions the site has the potential to resemble the Reference State. Interpretations for this site are based primarily on the 1.1 Reference Plant Community Phase. This community phase and the Reference State have been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been considered. Plant community phases, states, transitional pathways, and thresholds have been determined through similar studies and experience.


This site is very sensitive to precipitation fluctuations from year to year. With above average precipitation the site becomes very wet, leading to a much different plant community than what would be present with average to below average precipitation. In dry years, plant density becomes very low. The two plant communities influenced strongly by precipitation alone, Western Wheatgrass-Common Spikerush Plant Community Subphase and Common Spikerush-Foxtail Barley Plant Community Subphase make up the natural fluctuation of what could be considered the 1.1 Reference Plant Community Phase.

Following the state and transition diagram are narratives for each of the described states and community phases. These may not represent every possibility, but they are the most prevalent and repeatable states/community phases. The plant composition tables shown below have been developed from the best available knowledge at the time of this revision. As more data are collected, some of these community phases or states separately or the combination of both together may be revised or removed, and new ones may be added. The main purpose for including the descriptions here is to capture the current knowledge and experience at the time of this revision.

The following is a diagram that illustrates the common plant community phases that can occur on the site and the transition and community pathways between them. The ecological processes will be discussed in more detail in the plant community descriptions following the diagram.

State and transition model

Closed Depression – R053CY019SD



LEGEND

Closed Depression – R053CY019SD

HCG – Heavy continuous grazing
 IN – Invasion
 LTPG – Long-term prescribed grazing
 PB – Prescribed burning
 PC – Precipitation cycles
 PG – Prescribed grazing

Figure 9. State-And-Transition model

Code	Process
T1A	Heavy continuous grazing, invasion
T1B	Heavy continuous grazing, invasion
1.1A	Heavy continuous grazing
1.2A	Prescribed grazing, prescribed burning
2.1A	Heavy continuous grazing
2.2A	Prescribed grazing
T2A	Long-term prescribed grazing

Figure 10. Matrix

State 1
Reference State

This state represents the natural range of variability that dominated the dynamics of this ecological site (ES). This state is dominated by cool-season grasses, with warm-season grasses being subdominant. In pre-European times, the primary disturbance mechanisms for this site in the reference condition included periods of below or above average precipitation, periodic fire, and herbivory by insects and large ungulates. Timing of fires and herbivory coupled with weather events dictate the dynamics that occur within the natural range of variability. Wheatgrass species can decline and a corresponding increase in foxtail barley, short warm-season grasses, and forbs will occur. Under extended periods of disturbance, the main change is a reduction in vigor and production and an increase in bare ground and forb composition. Interpretations are based primarily on the 1.1 Reference Plant Community Phase. This plant community evolved with grazing by large herbivores and occasional fire, as well as periodic flooding and drying, and can be maintained with prescribed grazing, prescribed burning, or areas receiving occasional short periods of rest or deferment. This plant community phase has two sub-phases, referred to as plant communities here. These sub-phases are mainly driven by precipitation and flooding or drying sequences.

Community 1.1 Western Wheatgrass-Common Spikerush

Community Subphase 1.1 Western Wheatgrass-Common Spikerush – Interpretations are based primarily on the 1.1 Western Wheatgrass-Common Spikerush Plant Community Subphase. Following several years of above average precipitation, the plant community stabilizes and becomes dominated with perennial grasses such as western wheatgrass and common spikerush. Other grasses and grass-likes present include Nuttall’s alkaligrass (*Puccinellia nuttalliana*), sedge (*Carex*), rush (*Juncus*), and slender wheatgrass (*Elymus trachycaulus*). The occurrence of forbs will be considerably lower, including some species such as American licorice (*Glycyrrhiza lepidota*), curlytop knotweed (*Polygonum lapathifolium*), Pennsylvania smartweed (*Polygonum pensylvanicum*), Pursh seepweed (*Suaeda calceoliformis*), and western dock (*Rumex aquaticus*). The plant community is made up of about 80 to 90 percent grasses and grass-likes, and about 10 to 20 percent forbs. Community Subphase 1.1 Common Spikerush-Foxtail Barley – Interpretations are based primarily on the 1.1 Common Spikerush-Foxtail Barley Plant Community Subphase. This plant community often occurs after a period of higher precipitation that follows an extended dry cycle. Grasses and grass-likes commonly occurring include common spikerush, sedge, rush, foxtail barley, western wheatgrass, and bluegrasses. The forbs commonly found include western dock, mint (*Mentha*), Pursh seepweed, lambsquarters (*Chenopodium album*), knotweed (*Polygonum*), evening-primrose (*Oenothera*), buttercup (*Ranunculus*), and New England aster (*Symphotrichum novae-angliae*). The plant community is made up of about 5 to 10 percent grasses, 30 to 40 percent grass-likes, and about 50 to 60 percent forbs. Transitions or pathways leading to other plant communities are as follows: • Precipitation cycles will shift this community between the 1.1 Western Wheatgrass-Common Spikerush Plant Community Subphase and the 1.1 Common Spikerush-Foxtail Barley Plant Community Subphase. After several years of above average precipitation, the plant community stabilizes and perennial grasses or western wheatgrass will dominate the site with few grass-likes and forbs; and in the instance of higher precipitation received after extended years of drought, there will be an increase in the grass-likes and forbs components.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	2068	2845	3419
Forb	174	1078	1625
Total	2242	3923	5044

Figure 12. Plant community growth curve (percent production by month). SD5307, Southern Dark Brown Glaciated Plains, cool-season dominant, warm-season subdominant.. Cool-season dominant, warm-season subdominant, lowland..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	13	20	25	18	11	5	3	0	0

Community 1.2 Western Wheatgrass-Inland Saltgrass

This plant community is the result of heavy continuous grazing. Repeated defoliation depletes stored carbohydrates resulting in weakening and eventual death of the most palatable grasses. Lack of litter and reduced plant vigor result in higher soil temperatures, poor water infiltration rates, high evapotranspiration, and increased percolation of the high water table, which increases salt concentrations on the surface. This gives inland saltgrass (*Distichlis spicata*) and other salt tolerant species a competitive advantage over less tolerant species. Inland saltgrass drastically increases and competes with western wheatgrass as the dominant species. Other grass and grass-like species present will include Nuttall's alkaligrass, plains bluegrass (*Poa arida*), common spikerush, needle Spikerush (*Eleocharis acicularis*), and other sedges (Cyperaceae) and rushes (Juncaceae). Early cool-season grasses including foxtail barley, fowl bluegrass (*Poa palustris*), and Kentucky bluegrass (*Poa Pratensis*) begin to invade. Forbs that will invade are curly dock (*Rumex crispus*) and cocklebur (*Xanthium*). Common forbs to the site include lambsquarters, Pennsylvania smartweed, curlytop knotweed, plantain (*Plantago*), and povertyweed (*Iva axillaris*). This plant community is relatively stable and well adapted to increased salinity. Plant vigor, litter, frequency, and production have decreased. The biological integrity, water, and nutrient cycles of this plant community are becoming impaired. This plant community is less productive than the 1.1 Reference Plant Community Phase.

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1569	1973	2102
Forb	112	493	1037
Total	1681	2466	3139

Figure 14. Plant community growth curve (percent production by month). SD5303, Southern Dark Brown Glaciated Plains, cool-season/warm-season codominant.. Cool-season, warm-season codominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	10	20	28	21	10	5	3	0	0

Pathway 1.1A Community 1.1 to 1.2

Heavy continuous grazing (grazing at full to heavy levels for extended portions of the growing season without adequate recovery periods) will shift this community to the 1.2 Western Wheatgrass-Inland Saltgrass Plant Community.

Pathway 1.2A Community 1.2 to 1.1

Prescribed grazing or prescribed burning separately or the combination of both together occurring at relatively frequent intervals (3 to 5 years) and a return to normal disturbance regime levels and frequencies or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the 1.1 Reference Plant Community Phase.

State 2 Native/Invaded State

This state represents the range of variability that exists with reduced vigor and production of the dominant climax species as a result of grazing-induced disturbance and the introduction of nonnative species. This state is dominated by cool-season grasses. It can be found on areas that are impacted by extended periods of heavy continuous grazing. Grazing tolerant species become dominant, and non-native species are present.

Community 2.1 Foxtail Barley-Inland Saltgrass

This plant community developed with heavy continuous grazing where adequate recovery periods between grazing events were not allowed. Patches of inland saltgrass sod are typical and foxtail barley and fowl bluegrass are well distributed throughout the community. Nuttall's alkaligrass and western wheatgrass have been greatly reduced in production and vigor, but may persist in remnant amounts. This plant community is resistant to change due to the grazing tolerance of inland saltgrass and increased surface salts. A significant amount of production and diversity has been lost when compared to the 1.1 Reference Plant Community Phase. Loss of key cool-season grasses and increased bare ground have negatively impacted energy flow and nutrient cycling. Water infiltration is reduced significantly due to the massive shallow root system "root pan," characteristic of inland saltgrass and increased bare ground. It will take a long time to bring this plant community back to the Reference State (State 1) with management alone. Renovation (mechanical or chemical inputs separately or the combination of both together) is typically not effective due to high salt content of the soil and saltgrass persistence.

Table 7. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1042	1429	1743
Forb	78	252	499
Total	1120	1681	2242

Figure 16. Plant community growth curve (percent production by month). SD5303, Southern Dark Brown Glaciated Plains, cool-season/warm-season codominant.. Cool-season, warm-season codominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	10	20	28	21	10	5	3	0	0

Community 2.2 Curly Dock-Foxtail Barley

This plant community can be reached with heavy continuous grazing coupled with compaction due to grazing when the soil is saturated. This plant community can also result from long-term ponding and occasional subsequent drying as when this site is developed for a water source. The dominant vegetation includes pioneer annual grasses, forbs, invaders, and early successional biennial and perennial species. Grasses may include foxtail barley which may become dominant along with fowl bluegrass, Nuttall's alkaligrass, and western wheatgrass. The dominant forbs include curly dock, curlycup gumweed (*Grindelia squarrosa*), kochia (*Bassia scoparia*), cocklebur, and other early successional salt tolerant species. The community is susceptible to nonnative species due to severe soil disturbances and relatively high percent of bare ground. This plant community is resistant to change, as long as soil disturbance or severe vegetation defoliation persists, thus holding back secondary plant succession. Secondary succession is highly variable, depending upon availability and diversity of a viable seed bank of higher successional species within the existing plant community and neighboring plant communities.

Figure 17. Plant community growth curve (percent production by month). SD5302, Southern Dark Brown Glaciated Plains, cool-season dominant, warm-season subdominant.. Cool-season dominant, warm-season subdominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	10	23	34	15	6	5	4	0	0

Pathway 2.1A Community 2.1 to 2.2

Heavy continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites will shift this community to the 2.2 Curly Dock-Foxtail Barley Plant Community Phase.

Pathway 2.2A Community 2.2 to 2.1

Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the 2.1 Foxtail Barley-Inland Saltgrass Plant Community Phase.

Conservation practices

Prescribed Grazing

Transition T1A State 1 to 2

Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season, and often at the same time of year each year, typically beginning early in the season) or invasion of non-native plant species separately or the combination of both together will convert the 1.1 Common Spikerush-Foxtail Barley Plant Community Subphase within the Reference State (State 1) to the 2.2 Curly Dock-Foxtail Barley Plant Community Phase within the Native/Invaded State (State 2).

Restoration pathway T2A State 2 to 1

Long-term prescribed grazing (moderate stocking levels coupled with adequate recovery periods, or other grazing systems such as high-density, low-frequency intended to treat specific species dominance, or periodic light to moderate stocking levels possibly including periodic rest) may lead the Native/Invaded State (State 2) over a threshold to the Reference State (State 1).

Conservation practices

Prescribed Grazing

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Wheatgrass			785–3335	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	785–3335	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0–392	–
2	Cool-Season Bunchgrasses			196–1569	
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	78–1373	–
	Nuttall's alkaligrass	PUNU2	<i>Puccinellia nuttalliana</i>	78–392	–
3	Short Warm-Season Grasses			39–392	
	saltgrass	DISP	<i>Distichlis spicata</i>	39–392	–
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0–196	–
4	Other Native Grasses			78–392	
	fowl bluegrass	POPA2	<i>Poa palustris</i>	39–275	–
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	0–196	–
	plains bluegrass	POAR3	<i>Poa arida</i>	39–196	–
5	Grass-likes			196–1765	

	common spikerush	ELPA3	<i>Eleocharis palustris</i>	196–1569	–
	needle spikerush	ELAC	<i>Eleocharis acicularis</i>	39–588	–
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–392	–
	sedge	CAREX	<i>Carex</i>	78–392	–
	rush	JUNCU	<i>Juncus</i>	0–314	–
Forb					
6	Forbs			196–1961	
	Forb, native	2FN	<i>Forb, native</i>	0–785	–
	curlytop knotweed	POLA4	<i>Polygonum lapathifolium</i>	0–588	–
	Pennsylvania smartweed	POPE2	<i>Polygonum pensylvanicum</i>	0–588	–
	pale dock	RUAL4	<i>Rumex altissimus</i>	0–588	–
	western dock	RUAQ	<i>Rumex aquaticus</i>	0–588	–
	New England aster	SYNO2	<i>Symphotrichum novae-angliae</i>	0–392	–
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0–392	–
	lambsquarters	CHAL7	<i>Chenopodium album</i>	0–392	–
	bushy knotweed	PORA3	<i>Polygonum ramosissimum</i>	0–314	–
	cinquefoil	POTEN	<i>Potentilla</i>	0–196	–
	golden tickseed	COTI3	<i>Coreopsis tinctoria</i>	0–196	–
	evening primrose	OENOT	<i>Oenothera</i>	0–196	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0–196	–
	povertyweed	IVAX	<i>Iva axillaris</i>	0–196	–
	mint	MENTH	<i>Mentha</i>	0–196	–
	plantain	PLANT	<i>Plantago</i>	0–196	–
	Pursh seepweed	SUCA2	<i>Suaeda calceoliformis</i>	0–196	–
	bluebells	MERTE	<i>Mertensia</i>	0–118	–
	mealy goosefoot	CHIN2	<i>Chenopodium incanum</i>	0–118	–
	creeping woodsorrel	OXCO	<i>Oxalis corniculata</i>	0–118	–
	smooth horsetail	EQLA	<i>Equisetum laevigatum</i>	0–118	–

Table 9. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Wheatgrass			370–986	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	370–986	–
2	Cool-Season Bunchgrasses			123–493	
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	74–370	–
	Nuttall's alkaligrass	PUNU2	<i>Puccinellia nuttalliana</i>	25–247	–
3	Short Warm-Season Grasses			247–986	
	saltgrass	DISP	<i>Distichlis spicata</i>	247–986	–
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0–74	–
4	Other Native Grasses			0–123	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	0–123	–

	plains bluegrass	POAR3	<i>Poa arida</i>	0–74	–
	fowl bluegrass	POPA2	<i>Poa palustris</i>	0–74	–
5	Grass-likes			123–616	
	common spikerush	ELPA3	<i>Eleocharis palustris</i>	49–370	–
	sedge	CAREX	<i>Carex</i>	0–197	–
	needle spikerush	ELAC	<i>Eleocharis acicularis</i>	0–123	–
	rush	JUNCU	<i>Juncus</i>	0–123	–
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–123	–
6	Non-Native Grasses			25–247	
	brome	BROMU	<i>Bromus</i>	25–123	–
Forb					
7	Forbs			123–863	
	Forb, introduced	2FI	<i>Forb, introduced</i>	0–247	–
	Pennsylvania smartweed	POPE2	<i>Polygonum pennsylvanicum</i>	0–247	–
	western dock	RUAQ	<i>Rumex aquaticus</i>	0–197	–
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0–173	–
	curlytop knotweed	POLA4	<i>Polygonum lapathifolium</i>	0–173	–
	cocklebur	XANTH2	<i>Xanthium</i>	0–123	–
	Forb, native	2FN	<i>Forb, native</i>	0–123	–
	povertyweed	IVAX	<i>Iva axillaris</i>	0–123	–
	curly dock	RUCR	<i>Rumex crispus</i>	0–123	–
	Pursh seepweed	SUCA2	<i>Suaeda calceoliformis</i>	0–123	–
	bushy knotweed	PORA3	<i>Polygonum ramosissimum</i>	0–123	–
	plantain	PLANT	<i>Plantago</i>	0–123	–
	creeping woodsorrel	OXCO	<i>Oxalis corniculata</i>	0–99	–
	evening primrose	OENOT	<i>Oenothera</i>	0–74	–
	smooth horsetail	EQLA	<i>Equisetum laevigatum</i>	0–74	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0–74	–
	curlycup gumweed	GRSQ	<i>Grindelia squarrosa</i>	0–74	–
	lambsquarters	CHAL7	<i>Chenopodium album</i>	0–74	–
	pale dock	RUAL4	<i>Rumex altissimus</i>	0–74	–
	mealy goosefoot	CHIN2	<i>Chenopodium incanum</i>	0–49	–
	New England aster	SYNO2	<i>Symphotrichum novae-angliae</i>	0–49	–
	cinquefoil	POTEN	<i>Potentilla</i>	0–25	–

Table 10. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Wheatgrass			0–84	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–84	–
2	Cool-Season Bunchgrasses			336–841	
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	336–841	–
	Nuttall's alkaligrass	PUNU2	<i>Puccinellia nuttalliana</i>	0–50	–
3	Short Warm-Season Grasses			168–673	
	saltgrass	DISP	<i>Distichlis spicata</i>	168–673	–
4	Other Native Grasses			0–84	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	0–84	–
5	Grass-likes			84–336	
	common spikerush	ELPA3	<i>Eleocharis palustris</i>	34–252	–
	sedge	CAREX	<i>Carex</i>	0–84	–
	needle spikerush	ELAC	<i>Eleocharis acicularis</i>	0–84	–
	rush	JUNCU	<i>Juncus</i>	0–50	–
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–50	–
6	Non-Native Grasses			17–135	
	brome	BROMU	<i>Bromus</i>	17–84	–
	bluegrass	POA	<i>Poa</i>	0–84	–
Forb					
7	Forbs			84–420	
	curly dock	RUCR	<i>Rumex crispus</i>	0–202	–
	Forb, introduced	2FI	<i>Forb, introduced</i>	0–168	–
	lambsquarters	CHAL7	<i>Chenopodium album</i>	0–135	–
	cocklebur	XANTH2	<i>Xanthium</i>	0–118	–
	curlycup gumweed	GRSQ	<i>Grindelia squarrosa</i>	0–101	–
	Forb, native	2FN	<i>Forb, native</i>	0–84	–
	plantain	PLANT	<i>Plantago</i>	0–84	–
	smooth horsetail	EQLA	<i>Equisetum laevigatum</i>	0–67	–
	creeping woodsorrel	OXCO	<i>Oxalis corniculata</i>	0–50	–
	povertyweed	IVAX	<i>Iva axillaris</i>	0–50	–
	Pennsylvania smartweed	POPE2	<i>Polygonum pennsylvanicum</i>	0–50	–
	bushy knotweed	PORA3	<i>Polygonum ramosissimum</i>	0–50	–
	western dock	RUAQ	<i>Rumex aquaticus</i>	0–34	–
	evening primrose	OENOT	<i>Oenothera</i>	0–34	–
	mealy goosefoot	CHIN2	<i>Chenopodium incanum</i>	0–34	–
	Indianhemp	APCA	<i>Apocynum cannabinum</i>	0–17	–
	Pursh seepweed	SUCA2	<i>Suaeda calceoliformis</i>	0–17	–

Animal community

Animal Community – Grazing Interpretations

The following table lists annual, suggested initial stocking rates with average growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of conservation planning. Often, the current plant composition does not entirely match any particular plant community (as described in this ES description). Because of this, a resource inventory is necessary to document plant composition and production. More accurate carrying capacity estimates should eventually be calculated using the following stocking rate information along with animal preference data and actual stocking records, particularly when grazers other than cattle are involved. With consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity.

Western Wheatgrass – Grass-likes/ Forbs (1.1)
Average Annual Production (lbs./acre, air-dry): 3,500
Stocking Rate* (AUM/acre): 0.96

Western Wheatgrass/Inland Saltgrass (1.2)
Average Annual Production (lbs./acre, air-dry): 2,200
Stocking Rate* (AUM/acre): 0.60

Foxtail Barley/Inland Saltgrass (2.1)
Average Annual Production (lbs./acre, air-dry): 1,500
Stocking Rate* (AUM/acre): 0.41

Introduced Forbs/Foxtail Barley (2.2)
Average Annual Production (lbs./acre, air-dry): 800
Stocking Rate* (AUM/acre): 0.22

*Based on 912 lbs./acre (air-dry weight) per Animal Unit Month (AUM), and on 25 percent harvest efficiency (refer to United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS) National Range and Pasture Handbook).

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage. During the dormant period, the forage for livestock will likely be lacking protein to meet livestock requirements and added protein will allow ruminants to better utilize the energy stored in grazed plant materials. A forage quality test (either directly or through fecal sampling) should be used to determine the level of supplementation needed.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group D. Infiltration varies from very slow to slow, and runoff potential for this site varies from high to very high depending on soil hydrologic group, slope, and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Dominance by blue grama, buffalograss, bluegrass, and/or smooth brome grass will result in reduced infiltration and increased runoff. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

Recreational uses

This site provides hunting, hiking, photography, bird watching, and other opportunities. The wide varieties of plants that bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are typically present on this site.

Other products

Seed harvest of native plant species can provide additional income on this site.

Other information

- SD119 Sully County, SD did not use the (Ho) Hoven silt loam, 0 to 1 percent slopes (national symbol 2tlcc) as used in the adjoining SD107 Potter County, SD.
- SD069 Hyde County, SD did not use the (Nn) Nishon silt loam (national symbol cxlm) (R53BY003ND ESD) as used in the adjoining SD049 Faulk County. SD049 Faulk County, SD (Nn) Nishon silt loam (national symbol cxlm) (R53BY003ND ESD) will need to be split correlated to match SD069 Hyde County, SD.
- SD069 Hyde County, SD did not use the (WoB) Williams-Bowbells-Nishon complex, 1 to 6 percent slopes (national symbol cxm6) (R53BY003ND ESD) as used in the adjoining SD049 Faulk County. SD049 Faulk County, SD (WoB) Williams-Bowbells-Nishon complex, 1 to 6 percent slopes (national symbol cxm6) (R53BY003ND ESD) will need to be split correlated to match SD069 Hyde County, SD.
- SD059 Hand County, SD did not use the (GsA) Glenham-Prosper-Hoven complex, 0 to 4 percent slopes (national symbol cw4y) as used in the adjoining SD069 Hyde County, SD.
- SD059 Hand County, SD did not use the (Ma) Macken silty clay loam, 0 to 1 percent slopes (national symbol 2wkqj) as used in the adjoining SD069 Hyde County, SD.
- SD073 Jerauld County, SD did not use the (Pt) Plankinton-Prosper complex (national symbol cx89) (R55CY0019D ESD) as used in the adjoining SD003 Aurora County. SD003 Aurora County, SD (Pt) Plankinton-Prosper complex (national symbol cx89) (R55CY0019D ESD) will need to be split correlated to match SD073 Jerauld County, SD.
- SD073 Jerauld County, SD did not use the (Hw) Hoven-Plankinton silt loams (national symbol cx83) (R55CY0019D ESD) as used in the adjoining SD003 Aurora County. SD003 Aurora County, SD (Hw) Hoven-Plankinton silt loams (national symbol cx83) (R55CY0019D ESD) will need to be split correlated to match SD073 Jerauld County, SD.
- Reference and alternative states within the state and transition model are may not be fully documented and may require additional field sampling for refinement.

Inventory data references

There is no NRCS clipping data and other inventory currently available for this site. Information presented here has been derived using field observations from range-trained personnel. Those involved in developing this site include: Stan Boltz, Range Management Specialist (RMS), NRCS; Shane Deranleau, RMS, NRCS; and Mitch Faulkner, RMS, NRCS, and Bruce Kunze, Soil Scientist, NRCS.

Other references

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Coterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Gilbert, M. C., Whited, P. M., Clairain Jr, E. J., & Smith, R. D. (2006). A Regional Guidebook for Applying the Hydrogeomorphic Approach to Assessing Wetland Functions of Prairie Potholes. Washington DC.

Samson, F. B., & Knopf, F. L. (1996). Prairie Conservation Preserving North America's Most Endangered Ecosystem. Washington D.C.: Island Press.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions. Available online. Accessed March 2018.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2003. National Range and Pasture Handbook, Revision 1. Grazing Lands Technology Institute. 214 pps.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. 672pps.

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://soils.usda.gov/technical/nasis/>)

USDA, NRCS. 2018. The PLANTS Database (<http://plants.usda.gov>, 27 March 2018). National Plant Data Team,

Greensboro, NC 27401-4901 USA.

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>. (Accessed 1 March 2018).

Contributors

Stan Boltz

Approval

Suzanne Mayne-Kinney, 1/22/2024

Acknowledgments

Contact for Lead Authors: Natural Resources Conservation Service (USDA-NRCS), Redfield Soil Survey Office Redfield, SD; Lance Howe (Lance.Howe@usda.gov), Soil Survey Office Leader, USDA-NRCS, Redfield, SD; and Steve Winter (Steven.Winter@usda.gov), Soil Scientist, USDA-NRCS, Redfield, SD

Additional Information Acknowledgment: Jason Hermann (Jason.Hermann@usda.gov), Area Rangeland Management Specialist, USDA-NRCS, Redfield, SD.

This Provisional Ecological Site concept has passed both Quality Control and Quality Assurance processes. Quality Assurance was approved by David Kraft, NRCS Regional Ecologist as of 11/12/2020.

Non-discrimination Statement

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, available online and at any USDA office, or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632- 9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Boltz, Mitch Faulkner, Shane Deranleau
Contact for lead author	Stan Boltz, stanley.boltz@sd.usda.gov, 605-352-1236
Date	03/15/2011
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills should not be present.

2. **Presence of water flow patterns:** Barely observable or not present.

3. **Number and height of erosional pedestals or terracettes:** None.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Normally bare ground is less than 5 percent and patches less than two inches in diameter. Following well-above average or well-below average precipitation periods, bare ground can be very high for brief periods of time.

5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None present.

7. **Amount of litter movement (describe size and distance expected to travel):** Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil aggregate stability normally a 4 to 6 rating. Typically high root content and organic matter in the soil surface. Soil surface is very resistant to erosion.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Hoven soils have an E-horizon (leached) at the surface which is platy structure, and typically will not have mollic colors at the surface. Macken and Plankinton soils have variable surface structure, either platy or subangular blocky, but parting to granular, and mollic (higher organic matter) colors of A-horizon down to about 2 to 4 inches.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Healthy, deep-rooted native grass and grass-like species enhance infiltration

and reduce runoff.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer should be present. Surface horizon of Hoven soils may be platy and appear to be compacted and should not be confused with compaction. At 2 to 4 inches, Plankinton and Macken soils may have platy structure, and this should not be confused with a compaction layer.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Drier precipitation cycles: Wheatgrasses (mid, cool-season rhizomatous) >> mid, cool-season bunchgrasses >
- Wetter precipitation cycles: Grass-like species = forbs >
- Sub-dominant: Drier precipitation cycles: Short, warm-season grasses >
- Wetter precipitation cycles: Wheatgrasses (mid, cool-season rhizomatous) > short, warm-season grasses >
- Other: Drier precipitation cycles: Forbs > grass-like species
- Wetter precipitation cycles: Mid, cool-season bunchgrasses
- Additional: Other grasses in other functional groups occur in minor amounts.
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little to no evidence of decadence or mortality.
-
14. **Average percent litter cover (%) and depth (in):** 55-80 percent plant litter cover, roughly 0.5 to 1 inch in depth. Litter cover is in contact with the soil surface.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 3,500 pounds/acre (air-dry basis)
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Refer to State and local Noxious Weed List.
-
17. **Perennial plant reproductive capability:** Perennial grass and grass-like species have vigorous rhizomes and/or tillers.
-