

Ecological site R055BY056ND **Clayey**

Accessed: 05/17/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Classification relationships

Level IV Ecoregions of the Conterminous United States: 42a – Missouri Coteau; 42b – Collapsed Glacial Outwash; 42c – Missouri Coteau Slope; 42d – Northern Missouri Coteau; 42f – Southern Missouri Coteau Slope; 42g – Ponca Plains; and 42h – Southern River Breaks.

Associated sites

R055BY057ND	Claypan This site (Claypan) typically occurs slightly lower on the landscape. It has a root-restrictive claypan layer. Characteristic vegetation includes mostly cool-season grasses, with warm-season grasses being subdominant, and native forbs. Western Wheatgrass (<i>Pascopyrum smithii</i>) and Green Needlegrass (<i>Nassella viridula</i>) are in the reference state.
R055BY058ND	Limy Subirrigated This site (Limy Subirrigated) occurs lower on the landscape. It is highly calcareous in the upper part of the subsoil and has redoximorphic features at a depth of 18 to 30 inches. Characteristic vegetation includes tall and mid-statured, warm-season grasses and native forbs. Little Bluestem (<i>Schizachyrium Nees</i>) and Big Bluestem (<i>Adropodon gerardii</i>) are in the reference state.
R055BY059ND	Loamy Overflow

R055BY064ND	Loamy This site (Loamy) typically occurs somewhat higher on the landscape. It has <35% clay in subsoil layer. Characteristic vegetation includes predominantly cool-season grass, some warm-season grasses, and native forbs. Western Wheatgrass (<i>Pascopyrum smithii</i>) and Green Needlegrass (<i>Nassella viridula</i>) are in the reference state.
-------------	---

Similar sites

R055BY057ND	Claypan This site (Claypan) typically occurs slightly lower on the landscape. It has a root-restrictive claypan layer. Characteristic vegetation includes mostly cool-season grasses, with warm-season grasses being subdominant, and native forbs. Western Wheatgrass (<i>Pascopyrum smithii</i>) and Green Needlegrass (<i>Nassella viridula</i>) are in the reference state; but it has less green needlegrass; more blue grama; and lower production than Clayey.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Nassella viridula</i> (2) <i>Pascopyrum smithii</i>

Physiographic features

This site typically occurs on glaciated uplands – ground moraines and lake plains and colluvium from weathered shale and shaly till. This site is typically on linear back slopes, concave foot slopes, or on flats on till plains and lake plains and on side slopes of escarpments. Slopes are less than 15 percent. On ground moraines the parent material is either fine-loamy or clayey. On lake plains the parent material is either fine-silty or clayey. On escarpments the parent material is clayey.

The following data (tables) were obtained from the National Soil Information System.

Table 2. Representative physiographic features

Landforms	(1) Lake plain (2) Till plain (3) Flood plain
Elevation	305–610 m
Slope	0–15%
Water table depth	91–203 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 55B is considered to have a continental climate – cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature are characteristic. The climate is the result of this MLRA's location in the geographic center of North America. There are few natural barriers on the northern Great Plains. The air masses move unobstructed across the plains and account for rapid changes in temperature.

Annual precipitation ranges from 16 to 21 inches per year. The normal average annual temperature is about 41.5° F. January is the coldest month with average temperatures ranging from about 2° F (Maddock, ND) to about 11° F (Mellette, SD). July is the warmest month with temperatures averaging from about 67° F (Maddock, ND) to about 73° F (Redfield 2 NE, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 64° F. This large annual range attests to the continental nature of this MLRA's climate. Winds average about 11 miles per hour annually, ranging from about 13 miles per hour during the spring to about 10 miles per hour during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour.

Growth of native cool-season plants begins in late March and continues to early to mid July. Native warm-season plants begin growth in mid May and continue to the end of August. Green up of cool-season plants can occur in September and October when adequate soil moisture is present.

Table 3. Representative climatic features

Frost-free period (average)	140 days
Freeze-free period (average)	161 days
Precipitation total (average)	533 mm

Influencing water features

No significant water features influence this site.

Soil features

The common features of soils in this site are the silty clay loam to clay textured subsoils and slopes of 0 to 25 percent. The loam to silty clay surface layer is 5 to 18 inches thick. The soils have a moderately slow to very slow infiltration rate. The soils in this site are well to moderately well drained and formed in glaciolacustrine sediments and clayey till. When dry these soils crack. When the soils are wet, surface compaction can occur with heavy traffic. This site typically should show slight to no evidence of rills, wind scoured areas or pedestalled plants. Water flow paths should not be present, and the soil surface is stable and intact. Sub-surface soil layers are non-restrictive to water movement and root penetration. These soils are susceptible to water and wind erosion. The hazard of water erosion increases on slopes greater than about 5 percent. Access Web Soil Survey (<http://websoilsurvey.nrcs.usda.gov/app/>) for specific local soils information.

Table 4. Representative soil features

Surface texture	(1) Silty clay loam (2) Silt loam (3) Loam
Family particle size	(1) Clayey
Drainage class	Moderately well drained to well drained
Permeability class	Very slow to moderately slow
Soil depth	203 cm
Surface fragment cover <=3"	0–3%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	15.24–30.48 cm
Calcium carbonate equivalent (0-101.6cm)	0–20%
Electrical conductivity (0-101.6cm)	0–8 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–15
Soil reaction (1:1 water) (0-101.6cm)	5.6–9
Subsurface fragment volume <=3" (Depth not specified)	0–10%
Subsurface fragment volume >3" (Depth not specified)	0–5%


Ecological dynamics

The site developed under Northern Great Plains climatic conditions, and included natural influence of large herding herbivores and occasional fire. Changes will occur in the plant communities due to weather fluctuations and/or management actions. Under adverse impacts, a slow decline in vegetative vigor and composition will occur. Under favorable conditions the site has the potential to resemble the Reference State. Interpretations for this site are based on the Green Needlegrass/Western Wheatgrass Plant Community Phase (1.1). The Reference State and the interpretive plant community have been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been considered. Community phases, community pathways, states, transitions, thresholds and restoration pathways have been determined through similar studies and experience. The grazing and fire interaction especially when coupled with drought events, set up the dynamics discussed and displayed in the following state and transition diagram and descriptions.

This ecological site has been grazed by domestic livestock since introduced into the area. The introduction of domestic livestock and the use of fencing and reliable water sources have radically changed the disturbance regime of this site. Heavy continuous grazing and/or continuous seasonal (spring) grazing, without adequate recovery periods following each grazing occurrence causes this site to depart from the reference plant community. Blue grama and Kentucky bluegrass if present, will begin to increase. Western wheatgrass will increase initially and then begin to decrease. Green needlegrass will decrease in frequency and production. In time, heavy continuous grazing will likely cause upland sedges and blue grama and/or Kentucky bluegrass if present to dominate and pioneer perennials and annuals to increase. The resulting plant community is relatively stable and competitive advantage prevents other species from establishing. Extended periods of non-use and/or lack of fire will result in a plant community having high litter levels, which favors an increase in Kentucky bluegrass and/or smooth brome grass.

Following the state and transition diagram are narratives for each of the described states and community phases. These may not represent every possibility, but they are the most prevalent and repeatable states/community phases. The plant composition tables shown below have been developed from the best available knowledge at the time of this revision. As more data are collected, some of these community phases and/or states may be revised or removed, and new ones may be added. The main purpose for including the descriptions here is to capture the current knowledge and experience at the time of this revision.

State and transition model



Refer to narratives in the Plant Community Section for detailed descriptions of these transitions/pathways. **1.1a** – Spring fire followed by intensive grazing; **1.2a** – Return to normal fire & grazing frequencies; **T1** – Introduction of non-native species; **2.1a** – Heavy continuous grazing (normally late season or season-long, inadequate recovery periods); **2.1b** – Heavy season-long grazing or lack of grazing and fire; **2.2a** – Prescribed grazing (including adequate recovery periods); **2.2b** – No fire, non-use; **2.3a** – Prescribed burning, coupled with prescribed grazing; **T2** – Long-term rest from grazing and fire; **R3a** – Range seeding with native species with management to control invasive species; **3.1a** – No fire, non use.

State 1 Reference

This state represents the natural range of variability that dominated the dynamics of this ecological site. This state was dominated by cool-season grasses, with warm-season grasses being subdominant. In pre-European times, the primary disturbance mechanisms for this site in the reference condition included periods of below and/or above average precipitation, periodic fire, and herbivory by insects and large ungulates. Timing of fires and herbivory coupled with weather events dictated the dynamics that occurred within the natural range of variability. Cool-season and taller warm-season grasses would have declined and a corresponding increase in short, warm-season grasses would have occurred. Today, a similar state (State 2) can be found on areas that are properly managed with grazing and/or prescribed burning, and sometimes on areas receiving occasional short periods of rest.

Community 1.1 Green Needlegrass/Western Wheatgrass

This community phase was the most dominant both temporally and spatially. The prevailing climate and weather patterns favored the development of this community phase dominated by cool-season grasses such as green needlegrass and western wheatgrass. There are also other needlegrasses and wheatgrasses present as well as various amounts of warm-season grasses such as big bluestem, blue grama and sideoats grama. A variety of leguminous and non-leguminous perennial forbs are present but only in slight amounts. This is the reference plant community phase and is described in the “Plant Community Composition and Group Annual Production” portion of this ecological site description. This is a naturally nitrogen deficient plant community.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1849	2608	3351
Forb	140	219	308
Shrub/Vine	28	87	151
Total	2017	2914	3810

Figure 5. Plant community growth curve (percent production by month).
ND5502, Central Black Glaciated Plains, cool-season dominant, warm-season sub-dominant.. Cool-season dominant, warm-season sub-dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	7	23	42	15	5	4	1	0	0

Community 1.2 Western Wheatgrass/Blue Grama/Sedge/Green Needlegrass

This plant community was a result of concentrated grazing following a spring fire, from heavy continuous grazing or from over utilization during extended drought periods. The potential plant community was made up of approximately 85 percent grasses and grass-like species, 10 percent forbs, and 5 percent shrubs. Dominant grass and grass-like species included western wheatgrass, blue grama, sedge, buffalograss, and green needlegrass. Grasses of secondary importance included sideoats grama, needleandthread, and porcupine grass. Forbs commonly found in this plant community included cudweed sagewort, heath aster, and western yarrow. This plant community had similar plant composition to the 2.2 Western Wheatgrass/Kentucky Bluegrass/Blue Grama/Sedge Plant Community Phase. The main difference is that this plant community phase did not have the presence of non-native invasive species such as Kentucky bluegrass and smooth brome grass. When compared to the Green Needlegrass/Western Wheatgrass Plant Community Phase (1.1), blue grama and sedge increased. Green needlegrass decreased, and production was also reduced. This plant community was moderately resistant to change. The herbaceous species present were well adapted to grazing; however, species composition could be altered through long-term overgrazing. If the herbaceous component was intact, it tended to be resilient if the disturbance was not long-term. The increase of shorter-statured, more compact rooted species would have resulted in somewhat higher runoff and decreased infiltration. This would have caused the site to become drier. These species also would have been more competitive.

Figure 6. Plant community growth curve (percent production by month).
ND5502, Central Black Glaciated Plains, cool-season dominant, warm-season sub-dominant.. Cool-season dominant, warm-season sub-dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	7	23	42	15	5	4	1	0	0

Pathway 1.1a Community 1.1 to 1.2

This pathway occurs as a result of spring fire followed by heavy grazing, or a combination of moderate to heavy

grazing coupled with prolonged periods of below-average precipitation. The dominant cool-season grasses such as green needlegrass and western wheatgrass will decrease, and shorter statured species such as blue grama and sedge will increase. This pathway would have led to the 1.2 Western Wheatgrass/Blue Grama/Sedge/Green Needlegrass Plant Community Phase.

Pathway 1.2a
Community 1.2 to 1.1

This pathway occurred when grazing, precipitation, and/or fire returned to normal disturbance regime levels and frequencies or periodic light to moderate grazing possibly including periodic rest occurred. This would have led to the 1.1 Green Needlegrass/Western Wheatgrass Plant Community Phase.

State 2
Native/Invaded Grass

This state represents the more common range of variability that exists with higher levels of grazing management but in the absence of periodic fire due to fire suppression. This state is dominated by cool-season grasses with warm-season grasses being subdominant. It can be found on areas that are properly managed with grazing and/or prescribed burning, and sometimes on areas receiving occasional short periods of rest. Taller cool-season species can decline and a corresponding increase in short statured grass will occur.

Community 2.1
Green Needlegrass/Western Wheatgrass/Kentucky Bluegrass

This plant community phase is similar to the 1.1 Green Needlegrass/Western Wheatgrass Plant Community Phase, but it also contains minor amounts of non-native invasive grass species such as Kentucky bluegrass and smooth brome grass (up to about 10 percent by air-dry weight). The potential vegetation is about 85 percent grasses or grass-like plants, 10 percent forbs, and 5 percent shrubs. The community is dominated by cool-season grasses, with warm-season grasses being subdominant. The major grasses include western wheatgrass, green needlegrass, slender wheatgrass, big bluestem, and sideoats grama. Other grass or grass-like species include needleandthread, porcupine grass, blue grama, Kentucky bluegrass, buffalograss, prairie dropseed, and sedge. This plant community is resilient and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community in regards to site/soil stability, watershed function, and biologic integrity.

Figure 7. Plant community growth curve (percent production by month).
ND5502, Central Black Glaciated Plains, cool-season dominant, warm-season sub-dominant.. Cool-season dominant, warm-season sub-dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	7	23	42	15	5	4	1	0	0

Community 2.2
Western Wheatgrass/Kentucky Bluegrass/Blue Grama/Sedge

This plant community is a result of heavy continuous grazing, continuous season-long grazing or from over utilization during extended drought periods. The potential plant community is made up of approximately 80 percent grasses and grass-like species, 15 percent forbs, and 5 percent shrubs. Dominant grass and grass-like species include western wheatgrass, blue grama, sedge, and Kentucky bluegrass. Grasses of secondary importance include green needlegrass, needleandthread, porcupine grass, buffalograss, smooth brome grass, and sedge. Forbs commonly found in this plant community include cudweed sagewort, heath aster, prairie coneflower, and western yarrow. When compared to the Green Needlegrass/Western Wheatgrass Plant Community Phase (1.1), blue grama and sedge have increased, and Kentucky bluegrass has invaded. Green needlegrass and production of mid and tall grasses has also been reduced. This plant community is moderately resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term. The increase of shorter-statured species with a dense, shallower rooted species will result in somewhat higher runoff and decreased infiltration. This will cause the site to become drier. These species will also become more

competitive.

Figure 8. Plant community growth curve (percent production by month).
ND5502, Central Black Glaciated Plains, cool-season dominant, warm-season sub-dominant.. Cool-season dominant, warm-season sub-dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	7	23	42	15	5	4	1	0	0

Community 2.3

Western Wheatgrass/Kentucky Bluegrass/Green Needlegrass

This plant community is a result of continuous season-long grazing, typically at light levels, or prolonged periods (multiple years) of complete rest from grazing and elimination of fire. This community phase is characterized by an increase in the introduced cool-season sodgrass, Kentucky bluegrass. This community phase is the most dominant both temporally and spatially. Kentucky bluegrass has become nearly co-dominant with western wheatgrass and green needlegrass. Warm season grasses are present but minor and tap rooted perennial forbs have decreased. Production and infiltration both decrease and this community phase is at risk of transitioning across a state threshold. With natural or management actions that decrease the composition of the cool-season bunchgrasses and increase the composition of Kentucky bluegrass, transition T2 will be initiated.

Figure 9. Plant community growth curve (percent production by month).
ND5501, Central Black Glaciated Plains, cool-season dominant. Cool-season dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	8	24	45	10	3	5	2	0	0

Pathway 2.1a

Community 2.1 to 2.2

This pathway occurs as a result of heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season, and often at the same time each year), or continuous season-long grazing, or a combination of disturbances such as extended periods of below average precipitation coupled with periodic heavy grazing. This pathway will lead to the 2.2 Western Wheatgrass/Kentucky Bluegrass/Blue Grama/Sedge Plant Community Phase.

Pathway 2.1b

Community 2.1 to 2.3

Prolonged periods (multiple years) of continuous season-long grazing, or complete rest from grazing or grazing at very light levels coupled with elimination of fire results in increased litter levels and decreased vigor of less shade tolerant species. These factors favor cool-season species, and lead to the 2.3 Western Wheatgrass/Kentucky Bluegrass/Green Needlegrass Plant Community Phase. When continuous or light grazing is involved, this community will often occur in a patchy mosaic pattern, often referred to as patch grazing.

Pathway 2.2a

Community 2.2 to 2.1

The implementation of prescribed grazing, including adequate recovery periods between grazing events and season of use change, will initiate this pathway by shifting the competitive advantage away from the short statured grasses to the taller cool-season grasses.

Pathway 2.2b

Community 2.2 to 2.3

Prolonged periods (multiple years) of complete rest from grazing or grazing at very light levels coupled with elimination of fire results in increased litter levels and decreased vigor of less shade tolerant species. These factors

favor cool-season species, and lead to the 2.3 Western Wheatgrass/Kentucky Bluegrass/Green Needlegrass Plant Community Phase.

Pathway 2.3a
Community 2.3 to 2.1

The implementation of prescribed burning coupled with prescribed grazing including adequate recovery periods between grazing events and season of use change will initiate this pathway by shifting the competitive advantage away from the short statured grasses to the taller native grasses.

State 3
Invaded

This state is the result of invasion and dominance of introduced species. This state is characterized by the dominance of Kentucky bluegrass and smooth brome grass, and an increasing thatch layer that effectively blocks introduction of other plants into the system. Plant litter accumulation tends to favor the more shade tolerant introduced grass species. The nutrient cycle is also impaired, and the result is typically a higher level of nitrogen which also favors the introduced species. Increasing plant litter decreases the amount of sunlight reaching plant crowns thereby shifting competitive advantage to shade tolerant introduced grass species. Studies indicate that soil biological activity is altered, and this shift apparently exploits the soil microclimate and encourages growth of the introduced grass species. Once the threshold is crossed, a change in grazing management alone cannot cause a reduction in the invasive grass dominance. Once the state is well established, even drastic events such as high intensity fires driven by high fuel loads of litter and thatch will not result in more than a very short term reduction of Kentucky bluegrass. These events may reduce the dominance of Kentucky bluegrass, but due to the large amount of rhizomes in the soil there is no opportunity for the native species to establish and dominate before Kentucky bluegrass rebounds and again dominates the system.

Community 3.1
Kentucky Bluegrass/Smooth Brome grass

This plant community phase is a result of heavy, continuous seasonal grazing or heavy, continuous season-long grazing. It is characterized by a dominance of Kentucky bluegrass, smooth brome grass, sedge, and blue grama. The dominance is at times so complete that other species are difficult to find on the site. A relatively thick duff layer can sometimes accumulate at or above the soil surface. Nutrient cycling is greatly reduced, and native plants have great difficulty becoming established. Infiltration is greatly reduced and runoff is high. Production will be significantly reduced when compared to the interpretive plant community. The period that palatability is high is relatively short, as Kentucky bluegrass matures rapidly. Energy capture is also reduced. Biological activity in the soil is likely reduced significantly in this phase.

Figure 10. Plant community growth curve (percent production by month).
ND5501, Central Black Glaciated Plains, cool-season dominant. Cool-season dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	8	24	45	10	3	5	2	0	0

Community 3.2
Smooth Brome grass

This plant community phase is a result of extended periods of non-use and no fire. It is characterized by a dominance of smooth brome grass and Kentucky bluegrass. The dominance is at times so complete that other species are difficult to find on the site. A thick duff layer also accumulates at or above the soil surface. Nutrient cycling is greatly reduced, and native plants have great difficulty becoming established. When dominated by smooth brome grass, infiltration is moderately reduced and runoff is moderate. Production can be equal to or higher than the interpretive plant community. However, when dominated by Kentucky bluegrass, infiltration is greatly reduced and runoff is high. Production in this case will likely be significantly less. In either case, the period that palatability is high is relatively short, as these cool-season species mature rapidly. Energy capture is also reduced.

Figure 11. Plant community growth curve (percent production by month). ND5501, Central Black Glaciated Plains, cool-season dominant. Cool-season dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	8	24	45	10	3	5	2	0	0

Pathway 3.1a Community 3.1 to 3.2

Prolonged periods (multiple years) of complete rest from grazing or grazing at very light levels coupled with elimination of fire results in increase litter levels and decreased vigor of less shade tolerant species. These factors favor cool-season species, and lead to the 3.2 Smooth Brome grass Plant Community Phase.

Transition T1 State 1 to 2

This is the transition from the native herbaceous dominated reference state to the herbaceous dominated native/invaded state. This transition occurs when propagules of non-native species such as Kentucky bluegrass and/or smooth brome grass are present and become established on the site. This occurs as natural and/or management actions (altered grazing and/or fire regime) favor an increase in cool-season sodgrasses. Chronic season-long or heavy late season grazing facilitates this transition. Complete rest from grazing and no fire events can also lead to this transition. The threshold between states is crossed when the non-natives become established on the site.

Transition T2 State 2 to 3

This represents the transition from the more native dominated Native/Invaded State to a state dominated by non-native cool-season grasses. Preliminary studies would tend to indicate this threshold may exist when Kentucky bluegrass exceeds 30% of the plant community and native grasses represent less than 40% of the plant community composition.

Restoration pathway R3a State 3 to 2

It may be possible using selected plant materials and agronomic practices to approach something very near the functioning of the Native/Invaded State (State 2). Application of chemical herbicides and the use of mechanical seeding methods using adapted varieties of the dominant native bunchgrasses are possible and can be successful. After establishment of the native bunchgrasses, management objectives must include the maintenance of those species, the associated reference function and continued treatment of the introduced sodgrasses. This restoration pathway may also be initiated with high levels of prescribed grazing management over a long period of time, coupled with prescribed burning. The success of this restoration pathway depends on the presence of a remnant population of native grass and grass-like species, or the presence of these species in nearby similar sites. This remnant population may not be readily apparent without close inspection.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Needlegrass			729–1166	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	583–1166	–
	porcupinegrass	HESP11	<i>Hesperostipa spartea</i>	29–146	–
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	0–87	–

2	Wheatgrass			291–583	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	291–583	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	58–291	–
	slender wheatgrass	ELTRS	<i>Elymus trachycaulus</i> ssp. <i>subsecundus</i>	58–291	–
3	Tall/Mid Warm-Season Grasses			58–291	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	0–291	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–291	–
	prairie dropseed	SPHE	<i>Sporobolus heterolepis</i>	0–146	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–87	–
4	Short Warm-season Grasses			58–146	
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	29–146	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	29–146	–
5	Other Native Grasses			58–146	
	Graminoid (grass or grass-like)	2GRAM	<i>Graminoid (grass or grass-like)</i>	0–87	–
	plains reedgrass	CAMO	<i>Calamagrostis montanensis</i>	29–87	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	29–87	–
6	Grass-likes			29–146	
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–146	–
	sedge	CAREX	<i>Carex</i>	29–146	–
Forb					
7	Forbs			146–291	
	Forb, native	2FN	<i>Forb, native</i>	0–146	–
	western yarrow	ACMIO	<i>Achillea millefolium</i> var. <i>occidentalis</i>	29–58	–
	dotted blazing star	LIPU	<i>Liatris punctata</i>	29–58	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	29–58	–
	false boneset	BREU	<i>Brickellia eupatorioides</i>	29–58	–
	wavyleaf thistle	CIUN	<i>Cirsium undulatum</i>	29–58	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	29–58	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	29–58	–
	American vetch	VIAM	<i>Vicia americana</i>	29–58	–
	silverleaf Indian breadroot	PEAR6	<i>Pedimelum argophyllum</i>	29–58	–
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	29–58	–
	goldenrod	SOLID	<i>Solidago</i>	0–58	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0–29	–
	sanddune wallflower	ERCAC	<i>Erysimum capitatum</i> var. <i>capitatum</i>	0–29	–
	scarlet beeblossom	GACO5	<i>Gaura coccinea</i>	0–29	–
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	0–29	–
Shrub/Vine					
8	Shrubs			29–146	
	western snowberry	SYOC	<i>Symphoricarpos occidentalis</i>	0–87	–
	head-leaf	AMCAG	<i>Amorpha canescens</i>	0–87	–

	readaptant	AMCAB	<i>Amorpha canescens</i>	29–58	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	29–58	–
	prairie rose	ROAR3	<i>Rosa arkansana</i>	0–58	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–58	–

Animal community

Animal Community – Wildlife Interpretations

Major Land Resource Area (MLRA) 55B lies within the Northern mixed-grass prairie ecosystem. Prior to European settlement, this area consisted of diverse grassland habitats interspersed with varying densities of depressional wetlands and limited woody riparian corridors. These habitats provided critical life cycle components for many of its users. Many species of grassland birds and herds of roaming bison, elk, and pronghorn were among the inhabitants. These species, as well as several small mammal and insect species, were the primary consumers linking the grassland resources to predators such as wolves, mountain lions, and grizzly bears as well as smaller carnivores such as coyotes, bobcats, foxes and raptors. In addition, a wide variety of small mammals, reptiles, amphibians and insects were adapted to this semi-arid climate.

Historically, the Northern mixed-grass prairie was a disturbance-driven ecosystem with fire, herbivory and climate functioning as the primary disturbance factors either singly or in combination. Following European settlement, widespread conversion to cropland, elimination of fire, and habitat fragmentation influenced species composition and abundance. Introduced and invasive species further impacted plant and animal communities. Bison were historically a keystone species but have been extirpated as a free-ranging herbivore. The loss of bison and fire as ecological drivers greatly influenced the character of the remaining native plant community and the habitats that they provide. Fragmentation has reduced habitat quality for area-sensitive species.

Animal Community – Grazing Interpretations

This site is well adapted to managed grazing by domestic livestock. The predominance of herbaceous plants across all plant community phases best lends these sites to grazing by cattle but other domestic grazers with differing diet preferences may also be a consideration depending upon management objectives. Often, the current plant community does not entirely match any particular plant community (as described in the ecological site description). Because of this, a resource inventory is necessary to document plant composition and production. Proper interpretation of this inventory data will permit the establishment of a safe, initial stocking rate for the type and class of animals and level of grazing management. More accurate stocking rate estimates should eventually be calculated using actual stocking rate information and monitoring data.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group C. Infiltration varies from very slow to slow, and runoff potential for this site varies from high to very high depending on soil hydrologic group, slope and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Dominance by blue grama, buffalograss, bluegrass, and/or smooth brome grass will result in reduced infiltration and increased runoff. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

Seed harvest of native plant species can provide additional income on this site.

Inventory data references

Information presented here has been derived from NRCS and other federal/state agency clipping and inventory data. Also, field knowledge of range-trained personnel was used. All descriptions were peer reviewed and/or field-tested by various private, state and federal agency specialists. Those involved in developing this site description include: Stan Boltz, NRCS Range Management Specialist; David Dewald, NRCS State Biologist; Jody Forman, NRCS Range Management Specialist; Jeff Printz, NRCS State Range Management Specialist; Kevin Sedivec, Extension Rangeland Management Specialist; Shawn Dekeyser, North Dakota State University; Rob Self, The Nature Conservancy and Lee Voigt, NRCS Range Management Specialist.

Other references

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728.
(<http://hpccsun.unl.edu>)

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224.
(<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS, Various Published Soil Surveys.

Contributors

Jeff Printz

Megan Baxter

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Printz, Stan Boltz, Lee Voigt, Jody Forman
Contact for lead author	Jeff.printz@nd.usda.gov 701-530-2080
Date	04/19/2012
Approved by	Jeff Printz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

-
2. **Presence of water flow patterns:** None.
-

3. **Number and height of erosional pedestals or terracettes:** None.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground < 5%.
-
5. **Number of gullies and erosion associated with gullies:** None.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
-
7. **Amount of litter movement (describe size and distance expected to travel):** None.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use soil series description for depth, color and structure of A horizon/surface layer.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Mid cool-season bunchgrasses >>
- Sub-dominant: Mid cool-season rhizomatous grasses >>
- Other: Tall and mid warm-season grasses = forbs > short warm-season grasses = grass-like = shrubs = short, cool-season bunchgrasses
- Additional: Due to differing root structure and distribution, Kentucky bluegrass and smooth brome grass do not fit into reference plant community F/S groups.
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Little to no evidence of mortality or decadence.

14. **Average percent litter cover (%) and depth (in):** Plant litter is in contact with soil surface.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Representative value = 2600 lbs/ac air dry with a range of 1800 to 3400 lbs/acre air dry depending up growing conditions.

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious weeds, Kentucky bluegrass, smooth brome grass

17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.
