

Ecological site R055BY069ND Very Shallow

Last updated: 4/25/2025
Accessed: 04/30/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 055B–Central Black Glaciated Plains

Central Black Glaciated Plains MLRA is an expansive and agriculturally important region consisting of more than 10,000,000 acres and including all or a portion of 27 counties in east-central and southeastern North Dakota and northeastern South Dakota.

Most of MLRA 55B is covered by till: material that was moved and redeposited by the glaciers. Pre-glaciated bedrock (shale) is exposed on the breaks to some of the valleys and incised drainageways; but what covers the bedrock is glacial sediment, known as drift. These areas have the Late Wisconsin age till plain integrated drainage system in contrast to the closed drainage of much of the till plain and moraines. Some soils, particularly along the Elm, James and Sheyenne rivers, have weathered shale beds in the substratum.

The Drift Prairie Region consists of nearly level to gently rolling glacial till plains dissected by glacial outwash channels. MLRA 55B is located within the boundaries of the Prairie Pothole Region with numerous wetlands in areas without integrated drainage systems. Seven rivers flow through parts of the MLRA. The James and Sheyenne Rivers both have their headwaters in the northern part of the MLRA. A relatively narrow, low range of hills separates these rivers creating a continental watershed divide. The James River flows generally southward through the MLRA and empties into the Missouri River beyond the MLRA border. The Sheyenne River flows to the south and to the east; it empties into the Red River of the North in MLRA 56A. Major tributaries to the James River are the Pipestem

and Elm Rivers. The Sheyenne River receives additional water from Devils Lake (during periods of high lake levels) via two outlet pumping stations. Other important rivers in the MLRA are the Goose, Maple, and Wild Rice rivers which are also tributaries to the Red River of the North. The Wild Rice River begins in northeastern South Dakota and flows northward and eastward. In Sargent County, North Dakota, major ditch construction has served to straighten this river and more quickly drain water off adjacent farmland.

Surface and subsurface (tile) drainage systems have been constructed/installed in many areas to manage excess water and/or salinity on cropland. Soils that were poorly drained prior to wide-spread drainage may now function as somewhat poorly drained or moderately well drained soils. Restoration of hydrology to the natural conditions of the reference state may not be possible.

This region is utilized mostly by farms and ranches; about 75 percent is non-irrigated cropland. Cash-grain, bean and oil production crops are the principal enterprise on many farms, but other feed grains and hay are also grown. The vegetation on the steeper slopes, very stony areas, and thinner (or sandy) soils is still native rangeland. About 1 percent of this area is forested. Most forested areas occur along rivers, particularly the Sheyenne River Valley.

Classification relationships

Level IV Ecoregions of the Conterminous United States: 46c – Glacial Lake Basins; 46d – Glacial Lake Delta; 46e – Tewaukon Dead ice Moraine; 46f – End Moraine Complex; 46i – Drift Plains; and 46j – Glacial Outwash.

Ecological site concept

The Very Shallow ecological site is located on uplands – primarily on outwash plains and on terraces and moraines dissected by rivers, streams and drainageways; it also occurs on high beach ridges along some lake shores. The soils are very shallow (<14 inches) to layers high in coarse sand and gravel (15 to 60 percent gravel). In some soils this layer contains significant amounts (>25%) of shale fragments. The thin surface layer is typically sandy loam, gravelly sandy loam, or loamy coarse sand; but gravelly loam, cobbly loam, and coarse sandy loam also occur. Where the soils have been cultivated, the surface layer may be very gravelly. Soil on this site is excessively drained. Slopes range from 0 to 35 percent. On the landscape, this site is above the Shallow Gravel, Loamy, and Subirrigated ecological sites. On moraines, the Thin Loamy site occurs on similar landscape positions; it is highly calcareous with loamy or silty textures to a depth of 40 inches or more.

To see a full copy of the ecological site description with all tables and the full version 5 rangeland health worksheet. Please use the following hyperlink:

https://efotg.sc.egov.usda.gov/references/public/ND/55B_Very_Shallow_Narrative_FINALRef_FSG.pdf

Associated sites

R055BY064ND	Loamy This site typically occurs somewhat lower on the landscape. It either does not have a sand and gravel layer or it is deeper than 20 inches to that layer. The surface and subsoil layers form a ribbon 1 to 2 inches long.
R055BY068ND	Thin Loamy This site occurs on shoulder slopes of moraines. The soil does not have a sand and gravel layer. The surface and subsoil layers form a ribbon 1 to 2 inches long. It is highly calcareous (strong or violent effervescence) within a depth of 8 inches.
R055BY065ND	Subirrigated This site occurs lower on the landscape on beach ridges. It has redoximorphic features at a depth of 18 to 30 inches. Where a highly calcareous subsoil layer occurs, it is at a depth >16 inches. All textures are included in this site.
R055BY063ND	Shallow Gravel This site occurs lower on the landscape. The depth to sand and gravel is 14 to 20 inches.

Similar sites

R055BY063ND	Shallow Gravel This site occurs lower on the landscape. The depth to sand and gravel is 14 to 20 inches.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Hesperostipa comata</i> (2) <i>Pascopyrum smithii</i>

Physiographic features

This site occurs on uplands – on glacial outwash plains, on terraces, on dissected moraines, and on high beach ridges along lake shores. It is on upper back slopes, shoulder slope and summits. Parent materials are gravelly glaciofluvial or beach deposits. Slopes range from 0 to 35 percent.

Table 2. Representative physiographic features

Landforms	(1) Moraine (2) Beach ridge (3) Outwash plain
Runoff class	Negligible to medium
Flooding frequency	None
Ponding frequency	None
Elevation	299–651 m
Slope	0–35%
Water table depth	203 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 55B is considered to have a continental climate – cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature are characteristic of the MLRA. The climate is the result of this MLRA's location in the geographic center of North America. There are few natural barriers on the northern Great Plains. The air masses move unobstructed across the plains and account for rapid changes in temperature.

Annual precipitation ranges from 18 to 23 inches per year. The normal average annual temperature is about 41.5° F. January is the coldest month with average low temperature ranging from about -4.3° F (Petersburg, ND) to about 2.5° F (Mellette, SD). July is the warmest month with temperatures averaging from about 79° F (Petersburg, ND) to about 84° F (Mellette, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 64° F. This large annual range attests to the continental nature of this MLRA's climate. Winds average about 11 miles per hour annually, ranging from about 13 miles per hour during the spring to about 10 miles per hour during the summer. Daytime winds are generally stronger than nighttime and strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour.

Growth of native cool-season plants begins in late March and continues to early to mid-July. Native warm-season plants begin growth in mid-May and continue to the end of August. Green-up of cool-season plants can occur in September and October when adequate soil moisture is present.

Table 3. Representative climatic features

Frost-free period (characteristic range)	111-117 days
Freeze-free period (characteristic range)	128-134 days

Precipitation total (characteristic range)	483-559 mm
Frost-free period (actual range)	105-119 days
Freeze-free period (actual range)	124-135 days
Precipitation total (actual range)	457-584 mm
Frost-free period (average)	114 days
Freeze-free period (average)	131 days
Precipitation total (average)	533 mm

Climate stations used

- (1) BUTTE 5SE [USC00321225], Butte, ND
- (2) CARRINGTON [USC00321360], Carrington, ND
- (3) FORMAN 5 SSE [USC00323117], Forman, ND
- (4) HARVEY 4NE [USC00324013], Harvey, ND
- (5) LA MOURE [USC00324937], Lamoure, ND
- (6) MELLETTE 4 W [USC00395456], Northville, SD
- (7) PETERSBURG 2 N [USC00327027], Petersburg, ND
- (8) COLUMBIA 8 N [USC00391873], Columbia, SD

Influencing water features

This site does not receive additional water as runoff from adjacent slopes; it is on a run-off landscape position. Neither does it receive additional water from a seasonal high-water table. Depth to the water table is deeper than 6 feet throughout the growing season. Infiltration is moderately rapid to very rapid. Permeability below the surface layer is rapid or very rapid. Water loss is through percolation below the root zone and evapotranspiration.

Soil features

Soils associated with Very Shallow ecological site are in the Mollisol and Entisol orders. The Mollisols are classified further as Entic Hapludolls. The Entisols are classified further as Typic Udorthents. These soils were developed under prairie vegetation. They formed in glaciofluvial deposits or beach deposits.

The common feature of soils in this site is the very shallow depth (<14 inches) to layers high in coarse sand and gravel (15 to 60 percent gravel). In some soils this layer contains significant amounts (>25%) of shale fragments. Due to the high amounts of gravel, these soils are very droughty and have limited plant production.

The soils are excessively drained. The thin surface layer is typically sandy loam, gravelly sandy loam, or loamy coarse sand; but gravelly loam, cobbly loam and coarse sandy loam also occur. Where soils have been cultivated, the surface layer may be very gravelly.

Salinity and sodicity are none. Soil reaction is neutral to moderately alkaline. Calcium carbonate content is none too low in the surface soil and low to moderately high below. Some soils have layers of calcium carbonate accumulation in the upper part of the gravelly layers (typically as coatings on undersides of rock fragments).

The soil surface is very unstable but intact. Sub-surface soil layers are restrictive to root penetration. These soils are mainly susceptible to water erosion. The hazard of water erosion increases on slopes greater than about 15 percent. Very low available water capacity caused by the shallow rooting depth strongly influences the soil/water/plant relationship. Loss of the soil surface layer can result in a shift in species composition and/or production.

Major soil series correlated to the Very Shallow site are Coe, Sioux, and Wamduska.

Access Web Soil Survey

(<https://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>) for specific local soils information.

Table 4. Representative soil features

Parent material	(1) Glaciofluvial deposits (2) Outwash (3) Till
Surface texture	(1) Gravelly, very gravelly, cobbly sandy loam (2) Gravelly, very gravelly, cobbly loamy coarse sand (3) Gravelly, very gravelly, cobbly loam (4) Gravelly, very gravelly, cobbly coarse sandy loam
Family particle size	(1) Sandy
Drainage class	Excessively drained
Permeability class	Rapid to very rapid
Depth to restrictive layer	5–36 cm
Surface fragment cover <=3"	3–40%
Surface fragment cover >3"	0–30%
Available water capacity (0-101.6cm)	2.54–6.35 cm
Calcium carbonate equivalent (0-101.6cm)	0–20%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	6.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	15–60%
Subsurface fragment volume >3" (Depth not specified)	0–20%

Ecological dynamics

This ecological site description is based on nonequilibrium ecology and resilience theory and utilizes a State- and-Transition Model (STM) diagram to organize and communicate information about ecosystem change as a basis for management. The ecological dynamics characterized by the STM diagram reflect how changes in ecological drivers, feedback mechanisms, and controlling variables can maintain or induce changes in plant community composition (phases and/or states). The application of various management actions, combined with weather variables, impact the ecological processes which influence the competitive interactions, thereby maintaining or altering plant community structure.

Prior to European influence, the historical disturbance regime for MLRA 55B included frequent fires, both anthropogenic and natural in origin. Most fires, however, were anthropogenic fires set by Native Americans. Native Americans set fires in all months except perhaps January. These fires occurred in two peak periods, one from March-May with the peak in April and another from July-November with the peak occurring in October. Most of these fires were scattered and of small extent and duration. The grazing history would have involved grazing and browsing by large herbivores (such as American bison, elk, and whitetail deer). Herbivory by small mammals, insects, nematodes, and other invertebrates are also important factors influencing the production and composition of the communities. Grazing and fire interaction, particularly when coupled with drought events, influenced the dynamics discussed and displayed in the following state and transition diagram and descriptions.

Following European influence, this ecological site generally has had a history of grazing by domestic livestock, particularly cattle, which along with other related activities (e.g., fencing, water development, fire suppression) has changed the disturbance regime of the site. Changes will occur in the plant communities due to these and other factors.

Weather fluctuations coupled with managerial factors may lead to changes in the plant communities and may, under adverse impacts, result in a slow decline in vegetative vigor and composition. However, under favorable conditions the botanical composition may resemble that prior to European influence.

Four vegetative states have been identified for the site (Reference, Native/Invaded, Invaded, and Go-Back). Within each state, one or more community phases have been identified. These community phases are named based on the more dominant and visually conspicuous species; they have been determined by study of historical documents, relict areas, scientific studies, and ecological aspects of plant species and plant communities. Transitional pathways and thresholds have been determined through similar methods.

State 1: Reference State represents the natural range of variability that dominated the dynamics of this ecological site prior to European influence. Dynamics of the state were largely determined by variations in climate and weather (e.g., drought), as well as that of fire (e.g., timing, frequency) and grazing by native herbivores (e.g., frequency, intensity, selectivity). Due to those variations, the Reference State is thought to have shifted temporally and spatially between two plant community phases.

Currently the primary disturbances include widespread introduction of exotic plants, concentrated livestock grazing, lack of fire, and perhaps long-term non-use or very light grazing, and no fire. Because of these changes particularly the widespread occurrence of exotic plants, as well as other environmental changes, the Reference State is considered to no longer exist. Thus, the presence of exotic plants on the site precludes it from being placed in the Reference State. It must then be placed in one of the other states, commonly State 2: Native/Invaded State (T1A).

State 2: Native/Invaded State. Colonization of the site by exotic plants results in a transition from State 1: Reference State to State 2: Native/Invaded State (T1A). This transition was inevitable; it often resulted from colonization by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass) which have been particularly and consistently invasive under long-term non-use or very light grazing, and no fire. Other exotic plants (e.g., Canada thistle) may also invade the site.

Three community phases have been identified for this state; they are similar to the community phases in the Reference State but have now been invaded by exotic cool-season grasses. These exotic cool-season grasses can be expected to increase. As that increase occurs, plants more desirable to wildlife and livestock may decline. A decline in forb diversity can also be expected. Under non-use or minimal use management, mulch increases and may become a physical barrier to plant growth. This also changes the micro-climate near the soil surface and may alter infiltration, nutrient cycling, and biological activity near the soil surface. As a result, these factors coupled with shading cause desirable native plants to have increasing difficulty remaining viable and recruitment declines.

To slow or limit the invasion of these exotic grasses or other exotic plants, it is imperative that managerial techniques (e.g., prescribed grazing, prescribed burning) be carefully constructed, monitored, and evaluated with respect to that objective. If management does not include measures to control or reduce these exotic plants, the transition to State 3: Invaded State should be expected (T2A).

State 3: Invaded State. The threshold for this state is reached when both the exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass) exceed 30% of the plant community and native grasses represent less than 40% of the community. One community phase has been identified for this state.

The exotic cool-season grasses can be quite invasive and often form monotypic stands. As they increase, both forage quantity and quality of the annual production becomes increasingly restricted to late spring and early summer, even though annual production may increase. Forb diversity often declines. Under non-use or minimal use management, mulch can increase and become a physical barrier to plant growth which alters nutrient cycling, infiltration, and soil biological activity. As such, desirable native plants become increasingly displaced.

Once the state is well established, prescribed burning and prescribed grazing techniques have been largely ineffective in suppressing or eliminating the exotic cool-season grasses, even though some short-term reductions may appear successful. However, assuming there is an adequate component of native grasses to respond to treatments, a restoration pathway to State 2: Native/Invaded State may be accomplished with the implementation of long-term prescribed grazing in conjunction with prescribed burning (R3A).

State 4: Go-Back State often results following cropland abandonment and consists of only one plant community phase. This weedy assemblage may include noxious weeds that need control. Over time, the exotic cool-season grasses (Kentucky bluegrass, smooth brome, and/or crested wheatgrass) will likely predominate.

Initially, due to extensive bare ground and a preponderance of shallow rooted annual plants, the potential for soil erosion is high. Plant species richness may be high, but overall diversity (i.e., equitability) is typically low, with the site dominated by a relatively small assemblage of species. Due to the lack of native perennials and other factors, restoring the site with the associated ecological processes is difficult. However, a successful range planting may result in something approaching State 2: Native/Invaded State (R4A). Following planting, prescribed grazing, prescribed burning, haying, and the use of herbicides will generally be necessary to achieve the desired result and control weeds, some of which may be noxious weeds. A failed range planting and/or secondary succession will lead to State 4: Invaded State (R4B).

Woody Invasion. Historically, individual (or small patches of) shrubs and/or trees were scattered across the site. However, a marked increase in fire suppression, climate change, increase in non-use, and other factors enabled woody species to colonize, form patches (resistant to fire), and begin to encroach on the site. These changes have enabled these patches to expand and become more widespread. Encroachment of both native and exotic woody species (e.g., Rocky Mountain juniper, Russian olive, Siberian elm, western snowberry, silverberry, ponderosa pine, eastern red cedar, etc.) are examples of woody vegetation increasing in extent and impinging on the ecological integrity of the grassland biome. Windbreaks and other tree plantings can contain problematic and invasive species (such as eastern redcedar, Rocky Mountain juniper, ponderosa pine, Russian olive, etc.), which can contaminate surrounding grasslands. This results in increased long-term costs to maintain or restore this ecological site in native grasses and forbs.

The following state and transition model diagram illustrates the common states, community phases, community pathways, and transition and restoration pathways that can occur on the site. These are the most common plant community phases and states based on current knowledge and experience; changes may be made as more data are collected. Pathway narratives describing the site's ecological dynamics reference various management practices (e.g., prescribed grazing, prescribed fire, brush management, herbaceous weed treatment) which, if properly designed and implemented, will positively influence plant community competitive interactions. The design of these management practices will be site specific and should be developed by knowledgeable individuals; based upon management goals and a resource inventory; and supported by an ongoing monitoring protocol.

When the management goal is to maintain an existing plant community phase or restore to another phase within the same state, modification of existing management to ensure native species have the competitive advantage may be required. To restore a previous state, the application of two or more management practices in an ongoing manner will be required. Whether using prescribed grazing, prescribed burning, or a combination of both with or without additional practices (e.g., brush management), the timing and method of application needs to favor the native species over the exotic species. Adjustments to account for variations in annual growing conditions and implementing an ongoing monitoring protocol to track changes and adjust management inputs to ensure desired outcome will be necessary.

The plant community phase composition table(s) has been developed from the best available knowledge including research, historical records, clipping studies, and inventory records. As more data are collected, plant community species composition and production information may be revised.

State and transition model



Figure 8. Regardless of specific ecological site, Eastern red cedar and Russian olive invasion on native rangeland in a formerly treeless grassland biome in MLRA 55B. Eastern red cedar and Russian olive seed source likely translocated by birds.

Plant Communities and Transitional Pathways




Diagram Legend - MLRA 55B Very Shallow

T1A	Introduction of exotic cool-season grasses
T2A	Long-term non-use or very light grazing, no fire
T5A	Cessation of annual cropping
R3A	Long-term prescribed grazing and prescribed burning
R4A	Successful range planting
R4B	Failed range planting and/or secondary succession
CP 1.1 - 1.2 (1.1A)	Multiyear drought with or without heavy grazing
CP 1.2 - 1.1 (1.2A)	Return to average precipitation with light to moderate grazing
CP 2.1 - 2.2 (2.1A)	Multiyear drought with or without heavy grazing
CP 2.1 - 2.3 (2.1B)	Long-term non-use or very light grazing, no fire
CP 2.2 - 2.1 (2.2A)	Return to average precipitation and prescribed long-term grazing and prescribed burning
CP 2.2 - 2.3 (2.2B)	Long-term non-use or very light grazing, no fire
CP 2.3 - 2.2 (2.3A)	Long-term prescribed grazing and prescribed burning
CP 2.3 - 2.1 (2.3B)	Long-term prescribed grazing and prescribed burning

Reference

This state represents the natural range of variability that dominated the dynamics of this ecological site prior to European influence. The primary disturbance mechanisms for this site in the reference condition included frequent fire and grazing by large herding ungulates. Timing of fires and grazing, coupled with weather events, dictated the dynamics that occurred within the natural range of variability. These factors likely caused the community to shift both spatially and temporally between two community phases.

Characteristics and indicators. Because of changes in disturbances and other environmental factors (particularly the widespread occurrence of exotic species), the Reference State is considered to no longer exist.

Resilience management. If intact, the reference state should probably be managed with current disturbance regimes which has permitted the site to remain in reference condition as well as maintaining the quality and integrity of associated ecological sites. Maintenance of the reference condition is contingent upon a monitoring protocol to guide management.

Community 1.1

Needle and Thread-Blue Grama/Sedges (*Hesperostipa comata*- *Bouteloua gracilis*/*Carex* spp.)

This community phase was historically the most dominant both temporally and spatially. Cool-season grass and grass-like species dominated this plant community with warm-season grasses being subdominant. The major grass or grass-like species included needle and thread, blue grama, western wheatgrass, and threadleaf sedge. Other grasses included Fendler threeawn, plains muhly, slender wheatgrass, sand dropseed, and prairie Junegrass. Common forb and shrub species included dotted blazing star, hairy false goldaster, blacksamson echinacea, prairie clover, prairie sagewort, and rose. Annual production likely varied from about 800-1900 pounds per acre with grasses and grass-likes, forbs, and shrubs contributing about 80%, 15% and 5%, respectively. Both warm-season and cool-season grasses were well represented in the community; as a result production was distributed throughout the growing season. This community represents the plant community phase upon which interpretations are primarily based and is described in the "Plant Community Composition and Group Annual Production" portion of this ecological site description.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	796	1357	1805
Forb	73	157	241
Shrub/Vine	28	55	84
Total	897	1569	2130

Figure 10. Plant community growth curve (percent production by month).
ND5502, Central Black Glaciated Plains, cool-season dominant, warm-season sub-dominant.. Cool-season dominant, warm-season sub-dominant..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	3	7	23	42	15	5	4	1	0	0

Community 1.2

Blue Grama/Sedges/Forbs/Prairie Sagewort (*Bouteloua gracilis*/*Carex* spp./*Forbs*/*Artemisia frigida*)

This community phase formed during multiyear drought with or without heavy grazing. Short grass and grass-like species increased to dominate the site; annual production decreased compared to Community Phase 1.1. Blue grama and threadleaf sedge were the dominant grass/grass-like species. Other grasses often included western wheatgrass, needle and thread, prairie Junegrass, and Fendler threeawn. Common forbs included field sagewort, cutleaf ironplant, scurfpea, white heath aster, and common yarrow. Prairie sagewort was the dominant shrub.

Pathway 1.1a

Community 1.1 to 1.2

Community Phase Pathway 1.1 to 1.2 occurred during multiyear drought with or without heavy grazing. This led to marked increases in blue grama, sedges, and prairie sagewort with a corresponding decrease in needle and thread.

Pathway 1.2a

Community 1.2 to 1.1

Community Phase Pathway 1.2 to 1.1 occurred with the return to average precipitation with light to moderate grazing. This led to a marked increase in needle and thread with corresponding decreases blue grama, sedges, and prairie sagewort.

State 2

Native/Invaded

This state is similar to State 1: Reference State but has now been colonized by the exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass) which are now present in small amounts. Other exotic plants (e.g., leafy spurge) may also invade the site. Although the state is still dominated by native grasses, an increase in these exotic cool-season grasses can be expected. These exotic cool-season grasses can be quite invasive on the site and are particularly well adapted to heavy grazing. They also often form monotypic stands. As these exotic cool-season grasses increase, both forage quantity and quality become increasingly restricted to late spring and early summer due to the monotypic nature of the stand, even though annual production may increase. Native forbs generally decrease in production, abundance, diversity, and richness compared to that of State 1: Reference State. These exotic cool-season grasses have been particularly and consistently invasive under extended periods of non-use and no fire. To slow or limit the invasion of these exotic grasses, it is imperative that managerial techniques (e.g., prescribed grazing, prescribed burning) be carefully constructed, monitored, and evaluated with respect to that objective. If management does not include measures to control or reduce these exotic cool-season grasses, the transition to State 3: Invaded State should be expected. Annual production of this state can be quite variable, in large part due to the amount of exotic cool-season grasses. However, as the exotic cool-season grasses increase, peak production will shift to earlier in the growing season.

Characteristics and indicators. The presence of trace amounts of exotic cool-season grasses indicates a transition from State 1 to State 2. The presence of exotic biennial or perennial leguminous forbs (i.e., sweet clover, black medic) may not, on their own, indicate a transition from State 1 to State 2 but may facilitate that transition.

Resilience management. To slow or limit the invasion of these exotic grasses, it is imperative that managerial techniques (e.g., prescribed grazing, prescribed burning) be carefully constructed, monitored, and evaluated with respect to that objective. Grazing management should be applied that enhances the competitive advantage of native grass and forb species. This may include: (1) grazing when exotic cool-season grasses are actively growing and native cool-season grasses are dormant; (2) applying proper deferment periods allowing native grasses to recover and maintain or improve vigor; (3) adjusting overall grazing intensity to reduce excessive plant litter (above that needed for rangeland health indicator #14 – see Rangeland Health Reference Worksheet); (4) incorporating early heavy spring utilization which focuses grazing pressure on exotic cool-season grasses and reduces plant litter, provided that livestock are moved when grazing selection shifts from exotic cool-season grasses to native grasses. Prescribed burning should be applied in a manner that maintains or enhances the competitive advantage of native grass and forb species. Prescribed burns should be applied as needed to adequately reduce/remove excessive plant litter and maintain the competitive advantage for native species. Timing of prescribed burns (spring vs. summer vs. fall) should be adjusted to account for differences in annual growing conditions and applied during windows of opportunity to best shift the competitive advantage to the native species.

Community 2.1

Needlegrasses-Blue Grama/Prairie Sagewort (*Hesperostipa comata*- *Bouteloua gracilis*/*Artemisia frigida*)



Figure 11. Community Phase 2.1: Needlegrasses-Blue Grama/Prairie Sagewort (foreground), dominated by green needlegrass, needle and thread, blue grama and prairie sagewort.

This community phase is similar to Community Phase 1.1 but has been colonized by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass). However, these exotics are present in smaller amounts with the community still dominated by native grasses. Annual production may be comparable to that of Community Phase 1.1 (800-1900 pounds per acre). However, as the exotic cool-season grasses increase, peak production will shift to earlier in the growing season.

Community 2.2

Blue Grama/ Sedges/ Forbs/ Prairie Sagewort (*Bouteloua gracilis*/ *Carex* spp./ Forbs/ *Artemisia frigida*)

This community phase is similar to Community Phase 1.2 but has now been colonized by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass). These exotics, however, are present in smaller amounts with the community still dominated by native grasses. This community phase is often dispersed throughout a pasture in an overgrazed/ undergrazed pattern, typically referred to as patch grazing. Some overgrazed areas will exhibit the impacts of heavy use, while the ungrazed areas will have a build-up of litter and increased plant decadence. This is a typical pattern found in properly stocked pastures grazed season-long. As a result, Kentucky bluegrass tends to increase more in the undergrazed areas while the more grazing tolerant short-statured species (such as blue grama and sedges) increase in the heavily grazed areas. If present, Kentucky bluegrass may increase under heavy grazing. Increasing amounts of exotic cool-season grasses, particularly Kentucky bluegrass, can make this an “at risk” community, even though its presence may not be obvious. If management does not include measures to control or reduce Kentucky bluegrass, the transition to State 3: Invaded State should be expected.

Community 2.3

Needle and Thread/ Sedges/ Exotic Cool-Season Grasses (*Hesperostipa comata*/ *Carex* spp./ Exotic Cool-Season Grasses)

This community phase results from long-term non-use or very light grazing, and no fire, which leads to a marked increase in the exotic cool-season grasses. Needle and thread is the main grass. Mulch accumulation, particularly that of Kentucky bluegrass leads to a marked decline in blue grama. This community phase is approaching the threshold leading to a transition to State 3: Invaded State. As a result, it is an “at risk” community. If management does not include measures to control or reduce these exotic cool-season grasses, the transition to State 3: Invaded State should be expected.

Pathway 2.1a

Community 2.1 to 2.2

Community Phase Pathway 2.1 to 2.2 occurs during multiyear drought with or without heavy grazing. This leads to marked increases in blue grama, sedges, and prairie sagewort with a corresponding decrease in needle and thread.

Pathway 2.1b

Community 2.1 to 2.3

Community Phase Pathway 2.1 to 2.3 occurs during long-term non-use or very light grazing, and no fire. This results in marked increases in the exotic cool-season grasses and corresponding decreases in native grasses.

Pathway 2.2a

Community 2.2 to 2.1

Community Phase Pathway 2.2 to 2.1 occurs with the return to average precipitation and implementation of long-term prescribed grazing and prescribed burning leading to a marked increase in needle and thread with corresponding decreases blue grama, sedges, and prairie sagewort.

Pathway 2.2b

Community 2.2 to 2.3

Community Phase Pathway 2.2 to 2.3 occurs during long-term non-use or very light grazing, and no fire, which leads to marked increases in the exotic cool-season grasses and corresponding decreases in native grasses, particularly warm-season species.

Pathway 2.3b

Community 2.3 to 2.1

Community Phase Pathway 2.3 to 2.1 occurs with the implementation of long-term prescribed grazing and prescribed burning, leading to marked decreases in the exotic cool-season grasses and corresponding increases in the native grasses (particularly warm-season species). This pathway is similar to that of Community Phase Pathway 2.3A but differs in the abundance of blue grama in Community Phase 2.3.

Pathway 2.3a

Community 2.3 to 2.2

Community Phase Pathway 2.3 to 2.2 occurs with the implementation of long-term prescribed grazing and prescribed burning, leading to a marked decrease in the exotic cool-season grasses and a corresponding increase in the native grasses (particularly warm-season species). This pathway is similar to that of Community Phase Pathway 2.3B but differs in the abundance of blue grama in Community Phase 2.3.

State 3

Invaded

This state is the result of invasion and dominance by the exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass). Other exotic plants (e.g., leafy spurge) may also invade the site. These exotic cool-season grasses can be quite invasive on the site and are particularly well adapted to heavy grazing. They also often form monotypic stands. As these exotic cool-season grasses increase, both forage quantity and quality become increasingly restricted to late spring and early summer due to the monotypic nature of the stand, even though annual production may increase. Native forbs generally decrease in production, abundance, diversity, and richness compared to that of State 1: Reference State. Common forb and shrub species often include white heath aster, goldenrod, common yarrow, and rose. Once the state is well established, prescribed burning and grazing techniques have been largely ineffective in suppressing or eliminating these species, even though some short-term reductions may appear successful. Annual production of this state may vary widely, in part due to variations in the extent of invasion by exotic cool-season grasses. However, as the exotic cool-season grasses increase, peak production will shift to earlier in the growing season.

Characteristics and indicators. This site is characterized by exotic cool-season grasses constituting greater than 30 percent of the annual production and native grasses constituting less than 40 percent of the annual production.

Resilience management. Light or moderately stocked continuous, season-long grazing or a prescribed grazing system which incorporates adequate deferment periods between grazing events and proper stocking rate levels will maintain this State. Application of herbaceous weed treatment, occasional prescribed burning and/or brush

management may be needed to manage noxious weeds and increasing shrub (e.g., western snowberry) populations.

Community 3.1

Exotic Cool-Season Grasses/Forbs

This community phase is dominated by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass), often with a much-reduced forb and shrub component. Common forbs and shrub species often include white heath aster, goldenrod, common yarrow, and rose. Other exotic plants (e.g., leafy spurge) may also invade the site. Annual production of this state may vary widely, in part due to variations in the extent of invasion by exotic cool-season grasses. However, as the exotic cool-season grasses increase, peak production will shift to earlier in the growing season. The longer this community phase exists, the more resilient it becomes. Natural or management disturbances that reduce the cover of Kentucky bluegrass or smooth brome are typically short-lived.

State 4

Go-Back

This state is highly variable depending on the level and duration of disturbance related to the T5A transitional pathway. In this MLRA, the most probable origin of this state is plant succession following cropland abandonment. This plant community will initially include a variety of annual forbs and grasses, some of which may be noxious weeds and need control. Over time, the exotic cool-season grasses (Kentucky bluegrass, smooth brome, and/or crested wheatgrass) will likely predominate.

Characteristics and indicators. Tillage has destroyed the native plant community, altered soil structure and biology, reduced soil organic matter, and resulted in the formation of a tillage induced compacted layer which is restrictive to root growth. Removal of perennial grasses and forbs results in decreased infiltration and increased runoff.

Resilience management. Continued tillage will maintain the state. Control of noxious weeds will be required.

Community 4.1

Annual/Pioneer Perennial/Exotics

This community phase is highly variable depending on the level and duration of disturbance related to the T5A transitional pathway. In this MLRA, the most probable origin of this phase is secondary succession following cropland abandonment. This plant community will initially include a variety of annual forbs and grasses, including noxious weeds (e.g., leafy spurge) which may need control. Over time, the exotic cool-season grasses (Kentucky bluegrass, smooth brome, and/or crested wheatgrass) will likely predominate.

State 5

Any Plant Community

Transition T1A

State 1 to 2

This is the transition from the State 1: Reference State to the State 2: Native/Invaded State due to the introduction and establishment of exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass). This transition was inevitable and corresponded to a decline in native warm-season and cool-season grasses; it may have been exacerbated by chronic season-long or heavy late season grazing. Complete rest from grazing and suppression of fire could also have hastened the transition. The threshold between states was crossed when Kentucky bluegrass, smooth brome, crested wheatgrass, or other exotic species became established on the site.

Constraints to recovery. Current knowledge and technology will not facilitate a successful restoration to Reference State.

Transition T2A

State 2 to 3

This transition from the State 2: Native/Invaded State to State 3: Invaded State generally occurs during long-term non-use or very light grazing, and no fire. Exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass) become the dominant graminoids. Studies indicate that a threshold may exist in this transition when both the exotic cool-season grasses exceed 30% of the plant community and native grasses represent less than 40% of the plant community composition. This transition may occur under other managerial conditions, for example heavy season-long grazing (primarily Kentucky bluegrass).

Constraints to recovery. Variations in growing conditions (e.g., cool, wet spring) will influence effects of various management activities on exotic cool-season grass populations.

Restoration pathway R3A

State 3 to 2

This restoration pathway from State 3: Invaded State to State 2: Native/Invaded State may be accomplished with the implementation of long-term prescribed grazing and prescribed burning, assuming there is an adequate component of native grasses to respond to the treatments. Both prescribed grazing and prescribed burning are likely necessary to successfully initiate this restoration pathway, the success of which depends upon the presence of a remnant population of native grasses in Community Phase 3.1. That remnant population, however, may not be readily apparent without close inspection. The application of several prescribed burns may be needed at relatively short intervals in the early phases of this restoration process, in part because some shrubs (e.g., rose) sprout following one burn. Early season prescribed burns have been successful; however, fall burning may also be an effective technique. The prescribed grazing should include adequate recovery periods following each grazing event and stocking levels which match the available resources. If properly implemented, this will shift the competitive advantage from the exotic cool-season grasses to the native cool-season grasses.

Context dependence. Grazing management should be applied in a manner that enhances/maximizes the competitive advantage of native grass and forb species over the exotic species. This may include the use of prescribed grazing to reduce excessive plant litter accumulations above that needed for rangeland health indicator #14 (see Rangeland Health Reference Worksheet). Increasing livestock densities may facilitate the reduction in plant litter provided length and timing of grazing periods are adjusted to favor native species. Grazing prescriptions designed to address exotic grass invasion and favor native species may involve earlier, short, intense grazing periods with proper deferment to improve native species health and vigor. Fall (e.g., September, October) prescribed burning followed by an intensive, early spring graze period with adequate deferment for native grass recovery may shift the competitive advantage to the native species, facilitating the restoration to State 2: Native/Invaded. Prescribed burning should be applied in a manner that enhances the competitive advantage of native grass and forb species over the exotic species. Prescribed burns should be applied at a frequency which mimics the natural disturbance regime, or more frequently as is ecologically (e.g., available fuel load) and economically feasible. Burn prescriptions may need adjustment to: (1) account for change in fine fuel orientation (e.g., "flopped" Kentucky bluegrass); (2) fire intensity and duration by adjusting ignition pattern (e.g., backing fires vs head fires); (3) account for plant phenological stages to maximize stress on exotic species while favoring native species (both cool- and warm-season grasses).

Restoration pathway R4A

State 4 to 2

This restoration pathway from State 4: Go-Back State to the State 2: Native/Invaded State can be accomplished with a successful range planting. Following planting, prescribed grazing, prescribed burning, haying, or use of herbicides will generally be necessary to achieve the desired result and control any noxious weeds. It may be possible using selected plant materials and agronomic practices to approach something very near the functioning of State 2: Native/Invaded State. Application of chemical herbicides and the use of mechanical planting methods using adapted varieties of the dominant native grasses are possible and can be successful. After establishment of the native plant species, prescribed grazing should include adequate recovery periods following each grazing event and stocking levels which match the available resources; management objectives must include the maintenance of those species, the associated reference state functions, and continued treatment of exotic grasses.

Context dependence. A successful range planting will include proper seedbed preparation, weed control (both

prior to and after the planting), selection of adapted native species representing functional/structural groups inherent to the State 1, and proper planting technique. Management (e.g., prescribed grazing, prescribed burning) during and after establishment must be applied in a manner that maintains the competitive advantage for the seeded native species. Adding non-native species can impact the above and below ground biota. Elevated soil nitrogen levels have been shown to benefit smooth brome and Kentucky bluegrass more than some native grasses. As a result, fertilization, exotic legumes in the seeding mix, and other techniques that increase soil nitrogen may promote smooth brome and Kentucky bluegrass invasion. The method or methods of herbaceous weed treatment will be site specific to each situation; but generally, the goal would be to apply the pesticide, mechanical control, or biological control (either singularly or in combination) in a manner that shifts the competitive advantage from the targeted species to the native grasses and forbs. The control method(s) should be as specific to the targeted species as possible to minimize impacts to non-target species.

Restoration pathway R4B State 4 to 3

A failed range planting and/or secondary succession will lead to State 3: Invaded State.

Context dependence. Failed range plantings can result from many causes (both singularly and in combination) including drought, poor seedbed preparation, improper planting methods, seeded species not adapted to the site, insufficient weed control, herbicide carryover, poor seed quality (purity & germination), and/or improper management.

Transition T5A State 5 to 4

This transition from any plant community to State 4: Go-Back State. It is most commonly associated with the cessation of cropping without the benefit of range planting, resulting in a “go-back” situation. Soil conditions can be quite variable on the site, in part due to variations in the management/cropping history (e.g., development of a tillage induced compacted layer (plow pan), erosion, fertility, and/or herbicide/pesticide carryover). Thus, soil conditions should be assessed when considering restoration techniques.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Cool-season Grasses			235–706	
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	157–471	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	31–157	–
	slender wheatgrass	ELTR7	<i>Elymus trachycaulus</i>	0–78	–
	slender wheatgrass	ELTRS	<i>Elymus trachycaulus ssp. subsecundus</i>	0–78	–
2	Warm-season Grasses			235–549	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	78–314	–
	plains muhly	MUCU3	<i>Muhlenbergia cuspidata</i>	16–157	–
	Fendler threeawn	ARPUL	<i>Aristida purpurea var. longiseta</i>	16–47	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	16–31	–
3	Other Native Grasses			16–78	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–47	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	16–31	–
4	Grass-likes			78–235	
	threadleaf sedge	CAFI	<i>Carex filifolia</i>	78–204	–
	Grass-like (not a true grass)	2CI	Grass-like (not a true grass)	0–78	–

	Grass-like (not a true grass)	ZGL	Grass-like (not a true grass)	0-10	-
Forb					
5	Forbs			78-235	
	blazing star	LIATR	<i>Liatris</i>	16-126	-
	hairy false goldenaster	HEVIV	<i>Heterotheca villosa var. villosa</i>	16-63	-
	silverleaf Indian breadroot	PEAR6	<i>Pediomelum argophyllum</i>	0-63	-
	prairie clover	DALEA	<i>Dalea</i>	16-47	-
	blacksamson echinacea	ECAN2	<i>Echinacea angustifolia</i>	16-47	-
	velvety goldenrod	SOMO	<i>Solidago mollis</i>	16-47	-
	field sagewort	ARCA12	<i>Artemisia campestris</i>	16-31	-
	lacy tansyaster	MAPIP4	<i>Machaeranthera pinnatifida ssp. pinnatifida var. pinnatifida</i>	16-31	-
	spiny phlox	PHHO	<i>Phlox hoodii</i>	16-31	-
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	16-31	-
	white heath aster	SYER	<i>Symphotrichum ericoides</i>	16-31	-
	Forb, perennial	2FP	<i>Forb, perennial</i>	16-31	-
	Forb, annual	2FA	<i>Forb, annual</i>	16-31	-
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0-16	-
	onion	ALLIU	<i>Allium</i>	0-16	-
	pussytoes	ANTEN	<i>Antennaria</i>	0-16	-
	plains milkvetch	ASGI5	<i>Astragalus gilviflorus</i>	0-16	-
	rush skeletonplant	LYJU	<i>Lygodesmia juncea</i>	0-16	-
	scarlet beeblossom	OESU3	<i>Oenothera suffrutescens</i>	0-16	-
	cutleaf anemone	PUPAM	<i>Pulsatilla patens ssp. multifida</i>	0-16	-
Shrub/Vine					
6	Shrubs			31-78	
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	16-63	-
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0-47	-
	rose	ROSA5	<i>Rosa</i>	16-31	-

Animal community

Animal Community – Wildlife Interpretations

Landscape

The MLRA 55B landscape is characterized by mostly nearly level to gently rolling till plains with some steep slopes adjacent to streams and many poorly defined drainage channels. The continental drainage divide occurs in the east central part of the MLRA. The MLRA is located within the Prairie Pothole Region with temporary, seasonal, and semi-permanent wetlands throughout the MLRA. The MLRA includes areas of eskers, kames, and ground moraines. MLRA 55B is considered to have a continental climate with cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature are common and characteristic of the MLRA. This area supports mid- to tall-grass prairie vegetation with American elm, bur oak, green ash, and willow species growing along the riparian zones of river systems found throughout the MLRA. Complex intermingled ecological sites create diverse grass/shrub land habitats interspersed with varying densities of linear, slope, depressional, and in-stream

wetlands associated with headwater streams and tributaries of the James, Pipestem, Maple, Goose, Sheyenne, Wild Rice, and Elm Rivers. MLRA 55B is located within North and South Dakota and within the boundaries of the Prairie Pothole Region.

Three Hydrologic Unit Areas make up this MLRA. Approximately 6% drains into the Mouse River into MLRA 55A, with the balance split between the James and Sheyenne Rivers.

By the mid-19th century, over 76% of the MLRA had been converted from mid- to tall-grass prairie to annual crop production. To alleviate crop production loss from wetlands and overland flow, a system of shallow surface ditches, judicial ditches, and road ditches removes surface water in spring and during high rainfall events. Tile drainage systems have been or are being installed extensively throughout MLRA 55B for sub-surface field drainage to enhance annual crop production.

Historic Communities/Conditions within MLRA 55B:

The northern tall- and mixed-grass prairie were disturbance-driven ecosystems with fire, herbivory, and climate functions as the primary ecological drivers (either singly or often in combination). American bison roamed MLRA 55B, wintering along the Mouse River in MLRA 55A and migrating through MLRA 55B and into MLRA 56A. Many species of grassland birds, small mammals, insects, reptiles, amphibians, elk, moose, pronghorn, white-tailed deer, and large herds of American bison were historically among the inhabitants adapted to this region. Roaming herbivores, as well as several small mammals and insect species, were the primary consumers linking the grassland resources to large predators (such as the wolf, American black bear, grizzly bear) and smaller carnivores (such as the coyote, bobcat, red fox, and raptors). Extirpated species include free-ranging American bison and gray wolf (breeding). Extinct is the Rocky Mountain locust.

Present Communities/Conditions within MLRA 55B:

This area supports natural prairie vegetation characterized by western wheatgrass, green needlegrass, needle and thread, and blue grama. Little bluestem is an important species on the more sloping and shallower soils. Prairie cordgrass, northern reedgrass, big bluestem, and wheat sedge (aka slough sedge) are important species on wet soils. Western snowberry, leadplant, and prairie rose are commonly interspersed throughout the area.

Over 80% of MLRA 55B has been converted to annual crop production. These influences fragmented the landscape, reduced or eliminated ecological drivers (fire), and introduced exotic plant species including smooth brome, crested wheatgrass, Kentucky bluegrass, and leafy spurge; this further impacted plant and animal communities. The loss of the bison and fire as primary ecological drivers greatly influenced the character of the remaining native plant communities and the associated wildlife, moving towards a less diverse and more homogeneous landscape. Annual cropping is the main factor contributing to habitat fragmentation, reducing habitat quality for area-sensitive species.

Hydrological manipulation is extensive throughout the MLRA. Extensive wetland and subsurface tile drainage have taken place. Straightened segments of ephemeral and intermittent tributary streams of the James, Wild Rice, and Sheyenne River have reduced sinuosity, created oxbows, and enabled the conversion of riparian ecological sites to annual crop production. These anthropogenic impacts have reduced flood water detention and retention on the landscape. The results have been increasing storm water runoff sediment and nutrient loading to the James and Sheyenne Rivers and their tributaries (along with lakes and reservoirs within the MLRA). Large dams on the James, Pipestem and Sheyenne rivers, along with installation of instream structures have reduced aquatic species movement within the MLRA.

National wildlife refuges, waterfowl production areas, state wildlife management areas, and North and South Dakota Department of Trust Lands provide herbaceous and woody cover for wildlife. In addition, the United States Army Corps of Engineers and the United States Bureau of Reclamation manage three man-made reservoirs - Jamestown Reservoir, Pipestem Reservoir, and Lake Ashtabula for flood control, also providing fish habitat and adjacent uplands for wildlife cover. Lonetree Wildlife Management Area (WMA) is the largest state managed wildlife area covering 32,800 acres. Arrowwood National Wildlife Refuge is the largest refuge consisting of 16,000 acres.

Characteristic wildlife species in this area are:

Birds: Common goldeye, bufflehead, broad-winged hawk, alder flycatcher, mourning warbler, mallard, blue-winged teal, red-tailed hawk, American kestrel, killdeer, eastern and western kingbird, western meadowlark, American crow, common yellowthroat, clay-colored sparrow, vesper sparrow, red-necked grebe, Savannah sparrow, downy and hairy woodpeckers, black-capped chickadee, white-breasted nuthatch, and brown-headed cowbird.

Mammals: Northern short-tailed shrew, white-tailed jackrabbit, Franklin's ground squirrel, thirteen-lined ground squirrel, northern pocket gopher, plains pocket gopher, western harvest mouse, deer mouse, meadow vole, meadow jumping mouse, western jumping mouse, coyote, red fox, racoon, American badger, striped skunk, white-tailed deer, elk, moose, beaver, muskrat, mink, weasel, woodchuck, and red, eastern gray and fox squirrels.

Reptiles/Amphibians: American toad, Great Plains toad, northern leopard frog, chorus frog, tiger salamander, plains garter snake, smooth green snake, wood frog, and common garter snake.

Presence of wildlife species is often determined by ecological site characteristics including grass and forb species, tree and shrub species, hydrology, aspect, and other associated ecological sites. The home ranges of a majority of species are usually larger than one ecological site or are dependent upon more than one ecological site for annual life requisites. Ecological sites offer different habitat elements as the annual life requisites change. Habitat improvement and creation must be conducted within the mobility limits of a known population for the species.

Insects play an important role in providing ecological services for plant community development. Insects that are scavengers or aid in decomposition provide the food chain baseline sustaining the carnivorous insects feeding upon them. Many insects provide the ecological services necessary for pollination, keeping plant communities healthy and productive. Insects provide a protein food source for numerous species including grassland-nesting birds, woodpeckers, woodland edge and interior species, and their young. Extensive use of insecticides for specialty crops (such as soybeans, corn, and other crops) has greatly reduced insects within this MLRA.

Species of Concern within MLRA 55B:

The following is a list of species considered "species of conservation priority" in the North Dakota State Wildlife Action Plan (2015) and South Dakota State Wildlife Action Plan (2014); and species listed as "threatened, endangered, or petitioned" under the Endangered Species Act within MLRA 55B at the time this section was developed:

Invertebrates: Dakota skipper, Iowa skipper, monarch butterfly, northern sandy tiger beetle, Ottoe skipper, Poweshiek skipperling, regal fritillary, yellow-banded bumble bee, and western bumble bee.

Birds: American avocet, American bittern, American kestrel, American white pelican, Baird's sparrow, bald eagle, black-billed cuckoo, black tern, bobolink, burrowing owl, canvasback, chestnut-collared longspur, Dickcissel, ferruginous hawk, Franklin's gull, grasshopper sparrow, horned grebe, lark bunting, LeConte's sparrow, lesser scaup, marbled godwit, Nelson's sparrow, northern goshawk, northern harrier, northern pintail, osprey (migration), peregrine falcon (migration), piping plover (migration), red knot (migration), sharp-tailed grouse, short-eared owl, Swainson's hawk, upland sandpiper, western meadowlark, willet, Wilson's phalarope, whooping crane (migration), and yellow rail.

Mammals: Arctic shrew, big and little brown bats, Franklin's ground squirrel, northern river otter, plains pocket mouse, pygmy shrew, Richardson's ground squirrel, and silver-haired bat.

Amphibians and Reptiles: Canadian toad, plains hognose snake, smooth green snake, and snapping turtle.

Fish and Mussels: Black sandshell, blacknose shiner, Carmine shiner, creek heelsplitter, creeper, deertoe, fragile papershell, mapleleaf, northern pearl dace, northern redbelly dace, pink heelsplitter, threeridge, trout-perch, yellow sandshell, and Wabash pigtoe.

Grassland Management for Wildlife in MLRA 55B

Management activities within the community phase pathways impact wildlife but are essential for maintenance of healthy grassland ecosystems. Community phase, transitional, and restoration pathways are keys to long-term management within each State and between States. Timing, intensity, and frequency of these inputs can have

dramatic positive or negative effects on local wildlife species. Ranchers and other land managers must always consider the long-term beneficial management effects of grassland and woodland resources in comparison to typically short-term negative effects to the habitats of individual species.

Ecological sites occur as intermingled complexes on the landscape with gradual or sometimes abrupt transitions. Rarely do ecological sites exist in large enough acreage to manage independently. Ecological sites supporting a dominance of herbaceous vegetation (Loamy/Sandy) can be located adjacent to ecological sites that support medium to tall shrubs, Invaded Wooded States. Conversely, ecological sites that are dominated by short- to mid-statured grasses (Claypan) can be adjacent to sites with bare soil only supporting minor amounts of short grasses and forbs (Thin Claypan).

Management of these complex ecological sites can provide a heterogeneous or a homogenous landscape. Grassland bird use declines as the plant community transitions to a homogenous state. Managers must recognize ecological sites and the complexes in which they occur to properly manage the landscape. A management regime for one ecological site may negatively impact an adjacent site; for example, alteration of a grazing regime within the Invaded Wooded State ecological sites to encourage understory growth may encourage exotic cool-season grasses to increase or dominate an adjacent ecological site.

Life requisites and habitat deficiencies are determined for targeted species. Deficiencies need to be addressed along community phase, transitional, and restoration pathways as presented in specific state-and-transition models. Ecological sites should be managed and restored within the site's capabilities to provide sustainable habitat for targeted species or species guilds. Managers also need to consider vegetative associations provided by adjacent/intermingled ecological sites for species with home ranges or life requisites that may not be provided by one ecological site.

Grassland-nesting birds use various grass heights for breeding, nesting, foraging, or winter habitat. While most species use varying heights, many have a preferred vegetative stature height or sensitivity to woody vegetation. Understanding the sensitivity of grassland species to woody vegetation and preferred vegetative structure enables managers to determine which grassland-nesting bird species avoid grassland habitats adjacent to woody dominated plant community phases. The following chart provides sensitivity to woody vegetation and preferred vegetative stature heights.

To see the chart, please follow the hyperlink:

https://efotg.sc.egov.usda.gov/references/public/ND/55B_Very_Shallow_Narrative_FINALRef_FSG.pdf

Very Shallow Wildlife Habitat Interpretation:

Very Shallow ecological sites are excessively drained and have coarse sand and/or gravel within 14 inches of the surface. These sites are usually found on slopes ranging from 0 to 35 percent. Needlegrasses are the dominant mid cool-season grass. Associated ecological sites include Loamy, Thin Loamy, Shallow Gravel, and Subirrigated. This site, generally, is not associated with riparian, wetland, or water features; however, a few areas do occur on high beach ridges near lakes.

Due to the high amounts of coarse sand and gravel, these soils are very droughty and have limited plant production. Very Shallow habitat features and components commonly support grassland- nesting birds the prefer short-statured vegetation and bare ground. Insects rely on associated forbs and grasses for survival. Insect populations provide food sources for birds and their young. Very Shallow ecological sites provide forage for small and large herbivores.

1.0 Reference State

Community Phase 1.1 Needle and Thread-Blue Grama/Sedges: This plant community provides quality vegetative cover for wildlife; every effort should be made to maintain this ecological site within this community phase. Even though production is limited due to droughty soil conditions, this plant community retain high functionality through continued maintenance including prescribed grazing (with adequate recovery period), as well as prescribed fire. Prescribed fire frequency maintains a grass- dominated plant community providing habitat for bird species reliant on a diverse plant community (structure and native composition). Predominance of grass species favors grazers and mixed feeders (animals selecting grasses as well as forbs and shrubs). The structural diversity provides habitat for a migratory and resident birds the prefer short open density plant structure.

Invertebrates: Insects play a role in maintaining the forb community and provide a forage base for grassland birds, reptiles, and rodents. Ecological services, historically provided by bison, are simulated by domestic livestock. These services include putting plant material and dung in contact with mineral soil to be used by lower trophic level consumers (such as invertebrate decomposers, scavengers, shredders, predators, herbivores, dung beetles, and fungal-feeders).

Lack of host plants, such as little bluestem, limits use by Dakota and Ottoe skippers. Violet species are not common on this site, not supporting the needed habitat for the regal fritillary. Monarch butterfly may use flowering forbs; however, few milkweed species are found on this site to support caterpillar food. Bumblebees and other native bees utilize forbs as a nectar source; bare ground and nesting sites are available due to the dominance of bunch grasses.

Birds: This plant community provides nesting, foraging, and escape habitats favored by short-grass nesting birds. This plant community can provide suitable areas for sharp-tailed grouse lek sites; however, coarse soils and steep slopes may reduce suitability. This site is not preferred nesting habitat, brood-rearing, and loafing habitat. This site provides limited hunting opportunities for grassland raptors.

Mammals: The diversity of grasses and forbs provide high nutrition levels for small and large herbivores. Short- to mid-statured vegetation provides suitable food, thermal, protective, and escape cover for small herbivores, but provides no thermal, protective, or escape cover for large ungulates.

Amphibians and Reptiles: This ecological site is not typically found near Wet Meadow or Wetland ecological sites; therefore, they do not provide habitat for the northern leopard frog and Canadian toad. These sites do not provide habitat for the northern prairie skink and plains hog-nosed snake.

Fish and Mussels: This ecological site is not typically located adjacent to streams, rivers, or water bodies. They do not receive run-on hydrology from adjacent ecological sites and provide limited hydrology to adjacent down-slope ecological sites, such as Loamy and Subirrigated. Management on Very Shallow sites, in conjunction with neighboring run-on sites, will have an indirect effect on aquatic species in streams and/or tributaries receiving water from adjacent sites. Optimum hydrological function and nutrient cycling limit potential for sediment yield and nutrient loading to the any nearby aquatic ecosystems from Community Phase 1.1.

Community Phase 1.2 Blue Grama/Sedges/Forbs/Prairie Sagewort: Multiyear drought with or without heavy long-term grazing (via Community Phase Pathway 1.1A) causes an evident increase in short grasses and sedges and a corresponding decrease in mid-grass species. The reduction in bunch grasses and the increase in sod forming, warm season-grasses and sedges decrease bare ground and create a shorter statured plant community. Forb diversity is somewhat reduced, but forbs such as field sagewort, silverleaf Indian breadroot, white heath aster, common yarrow, and blazing star provide nectar and pollen. Prairie sagewort becomes the principal shrub. Return to normal precipitation and reduced grazing pressure with adequate recovery periods (via Community Phase Pathway 1.2A) will shift the competitive edge back to a needle and thread and sedge dominated plant community.

Invertebrates: Provides similar life requisites as Community Phase 1.1. Wind-pollinated prairie sagewort reduces pollen and nectar availability for native bees and larval food butterfly species. Forb species may produce fewer flowers reducing overall pollen and nectar sources during periods of multiyear drought and heavy grazing. Bare ground may decrease but due to the droughtiness of these soils, bare ground is still available for ground nesting bees.

Birds: A shift to shorter statured grasses provide nesting, foraging, and escape habitats favored by short-statured grassland nesting birds. This plant community provides suitable areas for sharp-tailed grouse lek sites, dependent upon slope steepness. This site provides hunting opportunities for grassland raptors.

Mammals: Provides similar life requisites as Community Phase 1.1; however, the increase in short- statured, sod-forming warm season grasses and sedges reduces plant diversity along with thermal, escape, and protective cover for small mammals. Foliage and stems of prairie sagewort provide browse for jackrabbits, cottontails, ground squirrels, and various mice species.

Amphibians and Reptiles: Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: The increase in sod-forming grasses and sedges may increase run-off and provide increased hydrology to adjacent down-slope ecological sites.

2.0 Native/Invaded State

Community Phase 2.1 Needle and Thread-Blue Grama/Prairie Sagewort: This plant community develops through Transition Pathway T1A due to changes in management and the presence of exotic, cool-season grasses. Lack of fire or heavy season-long grazing can facilitate this transition. The threshold between States 1.0 and 2.0 is crossed when Kentucky bluegrass, smooth brome, or other exotic species become established. This plant community phase has a very similar appearance and function to the Reference State of Community 1.1, except it has a minor amount of exotic cool-season grasses and forbs. This phase functions at a high level for native wildlife; therefore, managers should consider the State 2.0 community phase pathways to avoid transitioning to the 3.0 Invaded State. There is no known Community Phase Pathway back to 1.0 Reference State from 2.0 Native/Invaded State.

Invertebrates: Provides similar life requisites as Community Phase 1.1.

Birds: Provides similar life requisites as Community Phase 1.1.

Mammals: Provides similar life requisites as Community Phase 1.1.

Amphibians and Reptiles: Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: Provides similar life requisites as Community Phase 1.1.

Community Phase 2.2 Blue Grama/Sedges/Forbs/Prairie Sagewort: Multiyear drought with or without heavy grazing, via Community Phase Pathway 2.1A, causes an evident increase in blue grama and sedges and a corresponding decrease in mid-grass native species. The reduction in bunch grasses and the increases in sod forming warm season-grasses, sedges, and cool-season exotic grasses decrease bare ground and create a shorter statured plant community. Prairie sagewort becomes the principal shrub. Return to normal precipitation and reduced grazing pressure with adequate recovery periods (via Community Phase Pathway 2.2A) will shift the competitive edge back to a green needlegrass and western wheatgrass dominated plant community.

This community phase is often found in a mosaic in the pasture in an overgrazed/undergrazed pattern typical of properly stocked pastures grazed season-long. Some areas will be impacted by heavy use while other areas will have a build-up of litter and a high amount of plant decadence. This mosaic of grazed and ungrazed areas provides a short to tall vegetative stature. Dependent upon the patch size of overgrazed vs. undergrazed areas, grassland nesting birds preferring short/mid- vegetative stature may prefer this plant community phase.

Invertebrates: Grazing tolerant forbs remain in this plant community providing pollen and nectar sources for pollinating species. However, wind pollinator species (such as prairie sagewort) increasing could potentially reduce nectar and pollen availability. Sufficient bunch grasses remain to provide bare ground for ground nesting bees.

Birds: Provides similar life requisites as Community Phase 1.2. However, depending on the amount of overgrazed vs. undergrazed area, vegetative stature could favor short-grass nesting bird species on over-grassed areas.

Mammals: Vegetation stature is reduced due to the increase in sedges and Kentucky bluegrass reducing cover. However, depending on the amount of overgrazed vs. undergrazed area, vegetative stature in the undergrazed areas could provide thermal and escape cover for mammals, especially small mammals.

Amphibians and Reptiles: Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: Provides similar life requisites as Community Phase 1.1.

Community Phase 2.3 Needle and Thread/Sedges/Exotic Cool-Season Grasses: Extended periods of non-use or very light grazing, and no fire (via Community Pathway 2.1B) allows exotic cool-season grasses to increase while western wheatgrass and blue grama decrease. However, the reduction in fire frequency will allow forbs to increase. Multi-year prescribed fire and prescribed grazing can move this plant community back to 2.1. This is an "at risk"

plant community with high amounts of Kentucky bluegrass out-completing native grasses. Every effort should be made by managers to avoid implementing a grazing system that may favor reaching the 30 percent threshold for Kentucky bluegrass via Transitional Pathway T2A. The longer this plant community transitions over to State 3.0, the opportunity for restoration back to State 2.0 by implementing a prescribed grazing system and introducing prescribed fire decreases.

Invertebrates: An increase exotic cool-season grass and a reduction in forb diversity reduces pollen and nectar sources, limiting use by pollination insect species. Common forbs include common yarrow, white sagebrush, and field sagewort.

Birds: Grassland-nesting birds that favor short-statured vegetative cover will use this site. However, continuous non-use, no fire, increases exotic cool-season exotic grasses reducing use by grassland nesting birds.

Mammals: As exotic cool-season grasses increase, thermal, protective or escape cover becomes limited.

Amphibians and Reptiles: Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: Although, this ecological site is not typically adjacent to streams, rivers, or water bodies, runoff from Community Phase 2.3 increases due to management that increases short-statured grass and reduces bunch grasses. Management of this community phase, in conjunction with neighboring run-on sites, will have an indirect negative effect on aquatic species in streams and/or tributaries receiving water from Sands and adjacent sites.

3.0 Invaded State

Community Phase 3.1 Exotic Cool-Season Grasses/Forbs: Complete rest from grazing and elimination of fire (via Transitional Pathway T2A) are the two major contributors to this transition, allowing exotic cool-season grasses to dominate this site. Native grasses may still be present but with reduced vigor and numbers - not allowing for recovery without multi-year prescribed grazing and burning (via Restoration Pathway R3A). Continued non-use may allow Kentucky bluegrass to become the dominant herbaceous component.

Invertebrates: Overall, pollinator plant diversity is reduced, with forbs (such as common yarrow) and shrubs (such as rose) providing limited pollen and nectar production. Light or no grazing will increase litter amounts; dense Kentucky bluegrass sod and sedges limit bare ground for ground nesting bees. Pollinator species may need to rely on adjacent ecological sites to meet their habitat needs.

Birds: This homogenous landscape provides limited structural diversity and insect production for grassland nesting birds. Grassland nesting birds preferring increased plant litter and favoring short-statured grasses will benefit, but long-term continued rest reduces nesting habitat for most species. This site provides limited hunting opportunities for grassland raptors.

Mammals: This short-statured community provides limited food, thermal, protective, and escape cover for large herbivores; life requisites will not be met in this plant community. Higher litter amounts provide food, thermal, protective, and escape cover for small herbivores.

Amphibians and Reptiles: This site should provide similar life requisites as Community Phase 1.1.

Fish and Mussels: Provides similar life requisites as Community Phase 1.1; however, increased runoff from this plant community may increase water yields to adjacent ecological sites.

4.0 Go-Back State

Community Phase 4.1 Annual/Pioneer Perennial/Exotics: This plant community is the result of severe soil disturbance (such as cropping, recreational activity, or concentrated livestock activity) for a prolonged time period. Following cessation of disturbances, the resulting plant community is dominated by early pioneer annual and perennial plant species. Plant species composition and production are highly variable dependent upon duration of disturbance. Weedy plants can provide pollinator habitat along with spring and summer cover for many mammals and birds, and their young. Shallow Gravel ecological sites do not provide the tall-statured annual weeds for thermal cover for deer, small mammals, and over-wintering birds. The response by wildlife species will be dependent upon

plant community composition, vegetative stature, patch size, and management activities (such as prescribed grazing, burning, inter-seeding, haying, or noxious weed control).

Successful restoration of native species along Transition Pathway R4A can result in a native grass and forb community in State 2.0. Over time, with no management, the exotic cool-season perennial grasses (Kentucky bluegrass, smooth brome, and/or quackgrass) generally become established and dominate the community. Failed native grass seeding or with grazing and time, this annual/pioneer perennial plant community will result in an invaded plant community within Invaded State 3.0.

Animal Community – Grazing Interpretations

This site is well adapted to managed grazing by domestic livestock. The predominance of herbaceous plants across all plant community phases best lends these sites to grazing by cattle, but other domestic grazers with differing diet preferences may also be a consideration depending upon management objectives. Often, the current plant community does not match any particular plant community (as described in the ecological site description). Because of this, a resource inventory is necessary to document plant composition and production. Proper interpretation of this inventory data will permit the establishment of a safe, initial stocking rate for the type and class of animals and level of grazing management. More accurate stocking rate estimates should eventually be calculated using actual stocking rate information and monitoring data.

NRCS defines prescribed grazing as “managing the harvest of vegetation with grazing and/or browsing animals with the intent to achieve specific ecological, economic, and management objectives”. As used in this site description, the term ‘prescribed grazing’ is intended to include multiple grazing management systems (e.g., rotational grazing, twice-over grazing, conservation grazing, targeted grazing, etc.) provided that, whatever management system is implemented, it meets the intent of prescribed grazing definition.

The basic grazing prescription addresses balancing forage demand (quality and quantity) with available forage, varying grazing and deferment periods from year-to-year, matching recovery/deferment periods to growing conditions when pastures are grazed more than once in a growing season, implementation of a contingency (e.g., drought) plan, and a monitoring plan. When the management goal is to facilitate change from one plant community phase or state to another, then the prescription needs to be designed to shift the competitive advantage to favor the native grass and forb species.

Grazing levels are noted within the plant community narratives and pathways in reference to grazing/prescribed grazing management. “Degree of utilization” is defined as the proportion of the current years forage production that is consumed and/or destroyed by grazing animals (may refer to a single plant species or a portion or all the vegetation). “Grazing utilization” is classified as slight, moderate, full, close, and severe (see the following table for description of each grazing use category). The following utilization levels are also described in the Ranchers Guide to Grassland Management IV. Utilization levels are determined by using the landscape appearance method as outlined in the Interagency Technical Reference “Utilization Studies and Residual Measurements” 1734-3.

Utilization Level % Use Description

Slight (Light) 0-20 Appears practically undisturbed when viewed obliquely. Only choice areas and forage utilized.

Moderate 20-40 Almost all of accessible range shows grazing. Little or no use of poor forage. Little evidence of trailing to grazing.

Full 40-60 All fully accessible areas are grazed. The major sites have key forage species properly utilized (about half taken, half left). Points of concentration with overuse limited to 5 to 10 percent of accessible area.

Close (Heavy) 60-80 All accessible range plainly shows use and major sections closely cropped. Livestock forced to use less desirable forage, considering seasonal preference.

Severe > 80 Key forage species completely used. Low-value forages are dominant.

Hydrological functions

Available water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group A. Infiltration varies from moderately rapid to very rapid; runoff potential varies from negligible to medium depending on surface texture, slope percent, slope shape, and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Dominance by blue grama and/or sedge will result in reduced infiltration and increased runoff. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

Recreational uses

Hunting and Bird Watching: National wildlife refuges, waterfowl production areas, state wildlife management areas (WMA), and North Dakota and South Dakota Department of Trust Lands provide herbaceous and woody cover for wildlife. National Wildlife Refuges and waterfowl production areas are owned and managed by the United States Fish and Wildlife Service and are available for public hunting, hiking, and bird watching. In addition, the United States Army Corps of Engineers and the United States Bureau of Reclamation manage three man-made reservoirs - Jamestown Reservoir (2,036 acres), Pipestem Reservoir (1,027 acres), and Lake Ashtabula (5,174 acres) for flood control, also providing fish habitat and adjacent uplands for wildlife cover. Lonetree WMA is the largest state managed wildlife area covering 32,800 acres. Numerous WMAs in North Dakota and Game Production Areas in South Dakota are found within this MLRA. The largest refuges managed by the United States Fish and Wildlife service are Arrowwood National Wildlife Refuge (NWR) Complex consists of 75,000 acres and Tewaukon National NWR covers 8,363 acres.

Fishing: Approximately 100 lakes are managed for public fishing within MLRA 55B. Most of these lakes offer boat docks and ramps. These lakes contain various sport fish including walleye, northern pike, yellow perch, catfish, trout, crappie, and bluegill. Many of these lakes are known for excellent round-around walleye and yellow perch fishery.

Camping: Fort Ramson State Park, Pipestem Reservoir, Jamestown Reservoir, Spiritwood Lake, Clausen Springs, Little Yellowstone, Richmond Lake State Recreation Area, Mina Lake State Recreation Area, and other public and private campgrounds are found within the MLRA. Limited, primitive camping is available on wildlife management areas. Ft. Ransom State Park (North Dakota), located along the Sheyenne River has a designated horse park with 15 miles of trails.

Hiking/Biking/Horseback Riding: Horseback riders, hikers, and biker can enjoy over 15 miles of multi-use trails at Fort Ransom State Park. The Jamestown Reservoir (5 miles), Pipestem Reservoir (8 miles) and Arrowwood National Wildlife Refuge (9.4 miles) maintain hiking trails. The Lonetree Wildlife Management Area has a 32- mile segment of the North Country Trail. It is designed for hiking and non-motorized travel including mountain bikes or horseback riding.

Canoeing/Kayaking: The Sheyenne River offers 278 miles of canoeing/kayaking from May-July. A kayak kiosk is located at Valley City and canoe/kayak rentals are available at Fort Ransom State Park. The James River has a canoe trail starting in Grand Rapids and canoeing down to the James River Dam site in LaMoure; no rentals are available.

Auto Tour: A 63-mile scenic drive starts north of Valley City and heading south through Sheyenne River Valley. Audubon National Wildlife Refuge offers a 5.5-mile auto-tour route winding through both prairie grassland and wetland habitats of the lower portion of the James River Valley.

Wood products

There are no significant wood products found on this site.

Other products

Seed harvest of native plant species can provide additional income on this site.

Other information

Site Development and Testing Plan

- Further investigation is needed where areas of this site occur on beach ridges. These areas are periodically inundated for extended periods. When the progressive soil surveys were completed, these soils were not inundated; currently many of them are inundated and have been for many consecutive years. A major component in this site affected by recent inundation is Wamduska. The MLRA map unit needing investigation is:

- Wamduska-Mauvais complex, 0 to 9 percent slopes (map unit 2q5cp)

- Further investigation is needed in areas of bedded shale to determine if a Very Shallow minor component should be added to data map units with Kloten soil as a major component. A few, brief, pedon descriptions and field notes in Soil Survey Office files indicate shale beds may occur within a depth of 10 inches in steep areas. If a Very Shallow component is added, the physiographic and soil features narratives in the ESD will need to be revised accordingly. MLRA map units to investigate are:

- Buse-Kloten-Edgeley complex, 9 to 40 percent slopes (map unit 2sd4f)

- Edgeley-Kloten-Buse loams, 9 to 35 percent slopes (map unit 2q5rw)

- Kloten-Walsh-Edgeley loams, 6 to 35 percent slopes (map unit 2q5rz)

- Further evaluation and refinement of the State-and-Transition model is needed to identify disturbance driven dynamics. Additional states and/or phases may be required to address grazing response.

- Further documentation may be needed for plant communities in all states. Plant data has been collected in previous range-site investigations, including clipping data; however, this data needs review. If geo-referenced sites meeting Tier 3 standards for either vegetative or soil data are not available, representative sites will be selected for further investigation.

- Site concepts will be refined as the above noted investigations are completed.

- The long-term goal is to complete an approved, correlated Ecological Site Description as defined by the National Ecological Site Handbook.

This ESD is the best available knowledge. The site concept and species composition table have been used in the field and tested for more than five years. It is expected that as additional information becomes available revisions may be required.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS; Jeff Printz, Range Management Specialist, NRCS; and Lee Voigt, Range Management Specialist, NRCS.

Other references

Bakker, K.K. 2003. The effect of woody vegetation on grassland nesting birds: an annotated bibliography. The Proceedings of the South Dakota Academy of Science 82:119-141.

Barker, W.T. and W. C. Whitman. 1988. Vegetation of the Northern Great Plains. Rangelands 10(6): 266-272.

Bluemle, J.P. 2016. North Dakota's geologic legacy. North Dakota State University Press. 382 pages.

Briske, D.D. (editor). 2017. Rangeland systems – processes, management, and challenges. Springer Series on Environmental Management. 661 pages.

Burgess, R.L. 1965. A study of plant succession in the sandhills of southeastern North Dakota. Proceedings ND Academy of Science 19:62-80

DeKeyser, E.S., G. Clambey, K. Krabbenhoft, and J. Ostendorf. 2009. Are changes in species composition on central North Dakota rangelands due to non-use management? Rangelands 31:16-19

Dix, R.L. and F.E. Smeins. 1967. The prairie, meadow, and marsh vegetation of Nelson County, North Dakota. Canadian Journal of Botany 45:21-57.

Dornbusch, M.J., R.F. Limb, and C.K. Gasch. 2018. Facilitation of an exotic grass through nitrogen enrichment by

an exotic legume. *Rangeland Ecology & Management* 71:691-694.

Dyke, S.R., S.K. Johnson, and P.T. Isakson. 2015. North Dakota state wildlife action plan. North Dakota Game and Fish Department, Bismarck, ND. 468 pages.

Ehrenfeld, Joan G. 2002. Effects of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* 6:503-523.

Endangered and threatened wildlife and plants; designation of critical habitat for the Dakota skipper and Poweshiek skipperling; Vol. 79 No. Final Rule October 1, 2015, 50 CFR Part 17.

Ereth, C., J. Hendrickson, D. Kirby, E. DeKeyser, K. Sedevic, and M. West. Controlling Kentucky bluegrass with herbicide and burning is influenced by invasion level. *Invasive Plant Science and Management* 10: 80-89.

Ewing, J. 1924. Plant succession on the brush prairie in northwestern Minnesota. *Journal of Ecology* 12:228- 266.

Gilbert, W.; and S. Zack. 2010. Integrating multiple ecosystem services introduction ecological site descriptions. *Rangelands*: 32:49-54.

Grant, T.A. and R.K. Murphy. 2005. Changes on woodland cover on prairie refuges in North Dakota, USA. *Natural Areas Journal* 25:359-368.

Heitschmidt, R. K., K. D. Klement, and M. R. Haferkamp. 2005. Interactive effects of drought and grazing on Northern Great Plains rangelands. *Rangeland Ecology and Management* 58:11-19.

Hendrickson, J.R., P. S. Johnson, M. A. Liebig, K. K. Sedivec, and G. A. Halvorson. 2016. Use of ecological sites in managing wildlife and livestock: an example with prairie dogs. *Rangelands*

Hendrickson, J.R., S.L. Kronberg, and E.J. Scholljegerdes. 2020. Can targeted grazing reduce abundance of invasive perennial grass (Kentucky bluegrass) on native mixed-grass prairie? *Rangeland Ecology and Management*, 73:547-551.

Higgins, K.F. 1984. Lightning fires in grasslands in North Dakota and in pine-savanna lands in nearby South Dakota and Montana. *J. Range Manage.* 37:100-103.

Higgins, K.F. 1986. Interpretation and compendium of historical fire accounts in the Northern Great Plains. United States Department of Interior, Fish and Wildlife Service. Resource Publication 161. 39 pages.

Higgins, K.F., A.D. Kruse, and J.L. Piehl. 1989. Effects of fire in the Northern Great Plains. U.S. Fish and Wildlife Service and Cooperative Extension Service South, Dakota State University. Extension Circular 761. 48 pages.

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://hprcc.unl.edu>)

Johnson, Sandra. 2015. Reptiles and amphibians of North Dakota. North Dakota Game and Fish Department. 64 pages.

Jordan, N. R., D.L. Larson, and S.C. Huerd. 2008. Soil modification by invasive plants: effects on native and invasive species of mixed-grass prairies. *Biological Invasions* 10:177-190.

Mader, E., M. Shepherd, M. Vaughan, and S.H. Black. 2011. Attracting native pollinators: protecting North America's bees and butterflies. Accessed at <https://xerces.org>, May 1, 2017.

North Dakota Division of Tourism, Accessed on February 25, 2019. Available at <https://www.ndtourism.com/sports-recreation>

North Dakota Parks and Recreation Department, Accessed on February 25, 2019. Available at <http://www.parkrec.nd.gov/recreationareas/recreationareas.html>

- Palit, R., G. and E.S. DeKeyser. 2022. Impacts and drivers of smooth brome (*Bromus inermis* L.) invasion in native ecosystems. *Plants*: 10,3390. <http://https://www.mdpi.com/2223-7747/11/10/1340>
- Palit, R., G. Gramig, and E.S. DeKeyser. 2021. Kentucky bluegrass invasion in the Northern Great Plains and prospective management approaches to mitigate its spread. *Plants*: 10,817. <https://doi.org/10.3390/plants10040817>
- Printz, J.L. and J.R. Hendrickson. 2015. Impacts of Kentucky bluegrass Invasion (*Poa pratensis*) on ecological processes in the Northern Great Plains. *Rangelands* 37(6):226-232.
- Redmann, Robert E. 1975. Production ecology of grassland plant communities in western North Dakota. *Ecological Monographs* 45:83-106.
- Reeves, J.L., J.D. Derner, M.A. Sanderson, J.R. Hendrickson, S.L. Kronberg, M.K. Petersen, and L.T. Vermeire. 2014. Seasonal weather influences on yearling beef steer production in C3-dominated Northern Great Plains rangeland. *Agriculture, Ecosystems and Environment* 183:110-117.
- Royer, R. A., 2003. Butterflies of North Dakota: an atlas and guide. Minot State University, Minot, ND.
- Seabloom, R. 2020. Mammals of North Dakota. North Dakota Institute for Regional Studies, Fargo, ND. 470 pages.
- Sedivec, K.D., J.L. Printz. 2014. Ranchers guide to grassland management IV. NDSU Extension Service publication R1707.
- Severson, K. E. and C. Hull Sieg. 2006. The nature of eastern North Dakota: Pre-1880 Historical Ecology. North Dakota Institute for Regional Studies.
- South Dakota Dept. of Game, Fish and Parks. 2014. South Dakota wildlife action plan. Wildlife Division Report 2014-03.
- Spaeth, K.E., Hayek, M.A., Toledo, D., and Hendrickson, J. 2019. Cool season grass impacts on native mixedgrass prairie species in the Northern Great Plains. America's Grassland Conference: Working Across Boundaries. The Fifth Biennial Conference on the Conservation of America's Grasslands. Bismarck, ND. 20- 22 August.
- Tidwell, D., D.T. Fogarty, and J.R. Weir. 2021. Woody encroachment in grasslands, a guide for understanding risk and vulnerability. Oklahoma State University, Oklahoma Cooperative Extension Service publication E- 1054. 32 pages.
- Toledo, D., M. Sanderson, K. Spaeth, J. Hendrickson, and J. Printz. 2014. Extent of Kentucky bluegrass and its effect on native plant species diversity and ecosystem services in the Northern Great Plains of the United State. *Invasive Plant Science and Management* 7(4): 543-552.
- USDA, NRCS. 2021. National Range and Pasture Handbook, (<https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/landuse/rangepasture/?cid=stelprdb1043084>)
- USDA, NRCS. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.
- USDA, NRCS. National Soil Information System, 100 Centennial Mall North, Room 152, Lincoln, NE 68508- 3866. (https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/tools/?cid=nrcs142p2_053552)
- USDA, NRCS. National Water & Climate Center, 1201 NE Lloyd Blvd, Suite 802, Portland, OR 97232-1274. (<https://www.wcc.nrcs.usda.gov/>)
- USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS, Various Published Soil Surveys.

USDI BLM.1999. Utilization studies and residual measurements. Interagency Technical Reference 1734-3.

Vinton, M.A. and E.M. Goergen. 2006. Plant-soil feedbacks contribute to the persistence of Bromus inermis in tallgrass prairie. Ecosystems 9: 967-976.

Whitman, W.H., H. Hanson, and R. Peterson. 1943. Relation of drought and grazing to North Dakota range lands. North Dakota Agricultural Experimentation Bulletin 340.

Contributors

Jeff Printz
David Dewald
Jonathan Fettig
Alan Gulsvig
Mark Hayek
Chuck Lura
Steve Sieler
Hal Weiser

Approval

Suzanne Mayne-Kinney, 4/25/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	USDA-NRCS North Dakota
Contact for lead author	NRCS State Rangeland Management Specialist
Date	04/25/2025
Approved by	Jeff Printz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills are not expected on this site when slopes are less than 15%. On slopes greater than 15%, scattered, short (12 to 20 inches), and disconnected rills may be visible.

2. **Presence of water flow patterns:** On slopes less than 15%, water flow patterns may be present but are infrequent and difficult to find. On slopes greater than 15%, water flow patterns are more pronounced, frequent, short (10 feet or less), disconnected and irregular in appearance.

3. **Number and height of erosional pedestals or terracettes:** Neither pedestals nor terracettes are expected when

slopes are less than 15%. Some pedestalling and terracettes may be evident when slopes exceed 15%.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground ranges from 20 to 35%. Bare ground patches consist of randomly scattered patches no greater than 10 inches in diameter and not connected. Animal activity (burrows and ant mounds) may occasionally result in isolated bare patches of up to 24 inches in diameter.

5. **Number of gullies and erosion associated with gullies:** Active gullies are not expected on this site. If present, gully channel(s) are fully vegetated with no active erosion visible.

6. **Extent of wind scoured, blowouts and/or depositional areas:** No wind-scoured or depositional areas expected on this site.

7. **Amount of litter movement (describe size and distance expected to travel):** Plant litter movement not expected on this site when slopes are less than 15%. When slopes exceed 15%, some short movement (12 to 24 inches) of small/fine class of litter (forb and grass leaves) may be present.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability class anticipated to average 5 or greater.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use soil series description for depth, color, and structure of A-horizon.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Mid- and short-statured bunchgrass are dominant and well distributed across the site. Forbs, grass-likes, and mid- and short-statured rhizomatous grasses are subdominant.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layers occur naturally on this site.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Phase 1.1:
Mid & short C3 bunch grasses (2); Mid & short C4 bunch grasses (4)

Sub-dominant: Phase 1.1:
Forbs (10); Grass-likes (1); Mid & short C3 rhizomatous grasses (1)

Other: Minor - Phase 1.1:
Shrub

Additional: Due to differences in phenology, root morphology, soil biology relationships, and nutrient cycling Kentucky bluegrass, smooth brome, and crested wheatgrass are included in a new Functional/structural group, mid- and short-statured early cool-season grasses (MSeC3), not expected for this site.

To see a full version 5 rangeland health worksheet with functional/structural group tables. Please use the following hyperlink:

https://efotg.sc.egov.usda.gov/references/public/ND/55B_Very_Shallow_Narrative_FINALRef_FSG.pdf

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead or dying plants/plant parts occur rarely on this site. Some dead centers may be observed on warm-season bunchgrasses after one to several years of below normal precipitation.

14. **Average percent litter cover (%) and depth (in):** Plant litter cover is 15 to 35% with a depth of 0.1 to 0.25 inches. Litter is in contact with soil surface.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual air-dry production is 1400 lbs./ac (reference value) with normal precipitation and temperatures. Low and high production years should yield 800 to 1900 lbs./ac, respectively.

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious species, Kentucky bluegrass, smooth brome, crested wheatgrass, and Eastern red cedar/juniper.

17. **Perennial plant reproductive capability:** Noninvasive species in all functional/structural groups are vigorous and capable of reproducing annually under normal weather conditions.
