

Ecological site R055CY001SD Shallow Marsh

Last updated: 1/31/2024 Accessed: 05/14/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 055C-Southern Black Glaciated Plains

The Southern Black Glaciated Plains (55C) is located within the Northern Great Plains Region. It is entirely within South Dakota encompassing about 10,835 square miles (Figure 1). The elevation ranges from 1,310 to 1,970 square feet. The MLRA is on nearly level to undulating glacial till plains interrupted by steeper slopes adjacent to streams and moraines. The James River is an under-fit stream. Its valley was carved by floodwaters draining glacial Lake Dakota and is filled with glacial outwash and alluvial deposits. (USDA-NRCS, 2006).

The dominant soil order in this MLRA is Mollisols. The soils in the area dominantly have a mesic soil temperature regime, an ustic soil moisture regime, and mixed or smectitic mineralogy. They generally are very deep, well drained to very poorly drained, and clayey or loamy. This area supports natural prairie vegetation characterized by western wheatgrass (Pascopyrum smithii), green needlegrass (Nassella viridula), needle and thread (Hesperostipa comata), and porcupinegrass (Hesperostipa spartea) with Prairie cordgrass (Spartina pectinata), and reed canarygrass (Phalaris arundinacea) as the dominant vegetation on the poorly drained soils. (USDA-NRCS, 2006).

Classification relationships

Major Land Resource Area (MLRA): Southern Black Glaciated Plains (55C) (USDA-NRCS, 2006)

USFS Subregions: North Central Glaciated Plains Section (251B); Yankton Hills and Valleys Subsection (251Bf); Western Glaciated Plains Section (332B); James River Lowland Subsection (332Bb); North Central Great Plains Section (332D); Southern Missouri Coteau Slope Subsection (332Dd); Southern Missouri Coteau Subsection (332De) - (Cleland et al., 2007).

US EPA Level IV Ecoregion: Southern Missouri Coteau (42e); Southern Missouri Coteau Slope (42f); James River Lowland (46n) - (USEPA, 2013)

Ecological site concept

The Shallow Marsh ecological site typically occurs in a basin or closed depression, and receives water directly from precipitation, surface overland flow, and groundwater discharge. Soils are formed in local alluvium and are very poorly drained, and have a water table within 1 foot of the soil surface. Permeability is very slow due to the clayey subsoil and the site will pond water until early summer in most years. Ponded water conditions and very slow permeability strongly influences the soil-water-plant relationship. Vegetation in the Reference State is typically dominated by cool-season grass and grass-like species including Whitetop, slough sedge, woolly sedge, American mannagrass, prairie cordgrass, and spikerush. Forb species may include smartweeds, western dock, and white panicle aster. Non-native species such as quackgrass, creeping meadow foxtail, and barnyardgrass may invade the site due to changes in disturbance regime.

Associated sites

R055CY003SD	Subirrigated These sites occur in drainageways. Soils are somewhat poorly drained with a water table within 2 to 5 feet of the soil surface that persists longer than the wettest part of the growing season, typically until the month of August. The central concept soil series is Crossplain, but other series are included.
R055CY004SD	Wet Meadow These sites occur in a basin or closed depression. Soils are poorly drained and the site ponds water for 4 to 8 weeks in the spring of the year or after a heavy rain. The central concept soil series is Tetonka, but other series are included.
R055CY020SD	Loamy Overflow These sites occur in upland swales. Soils are moderately well drained and have water flow into and over or through the site. The central concept soil series are Prosper, Bonilla, and Onita, but other series are included.
R055CY006SD	Limy Subirrigated These sites occur along the edges of drainageways. Soils are somewhat poorly drained and have a water table within 2 to 5 feet of the soil surface that persists longer than the wettest part of the growing season, typically until the month of August. Soils will effervesce with acid at or near the surface. The central concept soil series is Davison but other series are included.

Similar sites

Wet Meadow
Wet Meadow is similar in landscape position, but the site ponds water only for 4 to 8 weeks in the spring
of the year or after a heavy rain. A Wet Meadow will have more prairie cordgrass and lower production
than a Shallow Marsh site.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) Scolochloa festucacea(2) Carex atherodes

Physiographic features

This site occurs on concave shallow swales or depressions.



Figure 2. Distribution Map of the Shallow Marsh site within MLRA 55C. In many cases, data is not spatially consistent across political boundaries due to the method with which soils were mapped; e. g. county subsets.

Landforms	(1) Pothole
Flooding frequency	None
Ponding duration	Long (7 to 30 days) to very long (more than 30 days)
Ponding frequency	Frequent
Elevation	396–610 m
Slope	0–1%
Ponding depth	0–61 cm
Water table depth	0–46 cm
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

MLRA 55C is considered to have a continental climate: Cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the Northern Great Plains, and air masses move freely across the plains and account for rapid changes in temperature.

Annual precipitation typically ranges from 19 to 25 inches per year. The average annual temperature is about 47°F. January is the coldest month with average temperatures ranging from about 15°F (Howard, South Dakota [SD]), to about 20°F (Wagner, SD). July is the warmest month with temperatures averaging from about 73°F (Howard, SD), to about 77°F (Wagner, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 58°F. This large annual range attests to the continental nature of this area's climate. Hourly winds are estimated to average about 12 miles per hour (mph) annually, ranging from about 13 mph during the spring to about 11 mph during the summer. Daytime winds are generally stronger than nighttime, and occasional strong storms may bring brief periods of high winds with gusts to more than 50 mph.

Growth of cool-season plants begins in early to mid-March, slowing or ceasing in late June. Warm-season plants begin growth about mid-May and continue to early or mid-September. Green-up of cool-season plants may occur in September and October when adequate soil moisture is present.

Table 3. Representative climatic features	Table	3.	Representative	climatic	features
---	-------	----	----------------	----------	----------

Frost-free period (characteristic range)	122-128 days
Freeze-free period (characteristic range)	135-147 days

Precipitation total (characteristic range)	559-686 mm		
Frost-free period (actual range)	114-130 days		
Freeze-free period (actual range)	133-151 days		
Precipitation total (actual range)	559-686 mm		
Frost-free period (average)	124 days		
Freeze-free period (average)	142 days		
Precipitation total (average)	635 mm		

Climate stations used

- (1) ACADEMY 2NE [USC00390043], Platte, SD
- (2) WAGNER [USC00398767], Wagner, SD
- (3) TYNDALL [USC00398472], Tyndall, SD
- (4) MENNO [USC00395481], Menno, SD
- (5) MARION [USC00395228], Marion, SD
- (6) ALEXANDRIA [USC00390128], Alexandria, SD
- (7) HOWARD [USC00394037], Howard, SD
- (8) HURON RGNL AP [USW00014936], Huron, SD
- (9) MILLER [USC00395561], Miller, SD
- (10) FAULKTON 1 NW [USC00392927], Faulkton, SD
- (11) REDFIELD [USC00397052], Redfield, SD
- (12) BRIDGEWATER [USC00391032], Bridgewater, SD
- (13) CHAMBERLAIN 5 S [USC00391621], Chamberlain, SD
- (14) SALEM 5NE [USC00395360], Salem, SD

Influencing water features

This section is under construction.

Wetland description

This section is under construction.

Soil features

The common features of soils in in the Shallow Marsh ES are the clay loam to clay-textured subsoil and slopes of 0 to 1 percent. The soils in this site are poorly to very poorly drained and formed in alluvium. The silt loam to silty clay loam surface layer is 8 to 20 inches thick. The soils have a very slow infiltration rate. This site should show no evidence of rills, wind-scoured areas, or pedestalled plants. The soil surface is stable and intact. Subsurface soil layers are nonrestrictive to water movement and root penetration.

These soils are not susceptible to the hazard of water erosion. Ponded water conditions and very slow permeability strongly influence the soil-water-plant relationship. The primary series associated with the Shallow Marsh site is Worthing.

Access Web Soil Survey (http://websoilsurvey.nrcs.usda.gov/app/HomePage.htm) for specific local soils information.

Table 4. Representative soil features

Surface texture	(1) Silty clay loam(2) Silt loam(3) Fine sandy loam		
Family particle size	(1) Clayey		

Drainage class	Poorly drained
Permeability class	Very slow
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	15.24–17.78 cm
Calcium carbonate equivalent (0-101.6cm)	0–5%
Electrical conductivity (0-101.6cm)	1–8 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	5.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–3%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

State and Community Phases

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

The Shallow Marsh ecological site, located in the Southern Black Glaciated Plains Region of the MLRA, developed under Northern Great Plains climatic conditions and included historic natural influence of large herding herbivores and occasional fire. Changes occur in the plant communities due to weather fluctuations and management actions. Under adverse impacts, a relatively rapid decline in vegetative vigor and composition can occur. Under favorable conditions the site has the potential to resemble the Reference State. Interpretations for this site are based primarily on the 1.1 Whitetop-Slough Sedge Plant Community Phase. This community phase and the Reference State have been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been considered.

This ecological site (ES) has been grazed by domestic livestock since they have been introduced into the area. The introduction of domestic livestock and the use of fencing and reliable water sources have changed the ecological dynamics of this site. Heavy, continuous grazing without adequate recovery periods following each grazing occurrence causes this site to depart from the Reference State due to the compaction and overgrazing. Species such as fowl bluegrass (*Poa palustris*), spikerush (Eleocharis), and Baltic rush (Juncus balticus) will initially increase. Whitetop and slough sedge will decrease in frequency and production. Continued heavy grazing eventually causes a dominance by spikerush, rushes (Juncus), and unpalatable forbs such as curly dock (*Rumex crispus*).

Following the state-and-transition diagram are narratives for each of the described states and community phases. These may not represent every possibility, but they are the most prevalent and repeatable states and community phases. The associated plant composition tables have been developed from the best available knowledge at the time of this revision. As more data are collected, some of these community phases and states may be revised or removed, and new ones may be added. The main purpose for including the descriptions here is to capture the current knowledge and experience at the time of this revision.

The following is a diagram that illustrates the common plant community phases that can occur on the site and the transition and community pathways between them. The ecological processes will be discussed in more detail in the plant community descriptions following the diagram.

State and transition model



Figure 9. State-And-Transition Model for the Shallow Marsh site in MLRA 55C.

Code	Process					
T1A	Heavy continuous grazing, inundation, no fire					
T1B	Tillage, artificial drainage (surface and subsurface)					
T2A	Long term prescribed grazing, wetland restoration					
T2B	Tillage, artificial drainage (surface and subsurface)					
T3A	Abandonment of cropping					
1.1A	Heavy continuous grazing					
1.2A	Prescribed grazing with recovery periods					
2.1A	Heavy continuous grazing, below normal precipitation period, no fire					
2.2A	Inundation					

Figure 10. The Matrix for the Shallow Marsh site in MLRA 55C.

State 1 **Reference State**

The Reference State represents the natural range of variability that dominates the dynamics of this ES. This state is typically dominated by cool-season grass and grass-like species. Before European settlement, the primary disturbance mechanisms for this site in the reference condition included periodic fire, grazing by large herding ungulates, and fluctuations in the water table and ponding frequency and duration. Frequent surface fires (3 to 5 years) and grazing coupled with weather events dictated the dynamics that occurred within the natural range of variability. Today, the primary disturbance is from a lack of fire, concentrated livestock grazing, and weather fluctuations. Species that are desirable for livestock and wildlife can decline and a corresponding increase in less desirable species will occur.

Community 1.1 Whitetop/Slough Sedge

Interpretations are based primarily on the 1.1 Whitetop/Slough Sedge Plant Community Phase (this is also considered to be the Reference Community). This plant community evolved with grazing by large herbivores, infrequent wild fires, and periodic flooding events, and is suited for grazing by domestic livestock. This plant community can be found on areas that are grazed and where the grazed plants receive adequate periods of rest during the growing season in order to recover. Historically, fires occurred infrequently. The potential vegetation is about 45 percent grasses, 40 percent grass-likes, and 15 percent forbs. The major grasses and grass-likes include whitetop (also known as common rivergrass), slough sedge (also known as wheat sedge), woolly sedge, American mannagrass, prairie cordgrass, Sartwell's sedge, Nebraska sedge, and spikerush. Key forbs include smartweeds, western dock, and white panicle aster. This plant community phase is diverse, stable, and productive, and is well adapted to the Northern Great Plains. The high water table supplies much of the moisture for plant growth. Community dynamics, nutrient and water cycles, and energy flow are functioning properly. Plant litter is properly distributed with very little movement offsite and natural plant mortality is very low. The diversity in plant species allows for the variability of both the fluctuations of the water table and reoccurring ponding. This is a sustainable plant community in terms of soil stability, watershed function, and biologic integrity.

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)				
Grass/Grasslike	5750	6456	7162				
Forb	303	717	1132				
Total	6053	7173	8294				

Table 5. Annual production by plant type

Figure 12. Plant community growth curve (percent production by month). SD5506, Southern Black Glaciated Plains, lowland cool-season dominant.. Cool-season dominant, lowland ..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	6	15	20	26	17	9	4	3	0	0

Community 1.2 Slough Sedge/Spikerush

This plant community will slowly develop from the adverse effects of continuous grazing without adequate recovery periods between each grazing event during the growing season. When compared to the 1.1 Whitetop/Slough Sedge Plant Community Phase, whitetop, prairie cordgrass, American mannagrass, and reedgrasses have decreased. The grass-like species, such as slough sedge, spikerush, woolly sedge, Sartwell's sedge, Nebraska sedge, and rushes have increased and tend to dominate this plant community.

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	4780	5394	6210
Forb	263	771	1076
Total	5043	6165	7286

Figure 14. Plant community growth curve (percent production by month). SD5506, Southern Black Glaciated Plains, lowland cool-season dominant.. Cool-season dominant, lowland..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	6	15	20	26	17	9	4	3	0	0

Pathway 1.1A Community 1.1 to 1.2

Heavy, continuous grazing that includes herbivory at moderate to heavy levels at the same time of year, each year without adequate recovery periods or during periods of below-average precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites, will shift this community to the 1.2 Slough Sedge/Spikerush Plant Community Phase.

Pathway 1.2A Community 1.2 to 1.1

Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing, possibly including periodic rest, will convert this plant community to the 1.1 Whitetop/Slough Sedge Plant Community Phase. This pathway could also occur with a return to more normal precipitation levels and frequencies.

Conservation practices

Prescribed Grazing

State 2 Invaded State

The Invaded State is characterized by the increase in bare ground due to trampling caused by excessive use or by inundation for extended periods, which causes a temporary shift in the plant composition and cover. This allows for the invasion of nonnative species which with continued heavy grazing can increase to eventual dominance. Loss or reduction of native cool- and warm-season species can negatively impact energy flow and nutrient cycling. Infiltration will be reduced and native plant mortality will increase. As the disturbance level increases, native plant density decreases even more, giving way to annual species and invasive perennial species, and a further increase

Community 2.1 Spikerush/Baltic Rush/Fowl Bluegrass/Forbs

This plant community developed with heavy, continuous grazing without adequate recovery periods between grazing events or inundation during periods of extended above-average precipitation. Spikerush, Baltic rush, bulrush, and other less desirable grass-likes, along with grasses such as fowl bluegrass and American sloughgrass, dominate the community. Quackgrass, creeping meadow foxtail, Kentucky bluegrass, and other nonnative species can invade on drier portions of the community. Whitetop, slough sedge, other sedges, prairie cordgrass, and reedgrasses will be virtually eliminated from the plant community. Smartweeds, dock, and cinquefoil have increased. Areas of bare ground can be present throughout the site. A significant amount of production and diversity has been lost when compared to the 1.1 Whitetop/Slough Sedge Plant Community Phase. Loss or reduction of native grasses, grass-likes, and forbs has negatively impacted energy flow and nutrient cycling. It will take a long time to restore this plant community with improved management or return of more normal precipitation patterns.

Table 7. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	2163	2869	3576
Forb	303	717	1132
Total	2466	3586	4708

Figure 16. Plant community growth curve (percent production by month). SD5506, Southern Black Glaciated Plains, lowland cool-season dominant.. Cool-season dominant, lowland..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	6	15	20	26	17	9	4	3	0	0

Community 2.2 Annual/Pioneer, Non-native

This plant community develops under severe disturbance, typically abandonment after cropping. The dominant vegetation includes pioneer annual or perennial grasses, forbs, invaders, and early successional biennial and perennial species. Grasses may include foxtail barley, barnyardgrass, quackgrass, fowl bluegrass, Kentucky bluegrass, Baltic rush, and sedges. The dominant forbs include knotweed, Canada thistle, and other early successional species. The community is susceptible to invasion of nonnative species due to severe soil disturbances and relatively high percentage of bare ground. This plant community is resistant to change, as long as soil disturbance or severe vegetation defoliation persists, thus holding back secondary plant succession. Significant economic inputs, management, and time would be required to move this plant community toward a higher successional stage. Secondary successional species. This plant community may be renovated to improve the production capability but management changes would be needed to maintain the new plant community. The total annual production ranges from 1,000 to 1,500 lbs./ac. (air-dry weight), depending upon growing conditions. No growth curve has been assigned to this plant community phase due to the highly variable nature of the plant community.

Pathway 2.1A Community 2.1 to 2.2

Heavy, continuous grazing which includes herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or during periods of below-normal precipitation when grazing frequency and intensity increases on these sites due to limited forage availability on adjacent upland sites and no surface fire for extended periods of time (typically for 10 years or more) causing litter levels to become high enough to reduce native grass vigor, diversity, and density will shift this community to the 2.2 Annual Pioneer-Perennial Pioneer Plant Community Phase.

Pathway 2.2A Community 2.2 to 2.1

Inundation for extended periods beyond normal ponding and drying patterns will convert this plant community to the 2.1 Spikerush-Baltic Rush-Fowl Bluegrass-Forbs Plant Community Phase within the Invaded State (State 2).

State 3 Crop Production State

This state is characterized by the production of annual crops using a variety of tillage and cropping systems along with management practices. Cropping on this site is enabled during years with drier than normal precipitation or with artificial drainage (surface or subsurface).

Community 3.1 Annual Crops

This plant community developed with the use of a variety of tillage and cropping systems for the production of annual crops including corn, soybeans, wheat, and a variety of other crops.

Transition T1A State 1 to 2

Heavy, continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season and often at the same time of year each year), no surface fire for extended periods of time (typically for 10 years or more) causing litter levels to become high enough to reduce native grass vigor, diversity, and density, or inundation for extended periods beyond normal ponding and drying patterns will eventually cause a shift over a threshold leading to the 2.1 Spikerush-Baltic Rush-Fowl Bluegrass-Forbs Plant Community Phase within the Invaded State (State 2). Grazing repeatedly in the early growing season can expedite this shift by causing mechanical disturbance due to trampling.

Transition T1B State 1 to 3

Tillage and artificial drainage (surface and subsurface) will cause a shift over a threshold leading to the 3.1 Annual Crops Plant Community Phase within the Crop Production State (State 3).

Restoration pathway R2A State 2 to 1

Long-term prescribed grazing (moderate stocking levels coupled with adequate recovery periods, or other grazing systems such as high-density, low-frequency intended to treat specific species dominance, or periodic light to moderate stocking levels possibly including periodic rest) may lead this plant community phase over a threshold to the Reference State (State 1). Wetland restoration techniques may be necessary to restore biotic integrity and plant diversity and productivity.

Conservation practices

Prescribed Grazing

Transition T2B State 2 to 3

Tillage and artificial drainage (surface and subsurface) will cause a shift over a threshold leading to the 3.1 Annual Crops Plant Community Phase within the Crop Production State (State 3).

Restoration pathway T3A State 3 to 2

Cropping followed by abandonment may lead this plant community phase over a threshold to the Invaded State (State 2) and more specifically to the 2.2 Annual Pioneer-Perennial Pioneer Plant Community Phase.

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike	•			
1	Tall Cool-Season Grasses			1435–2869	
	common rivergrass	SCFE	Scolochloa festucacea	1076–2511	_
	American mannagrass	GLGR	Glyceria grandis	359–717	_
2	Grass-Likes			1435–2869	
	wheat sedge	CAAT2	Carex atherodes	717–2152	_
	Nebraska sedge	CANE2	Carex nebrascensis	72–717	_
	woolly sedge	CAPE42	Carex pellita	143–717	_
	Sartwell's sedge	CASA8	Carex sartwellii	143–717	_
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–717	_
	spikerush	ELEOC	Eleocharis	143–717	_
	mountain rush	JUARL	Juncus arcticus ssp. littoralis	72–574	_
	green bulrush	SCAT2	Scirpus atrovirens	72–574	_
	rush	JUNCU	Juncus	0–359	_
	flatsedge	CYPER	Cyperus	0–359	_
3	Tall warm-season grasses	•		143–717	
	prairie cordgrass	SPPE	Spartina pectinata	143–717	_
	spiked muhly	MUGL3	Muhlenbergia glomerata	0–359	_
4	Reedgrasses			0–359	
	bluejoint	CACA4	Calamagrostis canadensis	0–359	_
	northern reedgrass	CASTI3	Calamagrostis stricta ssp. inexpansa	0–359	_
	slimstem reedgrass	CASTS5	Calamagrostis stricta ssp. stricta	0–359	_
5	Other native grasses			359–717	
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass-like)	143–574	_
	American sloughgrass	BESY	Beckmannia syzigachne	143–574	_
Forb					
7	Forbs			359–1076	
	Forb, native	2FN	Forb, native	72–287	_
	bur-reed	SPARG	Sparganium	72–287	_
	New England aster	SYNO2	Symphyotrichum novae-angliae	72–143	_
	pale dock	RUAL4	Rumex altissimus	0–143	_
	western dock	RUAQ	Rumex aquaticus	72–143	_
	Canadian anemone	ANCA8	Anemone canadensis	72–143	_
	Indianhemp	APCA	Apocynum cannabinum	0–143	_

Flodman's thistle	CIFL	Cirsium flodmanii	72–143	-
Rydberg's sunflower	HENUR	Helianthus nuttallii ssp. rydbergii	0–143	_
knotweed	POLYG4	Polygonum	72–143	_
Pennsylvania smartweed	POPE2	Polygonum pensylvanicum	72–143	-
giant goldenrod	SOGI	Solidago gigantea	72–143	-
broadleaf cattail	TYLA	Typha latifolia	0–143	-
cinquefoil	POTEN	Potentilla	0–72	-
Macoun's buttercup	RAMA2	Ranunculus macounii	0–72	-
smooth horsetail	EQLA	Equisetum laevigatum	0–72	-
hemlock waterparsnip	SISU2	Sium suave	0–72	-
blue-eyed grass	SISYR	Sisyrinchium	0–72	-
marsh arrowgrass	TRPA28	Triglochin palustris	0–72	-
white panicle aster	SYLA6	Symphyotrichum lanceolatum	0–72	_

Table 9. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike				
1	Tall Cool-Season Grasses			0–493	
	common rivergrass	SCFE	Scolochloa festucacea	0–493	_
	American mannagrass	GLGR	Glyceria grandis	0–308	_
2	Grass-Likes			2466–4624	
	wheat sedge	CAAT2	Carex atherodes	925–2466	_
	spikerush	ELEOC	Eleocharis	616–1849	_
	woolly sedge	CAPE42	Carex pellita	308–925	_
	Sartwell's sedge	CASA8	Carex sartwellii	0–616	_
	flatsedge	CYPER	Cyperus	0–616	_
	Nebraska sedge	CANE2	Carex nebrascensis	0–616	_
	rush	JUNCU	Juncus	0–616	_
	green bulrush	SCAT2	Scirpus atrovirens	0–616	_
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–493	_
4	Reed Grasses			0–123	
	northern reedgrass	CASTI3	Calamagrostis stricta ssp. inexpansa	0–123	_
	slimstem reedgrass	CASTS5	Calamagrostis stricta ssp. stricta	0–123	_
5	Other Native Grasses	•		0–308	
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass-like)	0–308	_
	American sloughgrass	BESY	Beckmannia syzigachne	0–308	_
6	Non-native Grasses			62–308	
	barnyardgrass	ECCR	Echinochloa crus-galli	62–308	_
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass-like)	0–185	_
Shrub	/Vine				
3	Tall Warm-Season Grasses			0–308	
	,	~~~~		0,000	

_

Forb					
7	Forbs			308–1233	
	broadleaf cattail	TYLA	Typha latifolia	62–247	-
	Forb, introduced	2FI	Forb, introduced	0–247	-
	Forb, native	2FN	Forb, native	62–247	-
	Pennsylvania smartweed	POPE2	Polygonum pensylvanicum	62–247	-
	giant goldenrod	SOGI	Solidago gigantea	62–185	_
	bur-reed	SPARG	Sparganium	0–185	_
	curly dock	RUCR	Rumex crispus	62–185	_
	knotweed	POLYG4	Polygonum	62–185	_
	narrowleaf cattail	TYAN	Typha angustifolia	62–185	_
	hemlock waterparsnip	SISU2	Sium suave	0–123	_
	white panicle aster	SYLA6	Symphyotrichum lanceolatum	0–123	_
	New England aster	SYNO2	Symphyotrichum novae-angliae	0–123	_
	pale dock	RUAL4	Rumex altissimus	0–123	_
	Indianhemp	APCA	Apocynum cannabinum	0–123	_
	Flodman's thistle	CIFL	Cirsium flodmanii	0–123	_
	splitlip hempnettle	GABI3	Galeopsis bifida	0–123	_
	Rydberg's sunflower	HENUR	Helianthus nuttallii ssp. rydbergii	0–62	_
	smooth horsetail	EQLA	Equisetum laevigatum	0–62	_
	northern water plantain	ALTR7	Alisma triviale	0–62	_
	Canadian anemone	ANCA8	Anemone canadensis	0–62	-
	western dock	RUAQ	Rumex aquaticus	0–62	-
	cinquefoil	POTEN	Potentilla	0–62	_

Table 10. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike	•	•		
2	Grass-Likes			897–1973	
	spikerush	ELEOC	Eleocharis	359–1076	_
	wheat sedge	CAAT2	Carex atherodes	179–538	_
	rush	JUNCU	Juncus	0–359	
	woolly sedge	CAPE42	Carex pellita	0–287	
	flatsedge	CYPER	Cyperus	0–287	
	Grass-like (not a true grass)	2GL	Grass-like (not a true grass)	0–179	
	Sartwell's sedge	CASA8	Carex sartwellii	0–179	
	Nebraska sedge	CANE2	Carex nebrascensis	0–179	
	green bulrush	SCAT2	Scirpus atrovirens	0–108	
5	Other Native Grasses	4		0–359	
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass- like)	0–359	
	American sloughgrass	BESY	Beckmannia syzigachne	0–359	
6	Non-native Grasses	4		72–538	I
	creeping meadow foxtail	ALAR	Alopecurus arundinaceus	0–359	
	barnyardgrass	ECCR	Echinochloa crus-galli	36–287	_
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass- like)	0–287	
	quackgrass	ELRE4	Elymus repens	0–215	
Forb	•	4	•		
7	Forbs			359–1076	
	giant goldenrod	SOGI	Solidago gigantea	36–363	_
	curly dock	RUCR	Rumex crispus	72–359	_
	Forb, introduced	2FI	Forb, introduced	36–359	_
	Pennsylvania smartweed	POPE2	Polygonum pensylvanicum	36–287	_
	narrowleaf cattail	TYAN	Typha angustifolia	36–287	_
	broadleaf cattail	TYLA	Typha latifolia	36–287	_
	knotweed	POLYG4	Polygonum	36–215	_
	Forb, native	2FN	Forb, native	36–215	_
	white panicle aster	SYLA6	Symphyotrichum lanceolatum	0–143	_
	New England aster	SYNO2	Symphyotrichum novae- angliae	0–143	_
	hemlock waterparsnip	SISU2	Sium suave	0–108	_
	Indianhemp	APCA	Apocynum cannabinum	0–108	_
	smooth horsetail	EQLA	Equisetum laevigatum	0–72	_
	cinquefoil	POTEN	Potentilla	0–72	
	bur-reed	SPARG	Sparganium	0–72	
	pale dock	RUAL4	Rumex altissimus	0–36	_
	Flodman's thistle	CIFL	Cirsium flodmanii	0–36	_

Animal community

Animal Community – Grazing Interpretations

The following table lists annual, suggested initial stocking rates for average growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of conservation planning. Often, the current plant composition does not entirely match any particular plant community (as described in this ES description). Because of this, a resource inventory is necessary to document plant composition and production. More accurate carrying capacity estimates should eventually be calculated using the following stocking rate information, along with animal preference data and actual stocking records, particularly when grazers other than cattle are involved. After consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity. Stocking rates are calculated using Animal-Unit-Month (AUM), which is the amount of air-dry forage required to feed a cow, with or without calf, for one month.

Prairie Cordgrass/Sedge/Northern Reedgrass (1.1) Average Annual Production (lbs./acre, air-dry): 6,400 Stocking rate (AUM/acre)*: 1.75

Fowl Bluegrass/Spikerush/Baltic Rush/Forbs (1.2) Average Annual Production (lbs./acre, air-dry): 5,500 Stocking rate (AUM/acre)*: 1.51

Reed Canarygrass/Sedge/Invasive Plants/Prairie Cordgrass (2.1) Average Annual Production (lbs./acre, air-dry): 3,200 Stocking rate (AUM/acre)*: 0.88

Annual/Pioneer, Non-Native Perennial (2.2) Average Annual Production (lbs./acre, air-dry): 1,500 Stocking rate (AUM/acre)*: 0.41

*Based on 912 lbs./acre (air-dry weight) per Animal Unit Month (AUM) and on 25 percent harvest efficiency (refer to United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS), National Range and Pasture Handbook).

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage. During the dormant period, the forage for livestock will likely be lacking protein to meet livestock requirements and added protein will allow ruminants to better utilize the energy stored in grazed plant materials. A forage quality test (either directly or through fecal sampling) should be used to determine the level of supplementation needed.

Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group D. Infiltration is very slow and runoff potential for this site is negligible. In many cases, areas with greater than 75 percent ground cover have the greatest potential for higher infiltration. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration. Runoff is negligible on this site (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

Recreational uses

This site provides hunting, hiking, photography, bird watching, and other opportunities. The wide varieties of plants that bloom from spring until fall have an aesthetic value that appeals to visitors.

Wood products

No appreciable wood products are typically present on this site.

Other products

Seed harvest of native plant species can provide additional income on this site.

Other information

Ecological Site Correlation Issues and Questions:

• SD069 Hyde County, SD did not use the (Wo) Worthing, sicl, 0 to 1 percent slopes (national symbol 2tlch) as used in the adjoining SD059 Hand County.

• SD005 Beadle County, SD did not use the (Wo) Worthing, sicl, 0 to 1 percent slopes (national symbol 2tlch) as used in the adjoining counties.

• The MLRA 55C – Worthing, sicl, 0 to 1 percent slopes (national symbol 2tlch) was also used in MLRA 53C and MLRA 102B. A future project is needed to assign the correct ESD in each MLRA.

• Reference and alternative states within the state and transition model are may not be fully documented and may require additional field sampling for refinement.

Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS; and Bruce Kunze, Soil Scientist, NRCS.

There are two SCS-RANGE-417s collected in 2005 from Beadle County, SD.

Other references

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Coterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC.

Gilbert, M. C., Whited, P. M., Clairain Jr, E. J., & Smith, R. D. (2006). A Regional Guidebook for Applying the Hydrogeomorphic Approach to Assessing Wetland Functions of Prairie Potholes. Washington DC.

Samson, F. B., & Knopf, F. L. (1996). Prairie Conservation Preserving North America's Most Endagered Ecosystem. Washington D.C.: Island Press.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions. Available online. Accessed March 2018.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2003. National Range and Pasture Handbook, Revision 1. Grazing Lands Technology Institute.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

USDA, NRCS. National Soil Information System, Information Technology Center, (http://soils.usda.gov/technical/nasis/)

USDA, NRCS. 2019. The PLANTS Database (http://plants.usda.gov, 13 March 2019).

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at http://www.epa.gov/eco-research/level-iii-and-iv-ecoregions- continental-united-states. (Accessed 13 March 2019).

High Plains Regional Climate Center, University of Nebraska. (http://www.hprcc.unl.edu/)

Contributors

Kenzie Cobb Stan Boltz

Approval

Suzanne Mayne-Kinney, 1/31/2024

Acknowledgments

Natural Resources Conservation Service (USDA-NRCS), Redfield Soil Survey Office Redfield, SD; Lance Howe (Lance.Howe@usda.gov), Soil Survey Office Leader, USDA-NRCS, Redfield, SD; and Steve Winter (Steven.Winter@usda.gov), Soil Scientist, USDA-NRCS, Redfield, SD.

Additional Information Acknowledgment: Jason Hermann (Jason.Hermann@usda.gov), Area Rangeland Management Specialist, USDA-NRCS, Redfield, SD.

This Provisional Ecological Site concept has passed both Quality Control and Quality Assurance processes. Officially approved by David Kraft as of 11/12/2020.

Non-discrimination Statement

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, available online and at any USDA office, or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632- 9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	David Schmidt, Tim Nordquist, Stan Boltz
Contact for lead author	david.schmidt@sd.usda.gov 605-352-1236

Date	12/07/2004
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills: Rills should not be present.
- 2. Presence of water flow patterns: Barely observable.
- 3. Number and height of erosional pedestals or terracettes: Essentially, non-existent.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground less than five percent and less than two inches in diameter.
- 5. Number of gullies and erosion associated with gullies: Active gullies should not be present.
- 6. Extent of wind scoured, blowouts and/or depositional areas: None.
- 7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Stability class 6. Typically high root content, and organic matter. Soil surface is very resistant to erosion.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use soil series description for depth and color of A-horizon.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Healthy, deep rooted native grasses enhance infiltration and reduce runoff.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer should be evident.

foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Tall cool-season rhizomatous grass >

Sub-dominant: Mid and tall cool-season rhizomatous grass-like >

Other: Tall warm-season rhizomatous grass > forb.

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very little to no evidence of decadence or mortality.

14. Average percent litter cover (%) and depth (in): Litter cover is in contact with soil surface.

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): 6,400–7,400 lbs./acre air-dry weight, average 6,400 lbs./acre air-dry weight.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Refer to State and Local Noxious Weed List, also reed canarygrass.
- 17. Perennial plant reproductive capability: All species are capable of reproducing.