

Ecological site R055CY008SD Sands

Last updated: 1/31/2024 Accessed: 05/03/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| Author(s)/participant(s) | David Schmidt, Tim Nordquist, Stan Boltz |
|---|--|
| Contact for lead author | david.schmidt@sd.usda.gov 605-352-1236 |
| Date | 12/07/2004 |
| Approved by | Stan Boltz |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

| 1110 | nuicators | | |
|------|--|--|--|
| 1. | Number and extent of rills: Rills should not be present. | | |
| 2. | Presence of water flow patterns: Barely observable. | | |
| 3. | Number and height of erosional pedestals or terracettes: Slight pedestalling on bunch grasses. | | |
| 4. | Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground less than 10 percent and less than 2 inches in diameter. | | |
| 5. | Number of gullies and erosion associated with gullies: Active gullies should not be present. | | |
| 6. | Extent of wind scoured, blowouts and/or depositional areas: None. | | |
| | | | |

7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant

litter remains in place and is not moved by erosional forces.

| 8. | Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Stability class usually 4-6. Typical high root content protects soil surface from erosion. |
|-----|--|
| 9. | Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Use soil series description for depth and color of A-horizon. |
| 10. | Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Healthy, deep rooted native grasses enhance infiltration and reduce runoff. |
| 11. | Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer should be evident. |
| 12. | Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to): |
| | Dominant: Tall warm-season rhizomatous grass >> mid warm-season bunch grass > |
| | Sub-dominant: Mid warm-season bunchgrasses > mid and tall cool-season bunch grass > |
| | Other: Short warm-season grasses = forb = shrub. |
| | Additional: Due to differing root structure and distribution, Kentucky bluegrass and smooth bromegrass do not fit into reference plant community F/S groups. |
| 13. | Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very little to no evidence of decadence or mortality. |
| 14. | Average percent litter cover (%) and depth (in): Litter cover is in contact with soil surface. |
| 15. | Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): 2,000–3,600 lbs./acre air-dry weight, average 2,800 lbs./acre air-dry weight. |
| 16. | Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Refer to State and Local Noxious Weed List. |

| 17. | Perennial plant reproductive capability: All species are capable of reproducing. |
|-----|--|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |