

Ecological site R058AC618MT Saline Overflow (SOv) RRU 58A-C 11-14" p.z.

Last updated: 6/14/2023
Accessed: 04/25/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Matt Ricketts, Tony Rolfes, Loretta Metz
Contact for lead author	
Date	04/11/2005
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** No rills present in the reference state.

- Presence of water flow patterns:** Due to the soil surface being well covered and minimal slope there is no evidence of past or current soil deposition or erosion for this site.

- Number and height of erosional pedestals or terracettes:** These should not be evident in the reference state.

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is less than 20% in the reference state.

- Number of gullies and erosion associated with gullies:** Gully erosion may be evident in the reference state, but only following storms of greater intensity than "normal".

- Extent of wind scoured, blowouts and/or depositional areas:** These are not present in the reference state.

7. **Amount of litter movement (describe size and distance expected to travel):** Because there is little bare ground, litter movement will be minimal at most. Because the site is dominated by the taller bunchgrasses and rhizomatous grasses, litter size will reflect the height and diameter of the reproductive culms and leaves of these grasses as well as the lesser dominate mid-size grasses.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values of 4 to 5 under plant canopies, and 2-3 in the plant interspaces.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface structure is granular. Organic matter is 2-4%. The A-horizon is 4 to 8 inches thick.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Deep-rooted native warm season perennial grasses with up to 15% woody species optimize infiltration and runoff.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer present in the reference state.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm season, mid grasses > warm season, short grasses = tall shrubs > mid shrubs > cool season, short grasses > forbs.
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Plant mortality is very low; decadence is minimal except in prolonged periods of drought (>5-6 years).
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1700 – 2000 #/acre. This would be the expected production for the reference state during average moisture years. 1800 pounds would be the expected production in a 12 inch average precipitation area.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if**

their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: inland saltgrass, alkali bluegrass, sandberg bluegrass, Kentucky bluegrass, foxtail barley, poverty sumpweed, seepweed, kochia, black greasewood, suaeda, dandelion, etc.

17. **Perennial plant reproductive capability:** This is not impaired in the reference state. Except in extended periods of drought, plants are able to reproduce sexually or vegetatively.
-