

# Ecological site R058BY204WY Clayey (Cy) 15-17" PZ

Last updated: 10/05/2023 Accessed: 05/18/2024

### **General information**

**Provisional**. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### **MLRA** notes

Major Land Resource Area (MLRA): 058B–Northern Rolling High Plains, Southern Part

MLRA 58B is located in northeastern Wyoming (95 percent) and extreme southeastern Montana (5 percent). It is comprised of sedimentary plains, scoria hills, and river valleys. The major rivers include the Powder, Tongue, Belle Fourche, Cheyenne, and North Platte. Tributaries include the Little Powder River, Little Missouri River, Clear Creek, Crazy Woman Creek, and others. This MLRA is traversed by Interstates 25 and 90, and U.S. Highways 14 and 16. The extent of MLRA 58B covers approximately 12.3 million acres. Major land uses include rangeland (approximately 93 percent), cropland, pasture, and hayland (approximately 2 percent), and forest, urban, and miscellaneous uses (approximately 5 percent). Cities include Buffalo, Casper, Sheridan, and Gillette, WY. Land ownership is mostly private. Federal lands include the Thunder Basin National Grassland (U.S. Forest Service) and lands administered by the Bureau of Land Management. Areas of interest in MLRA 58B in Wyoming include Fort Phil Kearny State Historic Site, Glendo State Park, and Lake DeSmet. The elevations in MLRA 58B increase gradually from north to south and range from approximately 2,900 to 5,900 feet. A few buttes are higher than 6,800 feet. The average annual precipitation in this area ranges from 10 to 17 inches per year. Precipitation occurs mostly during the growing season, often during rapidly developing thunderstorms. Mean annual air temperature is 46 degrees Fahrenheit. Summer temperatures may exceed 100 degrees Fahrenheit. Winter temperatures may drop to below zero. Snowfall averages 45 inches per year, but varies from 25 to over 70 inches in some locales.

### **Classification relationships**

USDA Natural Resources Conservation Service (NRCS): Land Resource Region – G Western Great Plains Range and Irrigation; Major Land Resource Area (MLRA) – 58B Northern Rolling High Plains, Southern Part (USDA, 2006)

Relationship to Other Classifications:

USDA Forest Service (FS) Classification Hierarchy:

Province – 331 Great Plains-Palouse Dry Steppe; Section – 331G-Powder River Basin; Subsections – 331Gb Montana Shale Plains, 331Ge Powder River Basin, 331Gf South Powder River Basin-Scoria Hills (Cleland et al, 1997)

Environmental Protection Agency (EPA) Classification Hierarchy:

Level III Ecoregion – 43 Northwestern Great Plains; Level IV Ecoregion – 43p Scoria Hills, 43q Mesic-Dissected Plains, 43w Powder River Basin (EPA, 2013) https://www.epa.gov/eco-research/ecoregions

### **Ecological site concept**

This ecological site occurs on nearly level to gently sloping sedimentary plains, hills, terraces, and fans at elevations ranging from 2,900 to 5,900 feet. Slopes range from 0 to 15 percent but are generally less than 8 percent. This site occurs on all aspects, although aspect is not a significant factor. The soils of this ecological site are moderately deep to very deep and are well drained. The soil surface textures are typically clay, clay loam, silty clay loam, silty clay, or sandy clay and contain greater than 35 percent clay.

### **Associated sites**

R058BY250WY	Sandy (Sy) 15-17" PZ The Sandy 15-17 ecological site occurs on slopes ranging from 0 to 15 percent, has a soil depth of 20 to 80 inches, and has higher total annual production. The Sandy 15-17 ecological site is generally positioned above the Clayey 15-17 ecological site.
R058BY258WY	Shallow Clayey (SwCy) 15-17" PZ The Shallow Clayey 15-17 ecological site occurs on slopes ranging from 0 to 60 percent, has a soil depth of 10 to 20 inches, and has lower total annual production. The Shallow Clayey 15-17 ecological site is positioned above the Clayey 15-17 ecological site.
R058BY222WY	<b>Loamy (Ly) 15-17" PZ</b> The Loamy 15-17 ecological site occurs on slopes ranging from 0 to 15 percent and has soils with less than 35 percent clay content. The Loamy 15-17 ecological site is located on similar landform positions as the the Clayey 15-17 ecological site.
R058BY228WY	<b>Lowland (LL) 15-17" PZ</b> Lowland 15-17 ecological site occurs on slopes ranging from 0 to 6 percent, has a soil depth of 80 inches, and has higher total annual production. The Lowland 15-17 ecological site is positioned below the Clayey 15-17 ecological site.

### Similar sites

R058BY222WY	Loamy (Ly) 15-17" PZ
	The Loamy 15-17 ecological site occurs on slopes ranging from 0 to 15 percent and has soils with less
	than 35 percent clay content. The Loamy 15-17 ecological site is located on similar landform positions as
	the the Clayey 15-17 ecological site.

### Table 1. Dominant plant species

Tree	Not specified
------	---------------

Shrub	(1) Artemisia tridentata (2) Krascheninnikovia lanata
Herbaceous	(1) Pascopyrum smithii (2) Nassella viridula

### **Physiographic features**

This ecological site occurs on nearly level to gently sloping fans, fan remnants, ridges, and hills on sedimentary plains at elevations ranging from 2,900 to 5,900 feet. Slopes range from 0 to 15 percent. This site occurs on all aspects. Aspect is not a significant factor.

Landforms	(1) Alluvial fan (2) Fan remnant (3) Ridge (4) Hill		
Runoff class	Medium to very high		
Flooding frequency	None		
Ponding frequency	None		
Elevation	884–1,798 m		
Slope	0–15%		
Water table depth	203 cm		
Aspect	Aspect is not a significant factor		

Table 2. Representative physiographic features

### **Climatic features**

The average annual precipitation ranges from 10 to 17 inches per year across MLRA 58B. There are two Precipitation Zones (PZs). The 10 to 14 inch precipitation zone is predominant across the MLRA, including portions of Sheridan, Johnson, and Natrona Counties; portions of Campbell and Converse Counties; and smaller portions of Weston and Niobrara Counties. The 15 to 17 inch precipitation zone occurs in northern and eastern portions of the MLRA, including portions of Sheridan, Campbell, and western Crook Counties. Wide fluctuations in precipitation may occur from year to year, and occasional periods of extended drought (longer than one year in duration) can be expected. Two-thirds of the annual precipitation occurs during the growing season from May through September. Mean Annual Air Temperature (MAAT) is 46 degrees Fahrenheit. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may also occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranching operations during late winter and spring. High-intensity afternoon thunderstorms may occur during the summer. Annual wind speeds average about 5 mph. Daytime winds are generally stronger than nighttime winds. Occasional strong storms may bring brief periods of high winds with gusts of more than 75 mph. The average length of the freeze-free period (28 degrees Fahrenheit) is 125 days and generally occurs from May 16 to September 19. The average frost-free period (32 degrees Fahrenheit) is 101 days and generally occurs from June 1 to September 9.

The growth of native cool-season plants begins in late April to early May with peak growth occurring in mid to late June. Native warm-season plants begin growth in late May to early June and continue into August. Regrowth of cool-season plants occurs in September in most years, depending upon moisture.

Note: The climate described here is based on historic climate station data and is averaged to provide an overview of the annual precipitation, temperatures, and growing season. Future climate is beyond the scope of this document. However, research to determine the effects of elevated CO2 and heating on mixed-grass prairie ecosystems, and how it may relate to future plant communities, is ongoing.

For detailed information, or to find a specific climate station, visit the Western Regional Climate Center (WRCC) website: Western Regional Climate Center, Historical Data, Western U.S. Climate summaries, NOAA Coop

Stations, Wyoming (Note: Montana climate stations are also listed under the Wyoming link). https://wrcc.dri.edu/summary/Climsmwy.html

Wind speed averages can be found at the WRCC home page, under the Specialty Climate tab: https://wrcc.dri.edu/

The following tables represent area-wide climate data for the 15 to 17 inch precipitation zone:

### Table 3. Representative climatic features

Frost-free period (characteristic range)	88-105 days		
Freeze-free period (characteristic range)	122-130 days		
Precipitation total (characteristic range)	381-406 mm		
Frost-free period (actual range)	83-109 days		
Freeze-free period (actual range)	119-130 days		
Precipitation total (actual range)	381-432 mm		
Frost-free period (average)	101 days		
Freeze-free period (average)	125 days		
Precipitation total (average)	381 mm		

### **Climate stations used**

- (1) DOUGLAS 1 SE [USC00482685], Douglas, WY
- (2) BIDDLE 8 SW [USC00240743], Biddle, MT
- (3) DILLINGER [USC00482580], Gillette, WY
- (4) GILLETTE 4SE [USC00483855], Gillette, WY
- (5) LEITER 9N [USC00485506], Clearmont, WY
- (6) SHERIDAN CO AP [USW00024029], Sheridan, WY

### Influencing water features

This upland ecological site is not influenced by a water table or run in from adjacent sites. Due to the semi-arid climate in which it occurs, the water budget is normally contained within the soil pedon. Soil moisture is recharged by spring rains, but it rarely exceeds field capacity in the upper 40 inches before being depleted by evapotranspiration. During intense precipitation events, precipitation rates frequently exceed infiltration rates and the site delivers moisture to downslope sites through surface runoff. Moisture loss through evapotranspiration exceeds precipitation for a majority of the growing season. Soil moisture is the primary limiting factor for vegetative production on this ecological site.

### Wetland description

N/A

### Soil features

The soils associated with this ecological site are typically moderately deep to very deep, well drained, and derived from alluvium or slope alluvium, or residuum derived from shale. The depth to a soil restrictive layer is greater than 20 inches from the soil surface. The surface layer ranges from a depth of 1 to 10 inches in thickness. The soil surface horizon textures are typically clay, clay loam, silty clay loam, or silty clay, but may include loam and contain greater than 32 percent clay. The subsoil horizons typically contain 35 percent clay or greater, have clay, silty clay, silty clay loam, or clay loam textures. Soils on this ecological site typically have carbonates at the surface, but a few soils may be leached to 33 inches. The soil moisture regime is typically ustic aridic and the soil temperature regime is mesic.

Major soil series correlated to this ecological site include Echeta, Leiter, Moorhead, Cromack, Sabatka, and

### Nuncho.

The attributes listed below represent 0 to 40 inches in depth or to the first restrictive layer.

### Table 4. Representative soil features

Parent material	(1) Alluvium–shale (2) Residuum–shale
Surface texture	<ul><li>(1) Clay loam</li><li>(2) Silty clay loam</li><li>(3) Silty clay</li><li>(4) Clay</li></ul>
Drainage class	Well drained
Permeability class	Slow to moderately slow
Depth to restrictive layer	51–203 cm
Soil depth	51–203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	13.21–21.34 cm
Calcium carbonate equivalent (0-101.6cm)	0–10%
Clay content (0-101.6cm)	35–60%
Electrical conductivity (0-101.6cm)	0–4 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–5
Soil reaction (1:1 water) (0-101.6cm)	6.6–8.4
Subsurface fragment volume <=3" (0-101.6cm)	0–5%
Subsurface fragment volume >3" (0-101.6cm)	0%

### **Ecological dynamics**

The Reference State is the plant community in which interpretations are primarily based and is used as a reference in order to understand the original potential of the site. The Reference State evolved under the combined influences of climatic conditions, periodic fire activity, grazing by large herbivores, and impacts from small mammals and insects. Changes may occur to the Reference State due to management actions such as continuous season-long or year-long grazing, increased stocking rates, climatic conditions such as drought, and natural events such as multiple fires in close succession. The reference plant community for this ecological site is dominated by a diversity of tall and medium height, cool-season and warm-season grasses which are tightly intermixed and well distributed over the site. Various forbs, half-shrubs, and shrubs are common on this site. The Reference Plant Community is not necessarily the management goal, as other vegetative states may be considered desired plant communities as long as critical resource concerns are met.

In addition to the Reference State, other plant communities can occur on this site and are usually the result of historic management practices. Grazing practices such as continuous season-long or year-long grazing, heavier stocking rates, or a combination of these factors on this ecological site results in a decrease of tallgrasses, mid-grasses, and more palatable forbs and in an increase of shortgrasses, sedges, and less palatable forbs. Half-shrubs and shrubs increase in the absence of prescribed fire and wildfire. More frequent fire intervals decreases the

shrub component resulting in a site dominated by grasses along with a substantial increase in the forb component (Porensky et al. 2018). There are various transitional stages which may occur on this ecological site. The information presented is representative of a dynamic set of plant communities that illustrate the complex interaction of several ecological processes.

### State and transition model

### Ecosystem states



- T1A Prolonged drought, heavier stocking rates, or a combination of these factors
- T1B Introduction of non-native, invasive species (annual bromes, crested wheatgrass, noxious weeds)
- T1C Tillage or herbicide application and seeding of annual crops or non-native hayland (frequently combined with irrigation practices)
- R2A Lower stocking rates in combination with rangeland seeding, grazing land mechanical treatment, and timely moisture (management intensive and costly)
- T2A Introduction of non-native, invasive species (annual bromes, crested wheatgrass, noxious weeds)
- T2B Tillage or herbicide application and seeding of annual crops or non-native hayland (frequently combined with irrigation practices)
- T3A Tillage or herbicide application and seeding of annual crops or non-native hayland (frequently combined with irrigation practices)
- T4A Cessation of annual cropping
- T5A Tillage or herbicide application and seeding of annual crops or non-native hayland (frequently combined with irrigation practices)

### State 1 submodel, plant communities



P1.1A - Prescribed fire and wildfire, mechanical and chemical treatments, biological processes

- P1.1B Drought, heavier stocking rates
- P1.2A Approximately 80 years post-fire regrowth (Porensky et al. 2018)
- P1.2B Drought, heavier stocking rates, multiple fires in close succession
- P1.3B Normal or above average precipitation, lighter stocking rates
- P1.3A Prescribed fire and wildfire, mechanical and chemical treatments, biological processes
- P1.4B Normal or above average precipitation; lower stocking rates
- P1.4A Approximately 80 years post-fire regrowth (Porensky et al. 2018)

### State 2 submodel, plant communities



P2.1A - Prescribed fire and wildfire, mechanical and chemical treatments, biological processes

P2.2A - Approximately 80 years post-fire regrowth (Porensky et al. 2018)

### State 3 submodel, plant communities

3.1. Bromus tectorum-
Bromus arvensis
(cheatgrass-field
brome)

#### State 4 submodel, plant communities

4.1. Cropland Community

State 5 submodel, plant communities

5.1. Abandoned Cropland Community

5.2. Perennial Grass Community

## State 1 Reference State

The Reference State (1) evolved under the combined influences of climatic conditions, periodic fire activity, grazing by large herbivores, and impacts from small mammals and insects. The Reference State (1) is the plant community in which interpretations are primarily based and is used as a reference in order to understand the original potential of the site. The plant communities and various successional stages represent the natural range of variability. The Reference State (1) for this ecological site consists of 4 community phases.

### Community 1.1 Artemisia tridentata wyomingensis/Nassella viridula-Pascopyrum smithii (Wyoming big sagebrush/green needlegrass-western wheatgrass)

Community Phase (1.1) is characterized by a mixed-grass community and Wyoming big sagebrush. The predominant species are cool-season grasses such as green needlegrass and western wheatgrass, with a smaller component of warm-season grasses. Shortgrasses and sedges such as blue grama, prairie junegrass, Sandberg bluegrass, threadleaf sedge, and needleleaf sedge are present at low cover. Forbs such as American vetch, upright prairie coneflower, large Indian breadroot, milkvetch, sulphur-flower buckwheat, purple prairie clover, white prairie clover, and tapertip hawksbeard are present. The dominant shrub species is Wyoming big sagebrush, although other shrubs such as fourwing saltbush and sub-shrubs such as winterfat occur at lower canopy covers. The potential vegetation (air-dry weight) is approximately 71 percent grasses, 10 percent forbs, and 19 percent shrubs. The total annual production (air-dry weight) is approximately 1,900 pounds per acre during an average year, but can range from approximately 1,500 pounds per acre in below average years to approximately 2,300 pounds per acre in above average years.

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1194	1512	1830
Shrub/Vine	319	405	490
Forb	168	213	258
Total	1681	2130	2578

### Table 5. Annual production by plant type

Figure 9. Plant community growth curve (percent production by month). WY1501, 15-17NP Upland sites.

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	25	40	10	5	5	5	0	0

### Community 1.2 Nassella viridula-Pascopyrum smithii (green needlegrass-western wheatgrass)

Community Phase 1.2 is characterized by a reduction in Wyoming big sagebrush and an increase in cool-season, rhizomatous wheatgrasses such as western wheatgrass and thickspike wheatgrass and mid-statured bunchgrasses such as green needlegrass and bluebunch wheatgrass. Shortgrasses and sedges such as blue grama, prairie junegrass, Sandberg bluegrass, buffalograss, threadleaf sedge, and needleleaf sedge occur at low cover. This community phase results in a substantial increase in the forb component with species such as American vetch, upright prairie coneflower, large Indian breadroot, milkvetch, sulphur-flower buckwheat, purple prairie clover, white

prairie clover, and tapertip hawksbeard (Porensky et al. 2018). Half-shrubs such as winterfat may occur. Wyoming big sagebrush is rare or absent. This community occurs following sagebrush mortality due to wildfire, prescribed fire, mechanical and chemical treatments, or biological processes and may persist for 80 years or longer (Porensky et al. 2018).

### Community 1.3 Artemicia tridenteta uncomingencia/Research/

# Artemisia tridentata wyomingensis/Pascopyrum smithii-Bouteloua gracilis (Wyoming big sagebrush/western wheatgrass-blue grama)

Community Phase 1.3 is characterized by an increase in shortgrass species and sedges such as blue grama, prairie junegrass, Sandberg bluegrass, threadleaf sedge, and needleleaf sedge, and a decrease in mid-statured grasses such as western wheatgrass and green needlegrass. Shortgrass species dominate the plant community and mid-statured grasses are less common and contribute little to overall production. This plant community occurs due to long-term drought, grazing management such as continuous season-long or year-long grazing, increased stocking rates, or a combination of these factors.

# Community 1.4 Pascopyrum smithii-Bouteloua gracilis (western wheatgrass-blue grama)

Community Phase 1.4 is characterized by an increase in shortgrass species and sedges such as blue grama, prairie junegrass, Sandberg bluegrass, threadleaf sedge, and needleleaf sedge, and a decrease in mid-statured grasses such as western wheatgrass and green needlegrass. Wyoming big sagebrush is rare or absent. This plant community occurs due to long-term drought, grazing management such as continuous season-long or year-long grazing, increased stocking rates, or a combination of these factors. Sagebrush mortality is evident due to wildfire, prescribed fire, mechanical and chemical treatments, or biological processes.

### Pathway P1.1A Community 1.1 to 1.2

Community Phase 1.1 will transition to Community Phase 1.2 through prescribed fire and wildfire, mechanical and chemical treatments, and biological processes. Wyoming big sagebrush is greatly reduced and perennial grasses will dominate the site.

### Pathway P1.1B Community 1.1 to 1.3

Drought, grazing practices such as continuous season-long or year-long grazing, heavier stocking rates, or a combination of these factors can shift Community Phase 1.1 to Community Phase 1.3. These factors favor an increase in shortgrasses such as blue grama and a decrease in cool-season mid-statured grasses. Wyoming big sagebrush cover will be similar to Community Phase 1.1.

### Pathway P1.2A Community 1.2 to 1.1

Approximately 80 years or more of natural vegetative regrowth will transition Community Phase 1.2 to Community Phase 1.1. Approximately 80 years or more without fire allows Wyoming big sagebrush to recolonize the site (Porensky et al. 2018).

### Pathway P1.2B Community 1.2 to 1.4

Drought, grazing practices such as continuous season-long or year-long grazing, heavier stocking rates, multiple fires in close succession, or a combination of these factors can shift Community Phase 1.2 to Community Phase 1.3. These factors favor an increase in blue grama and a decrease in cool-season mid-statured grasses. Wyoming big sagebrush cover will be similar to Community Phase 1.2.

### Pathway P1.3B Community 1.3 to 1.1

Approximately 80 years or more post-fire (Porensky et al. 2018), normal or above average precipitation, and lighter stocking rates transitions Community Phase 1.3 to Community Phase 1.1. This transition may also occur through natural succession.

# Pathway P1.3A Community 1.3 to 1.4

Community Phase 1.3 will transition to Community Phase 1.4 through prescribed fire and wildfire, mechanical and chemical treatments, and biological processes. Wyoming big sagebrush is greatly reduced and perennial grasses will dominate the site.

## Pathway P1.4B Community 1.4 to 1.2

Normal or above average precipitation and lower stocking rates transitions Community Phase 1.4 to Community Phase 1.2.

## Pathway P1.4A Community 1.4 to 1.3

Approximately 80 years or more of natural shrub regrowth will transition Community Phase 1.4 to Community Phase 1.3. Approximately 80 years or more without prescribed fire or wildfire allows Wyoming big sagebrush to recolonize the site (Porensky et al. 2018).

### State 2 Shortgrass State

The dynamics of the Shortgrass State (2) are driven by long-term drought, grazing management such as continuous season-long or year-long grazing, or a combination of these factors. The Shortgrass State (2) for this ecological site consists of 2 community phases.

# Community 2.1

# Artemisia tridentata wyomingensis/Bouteloua gracilis-Carex filifolia (Wyoming big sagebrush/blue grama-threadleaf sedge)

Community Phase 2.1 is characterized by a dominance of shortgrasses and sedges such as blue grama, prairie junegrass, Sandberg bluegrass, and threadleaf sedge. Mid-statured grasses such as western wheatgrass and green needlegrass are rare or absent. Prairie sagewort, broom snakeweed, and Wyoming big sagebrush are common. This plant community occurs when site conditions decline due to long-term drought, grazing management such as continuous season-long or year-long grazing, or a combination of these factors. This community phase results in a reduction of soil surface litter, soil organic matter, and infiltration and an increase of soil surface runoff. This plant community is capable of tolerating season-long, heavy grazing and therefore is highly resistant to change.

# Community 2.2 Bouteloua gracilis-Carex filifolia (blue grama-threadleaf sedge)

Community Phase 2.2 is characterized by a complete dominance of shortgrasses and sedges such as blue grama, prairie junegrass, Sandberg bluegrass, threadleaf sedge, and needleleaf sedge. Cool-season, mid-statured grasses such as western wheatgrass and green needlegrass are rare or absent. Prairie sagewort is common and Wyoming big sagebrush is rare. This plant community occurs when site conditions decline due to long-term drought, grazing management such as continuous season-long or year-long grazing, or a combination of these factors, and a fire has occurred on the site less than approximately 30 years prior. Sagebrush mortality is evident due to wildfire, prescribed fire, mechanical and chemical treatments, or biological processes. This community phase results in a

reduction of soil surface litter, soil organic matter, and infiltration and an increase of soil surface runoff. This plant community is capable of tolerating season-long, heavy grazing and therefore is highly resistant to change.

### Pathway P2.1A Community 2.1 to 2.2

Community Phase 2.1 will transition to Community Phase 2.2 through prescribed fire and wildfire, mechanical and chemical treatments, and biological processes. Wyoming big sagebrush is greatly reduced and perennial grasses will dominate the site.

# Pathway P2.2A Community 2.2 to 2.1

It is estimated that approximately 80 years or more of natural post-fire vegetative regrowth (Porensky et al. 2018) could transition Community Phase 2.2 to Community Phase 2.1. It is possible that this transition could occur over time, however, the processes are not fully understood at this time. Therefore, this pathway is considered hypothetical until further investigation can be completed.

### State 3 Invaded State

The Invaded State (3) occurs when invasive plant species invade native plant communities and displace the native species. The Invaded State (3) consists of 1 community phase.

### Community 3.1 Bromus tectorum-Bromus arvensis (cheatgrass-field brome)

Observations suggest that native species diversity declines significantly when invasive or noxious species exceed approximately 30 percent of the plant community. Non-native, perennial, drought tolerant grasses such as crested wheatgrass, non-native, annual, invasive species such as cheatgrass and field brome, and noxious weed species can eventually dominate the seedbank of this site and displace native species. Reduced plant species diversity, simplified structural complexity, and altered ecological processes result in a state that is substantially departed from the Reference State (1). The dominance of annual, invasive grasses such as cheatgrass and field brome increases the fire cycle frequency.

### State 4 Cropland State

The Cropland State (4) occurs when cultivation occurs to the land. The Cropland State (4) consists of 1 community phase.

### Community 4.1 Cropland Community

The land is cultivated and converted to crop production. Annual, cool-season cereal grains such as spring wheat, winter wheat, and barley are common crops which replace native plant communities.

# State 5 Post-Cropland State

The Post-Cropland State (5) occurs when cultivated cropland is abandoned and allowed to either re-vegetate naturally or is seeded back to perennial species for livestock grazing or wildlife use. This state can transition back to the Cropland State (4) if the site is returned to cultivation. No formal studies have been obtained regarding Wyoming big sagebrush recovery following cultivation and further investigation is needed to assess Wyoming big sagebrush recovery in the Post-Cropland State (5). The Post-Cropland State (5) has 2 community phases.

### Community 5.1 Abandoned Cropland Community

In the absence of active management, the site can re-vegetate naturally and potentially return to a perennial grassland community over time. Shortly after cropland is abandoned, annual and biennial forbs and annual brome grasses invade the site. The site is highly susceptible to erosion due to the absence of perennial species. Eventually, these pioneering annual species are replaced by perennial forbs and perennial shortgrasses. Depending on the historical management of the site, mid-statured perennial grasses may also return; however, species composition will depend upon the seed bank. Invasion of the site by exotic species, such as crested wheatgrass and annual bromes, will depend upon the site's proximity to a seed source. Approximately 50 or more years after cultivation, these sites may have species composition similar to phases in the Reference State (1) (Dormaar, J.F., and S. Smoliak. 1985). However, soil quality is consistently lower than conditions prior to cultivation and a shift to the Reference State (1) is unlikely.

# Community 5.2 Perennial Grass Community

When the site is seeded to perennial forage species this community phase can persist for several decades. Introduced perennial grasses, in particular, may form monocultures that persist for approximately 60 years or more (Samuel, M.J., and R.H. Hart. 1994). A mixture of native species may also be seeded to provide species composition and structural complexity similar to that of the Reference State (1). However, soil quality conditions have been substantially altered and will not return to pre-cultivation conditions.

### Transition T1A State 1 to 2

Prolonged drought, grazing management such as continuous season-long or year-long grazing, heavier stocking rates, or a combination of these factors weaken the resilience of the Reference State (1) and drive its transition to the Shortgrass State (2). The Reference State (1) transitions to the Shortgrass State (2) when mid-statured grasses are greatly reduced and shortgrasses and sedges such as blue grama, Sandberg bluegrass, prairie junegrass, and threadleaf sedge dominate the plant community.

### Transition T1B State 1 to 3

The Reference State (1) transitions to the Invaded State (3) when non-native grasses or noxious weeds invade the plant community. Exotic plant species dominate the site in terms of cover and production and site resilience has been substantially reduced. In addition, other rangeland health attributes, such as reproductive capacity of native grasses and soil quality, have been substantially altered from the Reference State (1).

### Transition T1C State 1 to 4

Tillage or application of herbicide followed by seeding of cultivated crops, such as winter wheat, spring wheat, and barley, transitions the Reference State (1) to the Cropland State (4).

# Restoration pathway R2A State 2 to 1

Blue grama can resist displacement by other species. A reduction in stocking rates alone may not be sufficient to reduce the cover of blue grama in the Shortgrass State (2) and mechanical treatments may be necessary. Therefore, returning the Shortgrass State (2) to the Reference State (1) could require considerable cost, energy, and time.

Transition T2A State 2 to 3 The Shortgrass State (2) transitions to the Invaded State (3) when non-native grasses, noxious weeds, and other invasive plants invade the Shortgrass State (2). Exotic plant species dominate the site in terms of cover and production. Site resilience has been substantially reduced.

### Transition T2B State 2 to 4

Tillage or application of herbicide followed by seeding of cultivated crops, such as winter wheat, spring wheat, and barley, transitions the Shortgrass State (2) to the Cropland State (4).

### Transition T3A State 3 to 4

The Invaded State (3) will transition to the Cropland State (4) when the site is placed under cultivation.

### Transition T4A State 4 to 5

The transition from the Cropland State (4) to the Post-Cropland State (5) occurs with the cessation of cultivation. The site may also be seeded to perennial forage species, such as crested wheatgrass and alfalfa, or a mix of native species.

### Transition T5A State 5 to 4

Tillage or application of herbicide followed by seeding of cultivated crops, such as winter wheat, spring wheat, and barley, transitions the Post-Cropland State (5) to the Cropland State (4).

### Additional community tables

Table 6. Community 1.1 plant community composition

Group	oup Common Name Symbol Scientific Name		Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike		•	-	
1	Cool-Season Rhizomato	ous Grasse	S	504–773	
	western wheatgrass	PASM	Pascopyrum smithii	504–773	5–30
	thickspike wheatgrass	ELLAL	Elymus lanceolatus ssp. lanceolatus	504–773	5–30
2	Cool-Season Bunchgras	sses	•	841–1289	
	green needlegrass	NAVI4	Nassella viridula	841–1289	10–50
3	Warm-Season Shortgra	sses	•	168–258	
	blue grama	BOGR2	Bouteloua gracilis	168–258	5–10
4	Miscellaneous Grasses		•	336–516	
	Grass, perennial	2GP	Grass, perennial	84–129	1–5
	plains reedgrass	CAMO	Calamagrostis montanensis	84–129	1–5
	buffalograss	BODA2	Bouteloua dactyloides	84–129	1–5
	prairie Junegrass	KOMA	Koeleria macrantha	84–129	1–5
	Sandberg bluegrass	POSE	Poa secunda	84–129	1–5
	needleleaf sedge	CADU6	Carex duriuscula	84–129	1–5
	threadleaf sedge	CAFI	Carex filifolia	84–129	1–5
	big bluestem	ANGE	Andropogon gerardii	84–129	1–5
	aidaaata arama	PACH	Poutolous outinondulo	0/ 100	1 5

I -	-0

Forb					
5	Forbs			252–387	
	western yarrow	ACMIO	Achillea millefolium var. occidentalis	84–129	1–5
	Forb, perennial	2FP	Forb, perennial	84–129	1–5
	American vetch	VIAM	Vicia americana	84–129	1–5
	upright prairie coneflower	RACO3	Ratibida columnifera	84–129	1–5
	aster	ASTER	Aster	84–129	1–5
	desertparsley	LOMAT	Lomatium	84–129	1–5
	large Indian breadroot	PEES	Pediomelum esculentum	84–129	1–5
	rosy pussytoes	ANRO2	Antennaria rosea	84–129	1–5
	milkvetch	ASTRA	Astragalus	84–129	1–5
	stemless mock goldenweed	STAC	Stenotus acaulis	84–129	1–5
	sulphur-flower buckwheat	ERUM	Eriogonum umbellatum	84–129	1–5
	scarlet beeblossom	GACO5	Gaura coccinea	84–129	1–5
	purple prairie clover	DAPU5	Dalea purpurea	84–129	1–5
	white prairie clover	DACA7	Dalea candida	84–129	1–5
	bluebells	MERTE	Mertensia	84–129	1–5
	textile onion	ALTE	Allium textile	84–129	1–5
	tapertip hawksbeard	CRAC2	Crepis acuminata	84–129	1–5
Shrub	/Vine				
6	Shrubs			168–258	
	Wyoming big sagebrush	ARTRW8	Artemisia tridentata ssp. wyomingensis	168–258	5–10
7	Miscellaneous Shrubs			336–516	
	winterfat	KRLA2	Krascheninnikovia lanata	84–129	1–5
	fourwing saltbush	ATCA2	Atriplex canescens	84–129	1–5
	Shrub (>.5m)	2SHRUB	Shrub (>.5m)	84–129	1–5
	Subshrub (<.5m)	2SUBS	Subshrub (<.5m)	84–129	1–5

### Animal community

Rhizomatous Wheatgrasses, Green needlegrass Community: The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include Western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here.

Heavy Sagebrush Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15% protein and 40-60% digestibility during that time. This community provides excellent escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for sage grouse.

Greasewood/ wheatgrass plant community: This plant community exhibits a low level of plant species diversity due

to the accumulation of salts in the soil. It may provide some thermal and escape cover for deer and antelope if no other woody community is nearby, but in most cases it is not a desirable plant community to select as a wildlife habitat management objective.

Mixed Sagebrush/Grass Plant Community: The combination of an overstory of sagebrush and an understory of grasses and forbs provides a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows nest in big sagebrush plants, and a host of other nesting birds utilize stands in the 20-30% cover range.

Blue Grama Sod Plant Community: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Western Wheatgrass/Cheatgrass Plant Community are limiting. Generally, these are not target plant communities for wildlife habitat management.

Go-back Land: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Western Wheatgrass/Cheatgrass Plant Community are limiting. Generally, these are not target plant communities for wildlife habitat management.

Introduced Pasture: These communities are highly variable depending on the species planted. Refer to Forage Suitability Groups for more information.

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community Production Carrying Capacity\* (lb./ac) (AUM/ac) Reference Plant Community 600-1400 .5 Heavy Sagebrush 600-1200 .33 Blue Grama Sod 400- 900 .25 Mixed Sagebrush/Grass 600-1200 .4 Greasewood/wheatgrass 500- 900 .25 Go-back Land 500- 900 .2 Introduced Pasture 800-2000 1

\* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

### Hydrological functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group C and D. Infiltration ranges from slow to moderately slow. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover

have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals should not be present. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

### **Recreational uses**

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

### Wood products

No appreciable wood products are present on the site.

### Other products

None noted.

### **Other information**

Site Development & Testing Plan

General Data (MLRA and Revision Notes, Hierarchical Classification, Ecological Site Concept, Physiographic, Climate, and Water Features, and Soils Data):

Updated. All "Required" items complete to Provisional level.

Community Phase Data (Ecological Dynamics, STM, Transition & Recovery Pathways, Reference Plant Community, Species Composition List, Annual Production Table):

Updated. All "Required" items complete to Provisional level.

Annual Production Table is from the "Previously Approved" ESD (2001).

The Annual Production Table and Species Composition List will be reviewed for future updates at the Approved level.

Each Alternative State/Community:

Complete to Provisional level.

Supporting Information (Site Interpretations, Assoc. & Similar Sites, Inventory Data References, Agency/State Correlation, References):

Updated. All "Required" items complete to Provisional level.

Wildlife Interpretations: Narrative is from "Previously Approved" ESD (2001). Wildlife species will need to be updated at the next Approved level.

Livestock Interpretations: Plant community names and stocking rates updated.

Hydrology, Recreational Uses, Wood Products, and Other Products carried over from previously "Approved" ESD (2001).

Existing NRI Inventory Data References updated. More field data collection is needed to support this site concept.

**Reference Sheet** 

Rangeland Health Reference Sheet carried over from previously "Approved" ESD (2005). It will be updated at the next "Approved" level.

"Future work, as described in a project plan, to validate the information in this provisional ecological site description is needed. This will include field activities to collect low and medium intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document." (NI 430\_306 ESI and ESD, April 2015)

### Inventory data references

Inventory information has been derived from data collection on private and federal lands by the following methods: • Double Sampling (Determining Vegetation Production and Stocking Rates, WY-ECS-1)

- Rangeland Health (Interpreting Indicators of Rangeland Health, Version 4, 2005)
- Soil Stability (Interpreting Indicators of Rangeland Health, Version 4, 2005)
- Line Point Intercept (Monitoring Manual for Grassland, Shrubland, and Savanna Ecosystems, Volume II, 2005)
- Soil Pedon Descriptions (Field Book for Describing and Sampling Soils, Version 3, 2012)
- SCS-RANGE-417 (Production & Composition Record for Native Grazing Lands)

National Resources Inventory (NRI)

- Number of Records: 5
- Sample Period: 2012-2015
- State: Wyoming
- Counties: Campbell, Sheridan, Weston

USDA - Agricultural Research Service (ARS)

Thunder Basin National Grassland

Plant Community Responses to Historical Wildfire in a Shrubland/Grassland Ecotone

- Number of Records: 140
- Sample Period: 2014-2021
- State: Wyoming
- Counties: Campbell, Converse, Crook, Niobrara, and Weston

Additional reconnaissance data collection includes ocular estimates and other inventory data; vegetative clipping data for NRCS program support; field observations from experienced rangeland personnel

Data collection for this ecological site was done in conjunction with the progressive soil surveys within MLRA 58B Northern Rolling High Plains (Southern Part)

Note: Revisions to soil surveys are on-going. For the most recent updates, visit the Web Soil Survey, the official site for soils information: http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx

### **Other references**

Cleland, D.T., et al. 1997. National hierarchical framework of ecological units. In: M.S. Boyce and A. Haney (eds.) Ecosystem Management Applications for Sustainable Forest and Wildlife Resources, Yale University Press, New Haven, CT.

Dormaar, J.F., and S. Smoliak. 1985. Recovery of vegetative cover and soil organic matter during revegetation of abandoned farmland in a semiarid climate. Journal of Range Management 38:487-491.

Federal Geographic Data Committee. 2008. The National Vegetation Classification Standard, Version 2. FGDC Vegetation Subcommittee. FGDC-STD-005-2008 (Version 2). pp. 126.

Herrick JE, Van Zee J, Havstad K, Burkett LM, Whitford WG. 2005. Monitoring Manual for Grassland, Shrubland,

and Savanna Ecosystems, Volume II. Tucson, AZ. University of Arizona Press – Distributor. 2005.

Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. USDA Handbook 296. USDA Natural Resources Conservation Service. 2006.

McNab, W.H., et al. 2007. Description of Ecological Sub-Regions: Sections of the Conterminous United States. USDA Forest Service. General Technical Report WO-76B.

Pellant M., Shaver P., Pyke D., Herrick JE. 2005. Interpreting Indicators of Rangeland Health, Version 4.0. Technical Reference 1734-6. Denver, CO. USDI Bureau of Land Management, NSTC, Division of Science Integration, Branch of Publishing Services.

Porensky, L. M., J. D. Derner, and D. W. Pellatz. 2018. Plant community responses to historical wildfire in a shrubland–grassland ecotone reveal hybrid disturbance response. Ecosphere 9(8):e02363. 10.1002/ecs2.2363.

Porensky, L. M., Rachel McGee, and David W. Pellatz. 2020. Long-term grazing removal increased invasion and reduced native plant abundance and diversity in a sagebrush grassland. Global Ecology and Conservation 24 (2020) e01267.

Samuel, M.J., and R.H. Hart. 1994. Sixty-one years of secondary succession on rangelands of the Wyoming High Plains. Journal of Range Management 47:184-191.

Schoeneberger, Wysocki, Benham, and Soil Science Division Staff. 2012. Field Book for Describing and Sampling Soils, Version 3. Washington, DC. United States Government Publishing Office (GPO).

Soil Science Division Staff. USDA Natural Resources Conservation Service. 2014. Keys to Soil Taxonomy, 12th Edition.

Soil Science Division Staff. USDA Natural Resources Conservation Service. 2017. Soil Survey Manual, Agriculture Handbook No. 18.

Soil Science Division Staff. USDA Natural Resources Conservation Service. 2019. National Soil Survey Handbook, Title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/?cid=nrcs142p2\_054242 (Accessed 16 January, 2018).

Soil Science Division Staff. USDA Natural Resources Conservation Service. National Soil Information System (NASIS) Database. https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geo/?cid=nrcs142p2\_053552 (Accessed 30 October, 2017).

Soil Science Division Staff. USDA Natural Resources Conservation Service. Official Soil Series Descriptions. https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2\_053587 (Accessed 15 November, 2017).

Soil Science Division Staff. USDA Natural Resources Conservation Service. Soil Survey Geographic (SSURGO) Database.

Soil Science Division Staff. USDA Natural Resources Conservation Service. Web Soil Survey. https://websoilsurvey.nrcs.usda.gov/app/HomePage.htm (Accessed 15 February, 2018).

Stubbendieck, James, S.L. Hatch, and L.M. Landholt. 2003. North American Wildland Plants. University of Nebraska Press, Lincoln and London.

United States Department of Commerce, National Oceanic and Atmospheric Administration (NOAA). Cooperative Climatological Data Summaries. NOAA Western Regional Climate Center, Reno, NV. http://www.wrcc.dri.edu/climatedata/climsum (Accessed 16 November, 2017).

United States Department of the Interior, Geological Survey. LANDFIRE 1.1.0 Existing Vegetation Types. 2011. http://landfire.cr.usgs.gov/viewer/. United States Department of the Interior, Geological Survey. LANDFIRE 1.1.0 Vegetation Dynamics Models. 2008. http://landfire.cr.usgs.gov/viewer/.

United States Environmental Protection Agency, National Health and Environmental Effects Research Laboratory. 2013. Level III and IV Eco-Regions of the Continental United States. https://www.epa.gov/eco-research/ecoregions (Accessed 30 January, 2019).

USDA Forest Service. Fire Effects Information System.http://www.fs.fed.us/database/feis/plants/shrub/amealn/all.html.

USDA Natural Resources Conservation Service, USDA Forest Service, USDI Bureau of Land Management. January 2013. Interagency Ecological Site Handbook for Rangelands.

USDA Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. USDA Handbook 296.

USDA Natural Resources Conservation Service. Climate Data. National Water and Climate Center. https://www.wcc.nrcs.usda.gov/climate (Accessed 13 October, 2017).

USDA Natural Resources Conservation Service. Glossary of Landform and Geologic Terms. National Soil Survey Handbook, Title 430-VI, Part 629.02c. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ref/? cid=nrcs142p2\_054242 (Accessed 16 January, 2018).

USDA Natural Resources Conservation Service. National Cooperative Soil Survey. https://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/survey/partnership/ncss/.

USDA Natural Resources Conservation Service. National Ecological Site Handbook, Title 190. March 2017. https://directives.sc.egov.usda.gov/ (Accessed 15 September, 2017).

USDA Natural Resources Conservation Service. National Range and Pasture Handbook. 1997, Revised 2003. http://www.glti.nrcs.usda.gov/technical/publications/nrph.html (Accessed 26 February, 2018).

USDA Natural Resources Conservation Service. NRCS Plants Database. https://plants.usda.gov/java/.

### Contributors

Everett Bainter Glenn Mitchell

### Approval

Kirt Walstad, 10/05/2023

### Acknowledgments

Project Staff:

Kimberly Diller, Ecological Site Specialist, NRCS MLRA SSO, Pueblo CO Mike Leno, Project Leader, NRCS MLRA SSO, Buffalo WY

Partners/Contributors:

Joe Dyer, Soil Scientist, NRCS MLRA SSO, Buffalo WY Arnie Irwin, Soil Scientist, BLM, Buffalo WY Blaine Horn, Rangeland Extension Educator, UW Extension, Buffalo WY Isabelle Giuliani, Resource Soil Scientist, NRCS, Douglas WY Mary Jo Kimble, Project Leader, NRCS MLRA SSO, Miles City MT Ryan Murray, Area Rangeland Management Specialist, NRCS, Buffalo WY Lauren Porensky, PhD, Ecologist, ARS, Fort Collins CO Chadley Prosser, Rangeland Program Manager, USFS, Bismarck ND Bryan Christensen, Ecological Site Specialist, NRCS MLRA SSO, Pinedale WY Marji Patz, Ecological Site Specialist, NRCS MLRA SSO, Powell WY Rick Peterson, Ecological Site Specialist, NRCS MLRA SSO, Rapid City SD Jeff Fenton, Ecological Site Specialist, NRCS MLRA SSO, Buffalo WY

Program Support:

John Hartung, WY State Rangeland Management Specialist (QC), NRCS, Casper WY David Kraft, Senior Regional Ecological Site Specialist (QA), NRCS, Emporia KS Carla Green Adams, Editor, NRCS-SSR5, Denver CO Chad Remley, Regional Director, Northern Central Soil Survey Region, Salina KS

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/18/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:

- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth ( in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability: