

## Ecological site R060AY020SD Loamy Overflow

Accessed: 05/18/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 060A–Pierre Shale Plains

#### MLRA Notes:

The Pierre Shale Plains (MLRA 60A) consists of approximately 10,150 square miles, the majority of which is located in South Dakota (70 percent) and small portions are in Montana (2 percent), Nebraska (8 percent), and Wyoming (20 percent). It encircles the Black Hills (MLRA 62) and the Dakota Hogback (MLRA 61). MLRA 60A includes portions of the Oglala, Buffalo Gap, and Thunder Basin National Grasslands. It also includes small sections of the Pine Ridge Indian Reservation, Badlands National Park, and Black Hills National Forest. The Cheyenne and Belle Fourche Rivers flow through the MLRA.

MLRA 60A is in the unglaciated section of the Missouri Plateau, of the Great Plains Province of the Interior Plains. It is an area of old plateaus and terraces that have been deeply eroded. Cretaceous Pierre Shale underlies almost all of this MLRA. This is a marine sediment with layers of volcanic ash that has been altered to smectitic clay. These clays shrink as they dry and swell as they receive moisture. Soils are shallow to very deep and generally are well drained and clayey.

Elevations generally range from 2,620 to 3,610 feet throughout the MLRA, but can range up to 4,260 feet. The average annual precipitation for the western side of the MLRA is 13 to 16 inches, whereas the eastern side receives 16 to 18 inches. A suite of ecological sites have been written specifically for these two precipitation zones. The Locator Map shows the break between the two precipitation zones.

This area supports a mixed natural prairie vegetation consisting of both cool- and warm-season grasses and forbs. Wyoming big sagebrush occurs primarily in the drier western portion of the MLRA; however, small remnant stands can be found in the eastern portion. Dominant land uses of the area are primarily ranching and, to a lesser extent, farming. Major resource concerns to this MLRA are wind erosion and surface water quality.

## Classification relationships

USDA - Land Resource Region G – Western Great Plains Range and Irrigated Region, Major Land Resource Area (MLRA) 60A – Pierre Shale Plains.

EPA - Level IV Ecoregions of the Continental United States: 43e – Sagebrush Steppe, 43g Semiarid Pierre Shale Plains, and 43k – Dense Clay Prairie.

## Ecological site concept

The Loamy Overflow ecological site occurs throughout the MLRA. It is located on Stream Orders 2 or greater. This site is a run-in site and receives additional moisture through runoff from adjacent sites and overflow during occasional flooding. Typical slopes range from 0 to 5 percent. The soil surface layer is 5 to 20 inches in depth with a texture range of clay loam to very fine sandy loam. When degraded, this site is susceptible to down cutting. The natural vegetation will gradually shift from almost exclusively herbaceous species in the upper reaches of a drainageway, to a mix of species including grasses, forbs, shrubs, and trees in the lower reaches. Vegetation in the Reference State (1.0) consists primarily of warm- and cool-season tall- and mid-grasses. Big bluestem and switchgrass are the dominant warm-season grasses, while western wheatgrass and needlegrasses are the dominant cool-season grasses. Forbs are common and very diverse. Patches of western snowberry, American plum, chokecherry, and willow are almost always present. In the western portion of the MLRA, Wyoming big sagebrush will likely be present. Tree species can exist throughout the site but are more likely to occur in the lower reaches. Major tree species include plains cottonwood, green ash, boxelder and willow. When disturbed, this site is very susceptible to invasion of non-native cool-season grasses, Canada thistle, hound's tongue, and other weedy forbs. Russian olive and saltcedar are invasive tree species that can dominate this site if not actively treated.

## Associated sites

R060AY003SD	<b>Subirrigated</b> The Subirrigated site can be found adjacent to or, in some cases, intermixed with the Loamy Overflow site.
R060AY010SD	<b>Loamy 13-16" P.Z.</b> The Loamy 13-16 PZ site will be located on upland landscape positions above the overflow site.
R060AY022SD	<b>Loamy Terrace</b> The Loamy Terrace site will be located on a stream or river terrace above the overflow site, but typically below the adjacent upland sites.
R060AY041SD	<b>Loamy 16-18" P.Z.</b> The Loamy 16-18 PZ site will be located on upland landscape positions above the overflow site.

## Similar sites

R060AY010SD	<b>Loamy 13-16" P.Z.</b> On the upper reaches of the Loamy Overflow, the adjacent Loamy 13-16 PZ will have more rhizomatous wheatgrass, more needlegrass, more big sagebrush, and lower forage production.
R060AY022SD	<b>Loamy Terrace</b> The Loamy Terrace site will have little, if any, tree regeneration, more sagebrush, more rhizomatous wheatgrass, more needlegrass, and lower forage production.
R060AY041SD	<b>Loamy 16-18" P.Z.</b> On the upper reaches of the Loamy Overflow site, the adjacent Loamy 16-18 PZ will have less big bluestem, more western wheatgrass, more needlegrass less shrubs, and lower forage production.
R060AY003SD	<b>Subirrigated</b> The Subirrigated site will have more big bluestem and prairie cordgrass and less green needlegrass.

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Pascopyrum smithii</i>

## Physiographic features

This site occurs on nearly level to gently sloping uplands and river valleys.

**Table 2. Representative physiographic features**

Landforms	(1) Alluvial fan (2) Flood plain (3) Stream terrace
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Rare to frequent
Ponding frequency	None
Elevation	762–1,311 m
Slope	0–5%
Water table depth	107–203 cm
Aspect	Aspect is not a significant factor

## Climatic features

The climate in this MLRA is typical of the drier portions of the Northern Great Plains, where sagebrush steppes to the west yield to grassland steppes to the east. Annual precipitation for the entire MLRA ranges from 13 to 18 inches per year, with most occurring during the growing season. Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air masses from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but the more severe occur during late fall, late winter, and spring. The normal average annual temperature is about 46°F. January is the coldest month with average temperatures ranging from about 19°F (Moorcroft CAA, WY) to about 22°F (Belle Fourche, SD). July is the warmest month with temperatures averaging from about 70°F (Moorcroft CAA, WY) to about 72°F (Belle Fourche, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 51°F. Hourly winds are estimated to average about 11 miles per hour annually, ranging from about 13 miles per hour during the spring to about 10 miles per hour during the summer. Daytime winds generally are stronger than nighttime, and occasional strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour. Growth of cool-season plants begins in early to mid-March, slowing or ceasing in late June. Warm-season plants begin growth about mid-May and can continue to early or mid-September. Green-up of cool-season plants may occur in September and October when adequate soil moisture is present.

**Table 3. Representative climatic features**

Frost-free period (average)	115 days
Freeze-free period (average)	133 days
Precipitation total (average)	432 mm

## Climate stations used

- (1) UPTON [USC00489205], Upton, WY

- (2) REDBIRD [USC00487555], Lance Creek, WY
- (3) ARDMORE 1 NW [USC00390236], Edgemont, SD
- (4) BELLE FOURCHE [USC00390559], Belle Fourche, SD
- (5) WASTA [USC00398911], Owanka, SD
- (6) MOORCROFT 3S [USW00024088], Moorcroft, WY

## Influencing water features

Stream Type: B6, C6  
(Rosgen System)

## Soil features

The soils in this site are moderately well to well drained and formed in alluvium. The clay loam to fine sandy loam surface layer is 5 to 20 inches thick. The soils have a moderately slow to moderately rapid infiltration rate. This site should show no evidence of rills, wind-scoured areas, or pedestalled plants. Water flow paths are broken, irregular in appearance, or discontinuous with numerous debris dams or vegetative barriers. The soil surface is stable and intact.

These soils are mainly susceptible to water erosion. The hazard of water erosion increases where vegetative cover is not adequate. A drastic loss of the soil surface layer on this site can result in a shift in species composition and/or production.

Major soils correlated this site include: Barnum, Clarkelen, Colombo, Craft, Draknab, Glenberg, Goshen, Haverdad, Haverson, Lomiller, Onita, Rockypoint, St.Onge, and Winetti.

More information can be found in the various soil survey reports. Contact the local USDA Service Center for soil survey reports that include more detail specific to your location.

**Table 4. Representative soil features**

Surface texture	(1) Silty clay loam (2) Loam (3) Fine sandy loam
Family particle size	(1) Loamy
Drainage class	Moderately well drained to well drained
Permeability class	Moderately slow to rapid
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	10.16–17.78 cm
Calcium carbonate equivalent (0-101.6cm)	0–25%
Electrical conductivity (0-101.6cm)	0–8 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–10
Soil reaction (1:1 water) (0-101.6cm)	6.6–9
Subsurface fragment volume <=3" (Depth not specified)	0–10%
Subsurface fragment volume >3" (Depth not specified)	0–5%

## Ecological dynamics

This site developed under Northern Great Plains climatic conditions, natural influences of large herbivores, occasional fire, and other biotic and abiotic factors that typically influence soil/site development. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plant and animal species, and management actions. While the following plant community descriptions specify more typical transitions between communities that will occur, severe disturbances, such as periods of well-below average precipitation, can cause significant shifts in plant communities and/or species composition.

Continuous grazing without adequate recovery periods following each grazing occurrence over several years causes this site to depart from the Reference Plant Community (1.1). Species such as western wheatgrass and blue grama will initially increase. Big bluestem, green needlegrass, and switchgrass will decrease in frequency and production. Heavy continuous grazing causes Kentucky bluegrass to increase and eventually develops into a sod condition. Extended periods of non-use and no fire will result in a plant community with high litter levels, which favors an increase in Kentucky bluegrass and annual bromes. In time, shrubs such as western snowberry and chokecherry will also increase. In relatively recent time, non-native tree species, principally Russian olive and saltcedar, have become very invasive in this site. Over time, and without treatment, these species can produce a closed canopy and out-compete native shrubs and trees.

The plant community upon which interpretations are primarily based is the Reference Plant Community. The Reference Plant Community has been determined by studying rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been used. Plant communities, states, transitional pathways, and thresholds have been determined through similar studies and experience.

The following is a diagram that illustrates the common plant communities that can occur on the site and the transition pathways between communities. The ecological processes are discussed in more detail in the plant community descriptions following the diagram.

## State and transition model

## Loamy Overflow R060AY020SD 7/24/17




Figure 6. Loamy Overflow - R060AY020SD.

Diagram Legend - Loamy Overflow - R060AY020SD		
T1A	Continuous season-long grazing or excessive haying, invasion and establishment of non-native, cool-season grasses.	
T1B	Flooding followed by long-term prescribed grazing that included proper stocking, change in season of use and deferment which provides time for seedling establishment, no fire, and time.	
T2A	Continuous season-long grazing or excessive haying, invasion of non-native cool-season grasses and possible non-native invasive tree species.	
T2B	Flooding followed by long-term prescribed grazing that included proper stocking, change in season of use and deferment which provides time for seedling establishment, no fire, and time.	
T3A	Invasion of non-native tree species, continuous season-long grazing, continuous seasonal grazing, and time.	
T4A	Invasion of non-native cool-season grasses, invasion on non-native tree species, continuous season-long grazing, and/or continuous seasonal grazing.	
T4B	Fire, invasion of non-native, cool-season grasses, continuous season-long grazing, continuous seasonal grazing and time.	
T5A	Fire, brush management, continuous season-long grazing, or continuous seasonal grazing.	
R3A	Prescribed grazing with proper stocking rates, change is season of use and time for adequate recovery or possibly prescribed burning followed by long-term prescribed grazing. This transition may not be rapid or feasible.	
CP 2.1A	2.1 - 2.2	No use and no fire
CP 2.2A	2.2 - 2.1	Prescribed grazing with proper stocking, change in season of use and adequate time for recovery, possibly prescribed burning followed by prescribed grazing.
CP 4.1A	4.1 - 4.2	No flooding, long-term prescribed grazing that included proper stocking, change in season of use, and deferment which provides opportunity for woody regeneration, and time.
CP 4.2A	4.2 - 4.1	Flooding, long-term prescribed grazing that included proper stocking, change in season of use, and deferment which provides opportunity for woody regeneration.
CP 4.2B	4.2 - 4.3	No flooding, long-term prescribed grazing that included proper stocking, change in season of use, and deferment which provides opportunity for woody regeneration and time.
CP 4.3A	4.3 - 4.1	Flooding, long-term prescribed grazing that included proper stocking, change in season of use, and deferment which provides opportunity for woody regeneration, and time.

Figure 7. Loamy Overflow - R060AY020SD

## State 1

### Reference State

This State represents what is believed to exist prior to European settlement. The Reference State may exist but is unlikely to function within the natural range of variability due to the spread of Kentucky bluegrass and smooth brome onto this site. In the Reference State, this site is dominated by warm-and cool- season grasses, and various shrub and tree species that are scattered across the site. Grazing or the lack of grazing, flooding or lack of flooding, fire, excessive haying, and invasion of non-native cool-season grasses are the major drivers of this State. Flooding and no fire on this site can cause a transition to a Woody Overstory State (4.0).

### Community 1.1

#### Big Bluestem-Rhizomatous Wheatgrass/Scattered Shrubs/Scattered Trees





Figure 8. Loamy Overflow - R060AY020SD - PCP 1.1

The plant community upon which interpretations are primarily based is the Big Bluestem-Rhizomatous Wheatgrass/Scattered Shrubs/Scattered Trees Plant Community (1.1). This is also considered to be the Reference Plant Community. This plant community can be found on areas that are properly managed with prescribed grazing that allows for adequate recovery periods following each grazing event. The potential vegetation is about 70 to 85 percent grasses and grass-like plants, 5 to 15 percent forbs, 5 to 10 percent shrubs, and 0 to 3 percent trees. Major grasses include big bluestem, wheatgrasses, green needlegrass, and switchgrass. Prairie sandreed may be prevalent on the more sandy textured sites. Other grasses occurring on this community include blue grama, Canada wildrye, needle and thread, and prairie Junegrass. Major forbs and shrubs include American licorice, goldenrod, purple prairie clover, silver sagebrush, western snowberry, and leadplant. Big sagebrush will be common in the western portion of the MLRA. Scattered green ash, plains cottonwood, and American elm may occur. This plant community is well adapted to the Northern Great Plains climatic conditions. Individual species can vary greatly in production depending on growing conditions (timing and amount of precipitation, and temperature). Community dynamics, nutrient cycle, water cycle, and energy flow are functioning properly. Plant litter is properly distributed with very little movement off-site and natural plant mortality is very low. The diversity in plant species allows for high drought tolerance. Run-off from adjacent sites and moderate or high available water capacity provides a favorable soil-water-plant relationship.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1939	2542	3099
Forb	151	314	504
Shrub/Vine	151	235	336
Tree	—	47	95
<b>Total</b>	<b>2241</b>	<b>3138</b>	<b>4034</b>

Figure 10. Plant community growth curve (percent production by month).  
SD6008, Pierre Shale Plains, lowland cool season/warm season co-  
dominant. Cool season, warm season co-dominant, lowland..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	4	11	19	23	20	12	6	5	0	0

## State 2 Native/Invaded State

This State has a significant amount of Kentucky bluegrass or smooth brome in the plant community but they have not become the dominate species. In this State, Kentucky bluegrass makes up less than 30 percent of the plant community composition (by weight). This State is at risk of transitioning to a bluegrass dominated Invaded State (3.0).



## Community 2.1

### Rhizomatous Wheatgrass-Kentucky Bluegrass-Remnant Big Bluestem/Scattered Trees

This plant community results from continuous season-long grazing without adequate recovery periods between each grazing event during the growing season, or excessive haying, and invasion of non- native cool-season grasses. Western wheatgrass is the dominant species, Kentucky bluegrass is subdominant. Big bluestem, little bluestem, green needlegrass, and switchgrass are greatly reduced. Forb species include western yarrow, asters, prairie coneflower, cudweed sagewort, and western ragweed. Leadplant is greatly reduced while other shrub species would tend to be heavily browsed. Scattered American elm, boxelder, cottonwood, and green ash are common on this site. Seedling and saplings can also be heavily browsed. This plant community is relatively stable and less productive than the Reference Plant Community (1.1). Reduction of litter and short plant heights result in higher soil temperatures, poor water infiltration rates, and increased runoff. This plant community can occur throughout the site, on spot grazed areas, and around water sources where season-long grazing patterns occur. Soil erosion will be minimal due to the sod forming habit of Kentucky bluegrass.

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	953	1453	1945
Forb	84	179	280
Shrub/Vine	84	135	185
Tree	—	27	56
<b>Total</b>	<b>1121</b>	<b>1794</b>	<b>2466</b>

Figure 12. Plant community growth curve (percent production by month).  
SD6007, Pierre Shale Plains, cool season dominant, warm season  
subdominant. Cool season dominant, warm season subdominant, lowland..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	13	20	25	18	11	5	3	0	0

## Community 2.2

### Rhizomatous Wheatgrass-Bluegrass/Mixed Shrub/Scattered Trees

This plant community develops after an extended period of non-use and exclusion of fire. This plant community will also develop with moderate or heavy continuous seasonal grazing. In either case, shrubs increase and can sometimes dominate the plant community. Cool-season grasses make up the majority of the understory with the balance made up of short- warm-season grasses, and miscellaneous forbs. Western wheatgrass and Kentucky bluegrass are the dominant grasses. Grasses of secondary importance include blue grama, prairie Junegrass, Sandberg bluegrass, green needlegrass, and needle and thread. Forbs commonly found in this plant community include cudweed sagewort, goldenrod, hairy goldenaster, and scurfpea. Woody plants such as silver sagebrush, and snowberry increase with canopy cover up to 20 percent. Wyoming big sagebrush will occur in the west. Scattered native trees will likely be present as will Russian olive and saltcedar. This plant community is resistant to change. Grazing in combination with brush management will reduce shrub cover if that is the management goal. Soil erosion is low. Runoff is similar to the Reference Plant Community (1.1). Once the advanced stage of this plant community is reached, time and external resources will be needed to see recovery in the diversity of the site.

Table 7. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	986	1689	2387
Shrub/Vine	241	493	729
Forb	118	247	392
Tree	—	37	78
<b>Total</b>	<b>1345</b>	<b>2466</b>	<b>3586</b>

Figure 14. Plant community growth curve (percent production by month).  
SD6008, Pierre Shale Plains, lowland cool season/warm season co-  
dominant. Cool season, warm season co-dominant, lowland..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	4	11	19	23	20	12	6	5	0	0

### Pathway 2.1A Community 2.1 to 2.2

Nonuse and no fire will move this plant community to the Rhizomatous Wheatgrass-Bluegrass/Snowberry/Scattered Trees/Excessive Litter Plant Community (2.2).

### Pathway 2.2A Community 2.2 to 2.1

Prescribed grazing that includes proper stocking, change in season of use, and adequate time for recovery, will move this plant community back to Plant Community Phase (PCP) 2.1. Brush management (reduction, not removal of shrubs) or prescribed burning followed by prescribed grazing will also move this plant community toward PCP 1.2. Either would require long-term management with prescribed grazing and/or prescribed burning under controlled conditions.

### State 3 Invaded State

This state is the result of invasion and dominance of introduced species. This state is characterized by the dominance of Kentucky bluegrass and smooth brome, and an increasing thatch layer that effectively blocks introduction of other plants into the system. Plant litter accumulation tends to favor the more shade tolerant, introduced grass species. The nutrient cycle is also impaired, and the result is typically a higher level of nitrogen which also favors the introduced species. Increasing plant litter decreases the amount of sunlight reaching plant crowns thereby shifting competitive advantage to shade tolerant, introduced grass species. Studies indicate that soil biological activity is altered, and this shift apparently exploits the soil microclimate and encourages growth of the introduced grass species. Once the threshold is crossed, a change in grazing management alone cannot cause a reduction in the invasive grass dominance. Preliminary studies would tend to indicate this threshold may exist when Kentucky bluegrass exceeds 30 percent of the plant community and native grasses represent less than 40 percent of the plant community composition. Plant communities dominated by Kentucky bluegrass have significantly less cover and diversity of native grasses and forb species (Toledo, D. et al., 2014).

### Community 3.1 Kentucky Bluegrass/Weedy Annual and Perennial Forbs

This plant community developed with heavy continuous season-long grazing, and/or excessive haying. Kentucky bluegrass dominates the community and can develop into a “sod-bound” appearance. Low vigor western wheatgrass can be found scattered throughout the community. Green needlegrass has been greatly reduced. Big bluestem may persist in minor amounts, greatly reduced in vigor and not readily seen. Western yarrow, scurfpea, ragweed, and goldenrod have increased. Non-native grasses and forbs such as annual bromes, curlycup gumweed, thistle, and cocklebur will invade this plant community. Western snowberry, American plum, and chokecherry may persist in the plant community if not removed during haying activities. Russian olive and saltcedar may be present

but in minor amounts. This plant community is resistant to change due to grazing tolerance of Kentucky bluegrass. A significant amount of production and diversity has been lost when compared to the Reference Plant Community (1.1). The dominance of non-native cool-season grasses and the loss of other desirable species has negatively impacted energy flow and nutrient cycling. Water infiltration is reduced significantly and soil loss may be accelerated where concentrated flows occur.

**Table 8. Annual production by plant type**

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	404	852	1295
Forb	106	168	230
Shrub/Vine	50	84	118
Tree	–	17	39
<b>Total</b>	<b>560</b>	<b>1121</b>	<b>1682</b>

**Figure 16. Plant community growth curve (percent production by month). SD6006, Pierre Shale Plains, lowland cool season dominant. Cool season dominant, lowland..**

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	6	15	20	26	17	9	4	3	0	0

## State 4 Woody Overstory State

This State is the result of the establishment of a tree overstory and shrub mid-story canopy. This State is more common on the lower reaches of the site. The dynamics of this State are largely due to flooding and successional changes, starting with cottonwood and shrub establishment, and eventually the development of a green ash and boxelder plant community. The successional process can restart following another flooding event. Water control structures which limit flooding, livestock grazing, heavy wildlife browse, fire, and the introduction of non-native, cool-season grasses can alter the dynamics of this site resulting in old remnant stands of trees with little, if any regeneration. Down cutting of the drainageway or stream channel will eventually leave the overflow site on a higher and drier landscape position. In this case the site will be described as the Loamy Terrace Ecological Site (R060AY022SD).

### Community 4.1 Cottonwood/Prunus-Snowberry/Forbs/Grasses

This plant community typically occurs after a flooding event. Flooding reduces herbaceous competition through scouring of the soil surface, which provides a site for regeneration and establishment of cottonwood and shrubs. Prescribed grazing, which prevents targeted grazing of cottonwood seedlings, is necessary for this plant community to establish. Trees will range from seedlings to saplings, and the herbaceous understory will still be productive as a result of the filtered canopy of the deciduous trees. Understory shrubs, primarily plum and/or chokecherry, and snowberry will likely establish. However, other species, including silver buffaloberry, silver sagebrush, big sagebrush, and currants can occur and make up a significance percentage of the shrub layer.

### Community 4.2 Mature Cottonwood-Immature Green Ash-Boxelder/Prunus-Snowberry/Forbs/Grass Understory

Mature cottonwood trees and woody shrubs provide a suitable microclimate for establishment of other deciduous trees. Green ash and boxelder are typically the first trees to establish, but other species such as hackberry and possibly bur oak will established as well. The herbaceous plant community will remain relatively productive, but will be reduced somewhat from the Reference Plant Community (1.1). This is due mainly to the competition from the woody shrub understory.

## **Community 4.3**

### **Green Ash-Boxelder/Prunus-Snowberry/Forbs/Grass Understory**

This plant community develops over time, with prescribed grazing and no flooding. Mature cottonwood trees will likely remain in lesser numbers, but the dominant trees will normally be green ash and boxelder. At times there will be a mix of all three species; however, some areas will be dominated by one or two of these species. Woody shrubs will remain in the understory, but typically in lesser amounts than in the previous two plant communities. While somewhat reduced, the herbaceous understory will remain relatively productive. The trees will mostly be in the mature stage, but regeneration will normally be evident (i.e., seedlings and saplings should be present).

## **Pathway 4.1A**

### **Community 4.1 to 4.2**

No flooding and long-term prescribed grazing that manages the herbaceous understory, but is not detrimental to woody regeneration, will allow this plant community to develop into the Mature Cottonwood-Immature Green Ash-Boxelder/Prunus-Snowberry/Forbs/Grass Understory Plant Community (4.2). Existing saplings must be large enough to avoid damage by livestock and wildlife for this pathway to occur.

## **Pathway 4.2A**

### **Community 4.2 to 4.1**

Flooding, which opens up the herbaceous understory and allows for woody regeneration, followed by long-term prescribed grazing that manages for woody regeneration and establishment will shift this plant community back to the Cottonwood/Prunus-Snowberry/Forbs/Grasses Plant Community (4.1).

## **Pathway 4.2B**

### **Community 4.2 to 4.3**

No flooding, long-term prescribed grazing that manages the herbaceous understory, but is not detrimental to woody regeneration, and time will transition this plant community to the Green Ash-Boxelder/Prunus-Snowberry/Forbs/Grass Understory Plant Community (4.3).

## **Pathway 4.3A**

### **Community 4.3 to 4.1**

Flooding, which opens up the herbaceous understory and allows for woody regeneration, followed by long-term prescribed grazing that manages for woody regeneration and establishment will shift this plant community back to the Cottonwood/Prunus-Snowberry/Forbs/Grasses Plant Community (4.1).

## **State 5**

### **Woody Invaded State**

This State developed as a result of the invasion of Russian olive and/or saltcedar in combination with continuous season-long grazing, or continuous seasonal grazing, or fire. With time, the cottonwood, boxelder and ash trees that survive become mature, and little or no regeneration occurs. This is due mainly to grazing of seedlings and saplings. Wildlife browse can also contribute to the loss of native tree and shrub regeneration. Grazing that limits regeneration also results in a reduction of the desirable native herbaceous species, often resulting in a dominance of species such as bluegrass and/or smooth brome, and forbs such as western ragweed, Canada thistle, burdock, and hound's tongue.

## **Community 5.1**

### **Scattered Mature Native Trees-Russian Olive-Saltcedar/Shrubs/Forbs/Bluegrass**



**Figure 17. Loamy Overflow - R060Y020SD - PCP 5.1**

This plant community developed due to the lack of woody regeneration, natural occurrences of flooding events, and continuous season-long grazing without adequate recovery periods. Older mature trees remain, including cottonwood, boxelder, and green ash. The trees are scattered, and the site may have a “park-like” appearance with few trees and reduced understory. If grazed during the winter, the increased durations of livestock loitering can result in manure accumulations and soil compaction which will reduce the vigor of the native understory plant community. Kentucky bluegrass and smooth brome continue to persist as dominant grass species at reduced production rates. The presence of non-desirable forb species such as Canada thistle, burdock, and hound’s tongue can be prolific and difficult to control. Russian olive and/or saltcedar will increase over time and will eventually dominate the site.

### **Transition T1A State 1 to 2**

Continuous season-long grazing and/or excessive haying and the invasion of non-native cool-season grasses will convert this plant community to the Native/Invaded State (2.0).

### **Transition T1B State 1 to 4**

T1B Flooding, followed by long-term prescribed grazing, and no fire can transition this plant community to the Woody Overstory State (4.0). This State is more likely to occur on the mid to lower reaches of a drainage.

### **Transition T2A State 2 to 3**

Continuous season-long grazing and/or excessive haying in addition to the spread of non-native cool-season grasses will result in a transition to the Invaded State (3.0). The spread of non-native invasive trees species is also a possibility.

### **Transition T2B State 2 to 4**

Flooding and no fire followed, by long-term prescribed grazing, including proper stocking, change in season of use, and adequate time for recovery will likely transition this site to the Woody Overstory State (4.0). Timed grazing is very important and must be followed for many years in order for saplings to attain a height where livestock will not damage and/or kill the trees. Wildlife browse can also be a concern if the management objective are to improve the overstory canopy. The Woody Overstory State is more likely to occur on the mid to lower reaches of a drainage.

### **Restoration pathway R3A State 3 to 2**

Long term prescribed grazing and/or prescribed burning may transition the plant community toward the Native/Invaded State (2.0). This is assuming an adequate seed/vegetative source is available. This transition may take an extended period of time and in the end not meet management objectives.

### Transition T3A State 3 to 5

Invasion of non-native tree species, continuous season-long grazing, continuous seasonal grazing and time will transition the Invaded State (3.0) to the Woody Invaded State (5.0).

### Transition T4B State 4 to 2

Fire, invasion of non-native, cool-season grasses, continuous season-long grazing, or continuous seasonal grazing resulting in little woody regeneration, and time will transition this state to the Native/Invaded State (2.0).

### Transition T4A State 4 to 5

Invasion of non-native tree species in combination with continuous season-long grazing, continuous seasonal grazing, heavy wildlife browse, or fire can cause any of the plant communities in the Woody Overstory State (4.0) to transition to the Woody Invaded State (5.0).

### Transition T5A State 5 to 3

Fire, brush management to remove Russian olive and salt cedar, continuous season-long grazing, or continuous seasonal grazing will transition this plant community to the Invaded State (3.0).

## Additional community tables

Table 9. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Tall Warm-Season Grasses</b>			785–1412	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	628–1255	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	157–628	–
2	<b>Wheatgrasses</b>			628–942	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	628–942	–
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	157–314	–
	slender wheatgrass	ELTRT	<i>Elymus trachycaulus ssp. trachycaulus</i>	157–314	–
3	<b>Needlegrasses</b>			157–314	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	157–314	–
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	63–157	–
4	<b>Native Grasses/Grass-Likes</b>			157–471	
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	63–251	–
	sedge	CAREX	<i>Carex</i>	63–251	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	31–157	–
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–157	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–157	–

	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0–94	–
	prairie sandreed	CALO	<i>Calamovilfa longifolia</i>	0–94	–
	dropseed	SPORO	<i>Sporobolus</i>	0–94	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–63	–
	spikerush	ELEOC	<i>Eleocharis</i>	0–63	–
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	0–63	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	31–63	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–31	–
	saltgrass	DISP	<i>Distichlis spicata</i>	0–31	–

#### Forb

6	<b>Forbs</b>			157–471	
	goldenrod	SOLID	<i>Solidago</i>	31–157	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	31–126	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	31–94	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–94	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	31–63	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	31–63	–
	American vetch	VIAM	<i>Vicia americana</i>	0–63	–
	blazing star	LIATR	<i>Liatris</i>	0–63	–
	mint	MENTH	<i>Mentha</i>	0–63	–
	Canadian anemone	ANCA8	<i>Anemone canadensis</i>	0–63	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	31–63	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	31–63	–
	milkvetch	ASTRA	<i>Astragalus</i>	0–31	–
	false boneset	BREU	<i>Brickellia eupatorioides</i>	0–31	–
	thistle	CIRSI	<i>Cirsium</i>	0–31	–
	scurfpea	PSORA2	<i>Psoralegium</i>	0–31	–
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	0–31	–
	blue-eyed grass	SISYR	<i>Sisyrinchium</i>	0–31	–
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	0–31	–

#### Shrub/Vine

7	<b>Shrubs</b>			157–314	
	Subshrub (<.5m)	2SUBS	<i>Subshrub (&lt;.5m)</i>	0–157	–
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	0–157	–
	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	0–157	–
	American plum	PRAM	<i>Prunus americana</i>	0–157	–
	chokecherry	PRVI	<i>Prunus virginiana</i>	0–157	–
	silver buffaloberry	SHAR	<i>Shepherdia argentea</i>	0–157	–
	snowberry	SYMPH	<i>Symphoricarpos</i>	31–157	–
	leadplant	AMCA6	<i>Amorpha canescens</i>	31–126	–
	currant	RIBES	<i>Ribes</i>	0–94	–
	rose	ROSA5	<i>Rosa</i>	31–94	–
	willow	SALIX	<i>Salix</i>	0–94	–



	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0–63	–
<b>Tree</b>					
8	<b>Trees</b>			0–94	
	Tree	2TREE	<i>Tree</i>	0–94	–
	boxelder	ACNE2	<i>Acer negundo</i>	0–94	–
	common hackberry	CEOC	<i>Celtis occidentalis</i>	0–94	–
	green ash	FRPE	<i>Fraxinus pennsylvanica</i>	0–94	–
	plains cottonwood	PODEM	<i>Populus deltoides ssp. monilifera</i>	0–94	–
	American elm	ULAM	<i>Ulmus americana</i>	0–94	–

Table 10. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Tall Warm-Season Grasses</b>			90–269	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	90–269	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	18–90	–
2	<b>Wheatgrasses</b>			448–628	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	359–628	–
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	36–90	–
	slender wheatgrass	ELTRT	<i>Elymus trachycaulus ssp. trachycaulus</i>	36–90	–
3	<b>Needlegrasses</b>			18–90	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	18–90	–
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	0–54	–
4	<b>Native Grasses/Grass-Likes</b>			90–179	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–90	–
	sedge	CAREX	<i>Carex</i>	18–90	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	18–90	–
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	18–90	–
	dropseed	SPORO	<i>Sporobolus</i>	18–90	–
	saltgrass	DISP	<i>Distichlis spicata</i>	18–54	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	18–54	–
	prairie sandreed	CALO	<i>Calamovilfa longifolia</i>	18–54	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–36	–
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0–36	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–36	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–36	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–18	–
5	<b>Non-Native Grasses</b>			269–448	
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	269–448	–
	cheatgrass	BRTE	<i>Bromus tectorum</i>	18–179	–
	smooth brome	BRIN2	<i>Bromus inermis</i>	0–90	–
<b>Forb</b>					

0	Forbs			90-209	
	cocklebur	XANTH2	<i>Xanthium</i>	0-90	-
	goldenrod	SOLID	<i>Solidago</i>	18-90	-
	white heath aster	SYER	<i>Symphotrichum ericoides</i>	36-90	-
	Forb, annual	2FA	<i>Forb, annual</i>	18-90	-
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	36-90	-
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	36-90	-
	thistle	CIRSI	<i>Cirsium</i>	0-90	-
	mint	MENTH	<i>Mentha</i>	0-90	-
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	0-72	-
	common yarrow	ACMI2	<i>Achillea millefolium</i>	36-72	-
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	18-54	-
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-54	-
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	18-36	-
	milkvetch	ASTRA	<i>Astragalus</i>	0-36	-
	scurfpea	PSORA2	<i>Psoraleidum</i>	18-36	-
	blazing star	LIATR	<i>Liatris</i>	0-36	-
	curlycup gumweed	GRSQ	<i>Grindelia squarrosa</i>	0-18	-
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0-18	-
	blue-eyed grass	SISYR	<i>Sisyrinchium</i>	0-18	-
	false boneset	BREU	<i>Brickellia eupatorioides</i>	0-18	-
	Canadian anemone	ANCA8	<i>Anemone canadensis</i>	0-18	-
	American vetch	VIAM	<i>Vicia americana</i>	0-18	-
<b>Shrub/Vine</b>					
7	<b>Shrubs</b>			90-179	
	Subshrub (<.5m)	2SUBS	<i>Subshrub (&lt;.5m)</i>	0-90	-
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	0-90	-
	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	0-90	-
	American plum	PRAM	<i>Prunus americana</i>	0-90	-
	chokecherry	PRVI	<i>Prunus virginiana</i>	0-90	-
	silver buffaloberry	SHAR	<i>Shepherdia argentea</i>	0-90	-
	snowberry	SYMPH	<i>Symphoricarpos</i>	18-90	-
	currant	RIBES	<i>Ribes</i>	0-54	-
	rose	ROSA5	<i>Rosa</i>	18-54	-
	leadplant	AMCA6	<i>Amorpha canescens</i>	18-54	-
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0-36	-
	willow	SALIX	<i>Salix</i>	0-36	-
<b>Tree</b>					
8	<b>Trees</b>			0-54	
	Tree	2TREE	<i>Tree</i>	0-54	-
	boxelder	ACNE2	<i>Acer negundo</i>	0-54	-
	common hackberry	CEOC	<i>Celtis occidentalis</i>	0-54	-
	green ash	FRPE	<i>Fraxinus pennsylvanica</i>	0-54	-
	slender cottonwood	POPEM	<i>Populus deltoides</i> var. <i>monilifera</i>	0-54	-

	plains cottonwood	PODEM	<i>Populus deltoides ssp. monilifera</i>	0–54	–
	American elm	ULAM	<i>Ulmus americana</i>	0–54	–

Table 11. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Tall Warm-Season Grasses</b>			49–247	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	25–197	–
	switchgrass	PAVI2	<i>Panicum virgatum</i>	25–123	–
2	<b>Wheatgrasses</b>			247–493	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	247–493	–
	thickspike wheatgrass	ELLAL	<i>Elymus lanceolatus ssp. lanceolatus</i>	0–123	–
	slender wheatgrass	ELTRT	<i>Elymus trachycaulus ssp. trachycaulus</i>	0–74	–
3	<b>Needlegrasses</b>			25–123	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	25–123	–
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	0–74	–
4	<b>Native Grasses/Grass-Likes</b>			123–370	
	dropseed	SPORO	<i>Sporobolus</i>	49–197	–
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	0–123	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	25–123	–
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–123	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	25–123	–
	prairie sandreed	CALO	<i>Calamovilfa longifolia</i>	25–123	–
	sedge	CAREX	<i>Carex</i>	49–123	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–49	–
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0–49	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–49	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–49	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–49	–
	spikerush	ELEOC	<i>Eleocharis</i>	0–25	–
	saltgrass	DISP	<i>Distichlis spicata</i>	0–25	–
5	<b>Non-Native Grasses</b>			370–493	
	smooth brome	BRIN2	<i>Bromus inermis</i>	0–247	–
	cheatgrass	BRTE	<i>Bromus tectorum</i>	25–247	–
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	0–247	–
<b>Forb</b>					
6	<b>Forbs</b>			123–370	
	Forb, annual	2FA	<i>Forb, annual</i>	49–247	–
	goldenrod	SOLID	<i>Solidago</i>	49–197	–
	mint	MENTH	<i>Mentha</i>	0–123	–
	scurfpea	PSORA2	<i>Psoralegium</i>	25–123	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	49–123	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	49–123	–

	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	49–123	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	25–123	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	25–123	–
	thistle	CIRSI	<i>Cirsium</i>	0–74	–
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	0–74	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	25–74	–
	American vetch	VIAM	<i>Vicia americana</i>	0–49	–
	cocklebur	XANTH2	<i>Xanthium</i>	0–49	–
	false boneset	BREU	<i>Brickellia eupatorioides</i>	0–49	–
	blazing star	LIATR	<i>Liatris</i>	0–49	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	0–25	–
	Canadian anemone	ANCA8	<i>Anemone canadensis</i>	0–25	–
	blue-eyed grass	SISYR	<i>Sisyrinchium</i>	0–25	–
	curlycup gumweed	GRSQ	<i>Grindelia squarrosa</i>	0–25	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0–25	–
<b>Shrub/Vine</b>					
7	<b>Shrubs</b>			247–740	
	snowberry	SYMPH	<i>Symphoricarpos</i>	49–370	–
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	49–370	–
	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	49–370	–
	American plum	PRAM	<i>Prunus americana</i>	25–123	–
	chokecherry	PRVI	<i>Prunus virginiana</i>	25–123	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0–123	–
	Subshrub (<.5m)	2SUBS	<i>Subshrub (&lt;.5m)</i>	25–123	–
	willow	SALIX	<i>Salix</i>	25–123	–
	silver buffaloberry	SHAR	<i>Shepherdia argentea</i>	0–123	–
	currant	RIBES	<i>Ribes</i>	0–74	–
	rose	ROSA5	<i>Rosa</i>	25–74	–
	leadplant	AMCA6	<i>Amorpha canescens</i>	0–49	–
<b>Tree</b>					
8	<b>Trees</b>			0–74	
	Tree	2TREE	<i>Tree</i>	0–74	–
	boxelder	ACNE2	<i>Acer negundo</i>	0–74	–
	common hackberry	CEOC	<i>Celtis occidentalis</i>	0–74	–
	green ash	FRPE	<i>Fraxinus pennsylvanica</i>	0–74	–
	plains cottonwood	PODEM	<i>Populus deltoides ssp. monilifera</i>	0–74	–
	American elm	ULAM	<i>Ulmus americana</i>	0–74	–

Table 12. Community 3.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Tall Warm-Season Grasses</b>			0–56	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	0–56	–

	switchgrass	PAVI2	<i>Panicum virgatum</i>	0–22	–
2	<b>Wheatgrasses</b>			56–112	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	56–112	–
3	<b>Needlegrasses</b>			0–56	
	green needlegrass	NAVI4	<i>Nassella viridula</i>	0–56	–
	needle and thread	HECOC8	<i>Hesperostipa comata</i> ssp. <i>comata</i>	0–34	–
4	<b>Native Grasses/Grass-Likes</b>			56–168	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–112	–
	dropseed	SPORO	<i>Sporobolus</i>	22–112	–
	sedge	CAREX	<i>Carex</i>	11–56	–
	saltgrass	DISP	<i>Distichlis spicata</i>	11–56	–
	foxtail barley	HOJU	<i>Hordeum jubatum</i>	22–56	–
	prairie sandreed	CALO	<i>Calamovilfa longifolia</i>	11–34	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–22	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–22	–
5	<b>Non-Native Grasses</b>			392–560	
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	392–560	–
	smooth brome	BRIN2	<i>Bromus inermis</i>	0–112	–
	cheatgrass	BRTE	<i>Bromus tectorum</i>	22–112	–
<b>Forb</b>					
6	<b>Forbs</b>			112–224	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	56–168	–
	thistle	CIRSI	<i>Cirsium</i>	0–112	–
	curlycup gumweed	GRSQ	<i>Grindelia squarrosa</i>	0–112	–
	cocklebur	XANTH2	<i>Xanthium</i>	0–90	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	22–56	–
	goldenrod	SOLID	<i>Solidago</i>	22–56	–
	white heath aster	SYER	<i>Symphyotrichum ericoides</i>	22–56	–
	mint	MENTH	<i>Mentha</i>	0–56	–
	scurfpea	PSORA2	<i>Psoralegium</i>	22–56	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	22–56	–
	Forb, annual	2FA	<i>Forb, annual</i>	22–56	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	11–34	–
	upright prairie coneflower	RACO3	<i>Ratibida columnifera</i>	11–34	–
	blazing star	LIATR	<i>Liatris</i>	0–11	–
<b>Shrub/Vine</b>					
7	<b>Shrubs</b>			56–112	
	silver sagebrush	ARCA13	<i>Artemisia cana</i>	0–56	–
	snowberry	SYMPH	<i>Symphoricarpos</i>	11–56	–
	rose	ROSA5	<i>Rosa</i>	11–34	–
	Subshrub (<.5m)	2SUBS	<i>Subshrub (&lt;.5m)</i>	0–22	–
	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0–22	–

	big sagebrush	ARTR2	<i>Artemisia tridentata</i>	0–11	–
	chokecherry	PRVI	<i>Prunus virginiana</i>	0–11	–
	willow	SALIX	<i>Salix</i>	0–11	–
	silver buffaloberry	SHAR	<i>Shepherdia argentea</i>	0–11	–
<b>Tree</b>					
8	<b>Trees</b>			0–34	
	Tree	2TREE	<i>Tree</i>	0–34	–
	boxelder	ACNE2	<i>Acer negundo</i>	0–34	–
	common hackberry	CEOC	<i>Celtis occidentalis</i>	0–34	–
	green ash	FRPE	<i>Fraxinus pennsylvanica</i>	0–34	–
	plains cottonwood	PODEM	<i>Populus deltoides ssp. monilifera</i>	0–34	–
	American elm	ULAM	<i>Ulmus americana</i>	0–34	–

## Animal community

The following table lists annual suggested initial stocking rates with average growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of conservation planning. Often, the current plant composition does not entirely match any particular plant community (as described in this Ecological Site Description). Therefore, a resource inventory is necessary to document plant composition and production. More accurate carrying capacity estimates should eventually be calculated using the following stocking rate information along with animal preference data and actual stocking records, particularly when grazers other than cattle are involved. With consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity.

Plant Community = Big Bluestem-Rhizomatous Wheatgrass/Scattered Shrubs/Scattered Trees (1.1)

Average Annual Production (lbs./ac, air-dry) = 2800

Stocking Rate (AUM/ac) = 0.77

Plant Community = Rhizomatous Wheatgrass-Kentucky Bluegrass-Remnant Big Bluestem/Scattered Trees (2.1)

Average Annual Production (lbs./ac, air-dry) = 1600

Stocking Rate (AUM/ac) = 0.44

Plant Community = Rhizomatous Wheatgrass-Bluegrass/Mixed Shrubs/Scattered Trees (2.2)

Average Annual Production (lbs./ac, air-dry) = 2200

Stocking Rate (AUM/ac) = 0.60\*

Plant Community = Kentucky Bluegrass/Weedy Annuals and Perennial Forbs (3.1)

Average Annual Production (lbs./ac, air-dry) = 1000

Stocking Rate (AUM/ac) = 0.27\*

Based on 912 lbs./acre (air-dry weight) per Animal Unit Month (AUM), and on 25 percent harvest efficiency of preferred and desirable forage species (refer to USDA NRCS, National Range and Pasture Handbook).

\* Highly variable production, stocking rates needs to be determined based from on-site production data.

Total annual production on-site may contain vegetation deemed undesirable or untargeted by the grazing animal. Therefore, AUM values may have been reduced to reflect only preferred or desirable forage species.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage. During the dormant period, the forage for livestock will likely be lacking protein to meet livestock requirements, and added protein will allow ruminants to better utilize the energy stored in grazed plant materials. A forage quality test (either directly or through fecal sampling) should be used to determine the level of supplementation needed.

## Hydrological functions

Water is the principal factor limiting herbage production on this site. The site is dominated by soils in hydrologic groups B and C. Infiltration and runoff potential for this site varies from moderate to high depending on soil hydrologic group, slope, and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An exception would be where shortgrasses form a dense sod and dominate the site. Normally areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

## Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an aesthetic value that appeals to visitors.

## Other products

Seed harvest of native plant species can provide additional income on this site.

## Other information

Revision Notes: "Previously Approved" Provisional

This Provisional ecological site concept has passed Quality Control (QC) and Quality Assurance (QA) to ensure that the site meets the 2014 NESH standards for a Provisional ecological site. This is an updated "Previously Approved" ESD which represents a first-generation tier of documentation that, prior to the release of the 2014 National Ecological Site Handbook (NESH), met all requirements as an "Approved" ESD as laid out in the 1997, rev.1, 2003 National Range and Pasture Handbook (NRPH). The document fully described the Reference State and Community Phase in the State-and-Transition model. All other alternative states are at least described in narrative form. The "Previously Approved" ESD has been field-tested for a minimum of five years and is a proven functional document for conservation planning. The "Previously Approved" ESD does not contain all tabular and narrative entries as required in the current "Approved" level of documentation but it is expected that the "Previously Approved" ESD will continue refinement towards an "Approved" status.

Site Development and Testing Plan:

Future work, as described in a Project Plan, is needed to validate the information in this Provisional Ecological Site Description. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. The final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document.

### Non-discrimination Statement

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, available online and at any USDA office, or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632- 9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for



Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

## **Inventory data references**

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site description include: Stan Boltz, Range Management Specialist, NRCS; Darrel DuVall, Range Management Specialist, NRCS; Jill Epley, Range Management Specialist, NRCS; Cheryl Nielsen, Range Management Specialist, NRCS; Rick Peterson, Range Management Specialist, NRCS; and Mike Stirling, Range Management Specialist, NRCS.

## **Other references**

EPA – Level III and Level IV Ecoregions of the Continental United States, (<https://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>). Available online. Accessed 01/03/17

High Plains Regional Climate Center, University of Nebraska, (<http://www.hprcc.unl.edu/>). Available online. Accessed 02/27/17.

Teledo, D., Sanderson, M., Spaeth, K., Hendrickson, J., Printz, J. 2014. Extent of Kentucky Bluegrass and Its Effect on Native Plant Species Diversity and Ecosystem Services in the Northern Great Plains of the United States. *Invasive Plant Science and Management*. 7(4):543-522. Weed Science Society of America.

USDA, NRCS. Soil Survey Staff. Official Soil Series Descriptions. Available online. Accessed 07/17/17.

USDA, NRCS. Soil Survey Staff. Web Soil Survey. Available online. Accessed 07/17/17.

USDA, NRCS. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. Ag Handbook 296.

USDA, NRCS. 2014. National Ecological Site Handbook, 1st Ed.

USDA, NRCS. National Water and Climate Center. (<http://www.wcc.nrcs.usda.gov/>). Available online. Accessed 02/27/17.

USDA, NRCS. 1997, rev. 1, 2003. National Range and Pasture Handbook.

USDA, NRCS. National Soil Information System, Information Technology Center. (<http://nasis.nrcs.usda.gov>).

USDA, NRCS. 2017. The PLANTS Database (<http://plants.usda.gov>, 2017). Available online. Accessed 7/14/17.

USDA, NRCS. Various Published Soil Surveys.

## **Contributors**

Stan C. Boltz

Rick L. Peterson

## **Acknowledgments**

ESD updated by Rick L. Peterson on 7/24/17.

## **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community

cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Boltz, Ryan Beer, Mitch Iverson, Thad Berrett, Cheryl Nielsen
Contact for lead author	stanley.boltz@sd.usda.gov, 605-352-1236
Date	06/04/2008
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** None.
- 

2. **Presence of water flow patterns:** Typically none or barely visible. Evidence of water flow may be present after high overland flow events or flooding from adjacent streams, but vegetation normally remains intact.
- 

3. **Number and height of erosional pedestals or terracettes:** None.
- 

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0 to 5 percent is typical.
- 

5. **Number of gullies and erosion associated with gullies:** None typical, however limited headcutting may form after high runoff or flooding events. Existing gullies should be stabilized with good vegetative cover.
- 

6. **Extent of wind scoured, blowouts and/or depositional areas:** None typical, but limited deposition may occur after major runoff or flooding events.
- 

7. **Amount of litter movement (describe size and distance expected to travel):** Litter of small and medium size classes will move after average to high rainfall events. Litter does not travel far, typically being trapped in small bunches by the extensive vegetative cover. Litter movement may be fairly extensive after major runoff or flooding events.
- 

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil aggregate stability ratings should typically be 5 to 6, normally 6. Surface organic matter adheres to the soil surface. Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.
- 

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon should be 5 to 20 inches thick with mollic (dark) colors when moist. Structure typically is medium to fine granular in the upper A-horizon.

- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None – when dry, B horizons can be hard and appear to be compacted, but no platy structure will be present.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Tall warm-season rhizomatous >
- Sub-dominant: Mid cool-season rhizomatous >> mid/tall cool-season bunchgrasses > forbs >
- Other: Shrubs = short cool-season grasses/grass-likes
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little evidence of decadence or mortality. Bunch grasses have strong, healthy centers and shrubs are vigorous.
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Production ranges from 2,000-3,600 lbs./acre (air-dry weight). Reference value production is 2,800 lbs./acre (air-dry weight).
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious weeds, Kentucky bluegrass, snowberry
- 
17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.
-