


# **Ecological site R064XY046NE** **Thin Claypan**

Accessed: 05/18/2024

## **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



**Figure 1. Mapped extent**

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

## **Classification relationships**

Level IV Ecoregions of the Conterminous United States: 25a – Pine Ridge Escarpment, 43h – White River Badlands, and 43i – Keya Paha Tablelands.

## **Associated sites**

R064XY014NE	<b>Clayey 14-17" PZ</b>
R064XY035NE	<b>Clayey 17-20 PZ</b>
R064XY044NE	<b>Claypan</b>
R064XY045NE	<b>Dense Clay</b>

## **Similar sites**

R064XY014NE	<b>Clayey 14-17" PZ</b> [More green needlegrass; higher production.]
R064XY035NE	<b>Clayey 17-20 PZ</b> [More green needlegrass; higher production.]

R064XY044NE	<b>Claypan</b> [More production; more western wheatgrass and green needlegrass.]
-------------	---

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Pascopyrum smithii</i> (2) <i>Bouteloua gracilis</i>

## Physiographic features

This site occurs on nearly level to gently undulating or rolling sedimentary uplands.

**Table 2. Representative physiographic features**

Landforms	(1) Alluvial fan (2) Alluvial flat (3) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	884–1,219 m
Slope	0–6%
Water table depth	122–203 cm
Aspect	Aspect is not a significant factor

## Climatic features

MLRA 64 is considered to have a continental climate – cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the northern Great Plains and air masses move freely across the plains and account for rapid changes in temperature.

Annual precipitation ranges from 14 to 20 inches per year. The normal average annual temperature is about 47° F. January is the coldest month with average temperatures ranging from about 21° F (Wood, SD) to about 25° F (Hemingford, NE). July is the warmest month with temperatures averaging from about 70° F (Keeline 3 W, WY) to about 76° F (Wood, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 55° F. This large annual range attests to the continental nature of this area's climate. Hourly winds average about 11 miles per hour annually, ranging from about 13 miles per hour during the spring to about 10 miles per hour during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour.

Growth of cool season plants begins in early to mid March, slowing or ceasing in late June. Warm season plants begin growth about mid May and continue to early or mid September. Green up of cool season plants may occur in September and October when adequate soil moisture is present.

**Table 3. Representative climatic features**

Frost-free period (average)	143 days
Freeze-free period (average)	163 days
Precipitation total (average)	508 mm

## Influencing water features

No riparian areas or wetland features are directly associated with this site.

## Soil features

The common features of soils in this site are the very fine sandy loam to clay textured subsoils and slopes of 0 to 6 percent. The soils in this site are moderately well to well drained and formed in alluvium or residuum derived from soft sandstone, siltstone or shale. The silt loam to loam surface layer is 1 to 4 inches thick. The extremely hard clayey Btn horizon has round-topped or “bun shaped” columnar or prismatic structured subsoil. These Btn horizons are high in sodium. The soils have a moderate to slow infiltration rate and very slow saturated hydraulic conductivity. Wet surface compaction can occur with heavy traffic. This site should show slight to no evidence of rills, wind scoured areas or pedestalled plants. Water flow paths are broken, irregular in appearance or discontinuous with numerous debris dams or vegetative barriers. The soil surface is stable and intact. These soils are mainly susceptible to water erosion. The hazard of water erosion increases on slopes greater than about 4 percent. Loss of 30 percent or more of the surface layer of the soils on this site can result in a shift in species composition and/or production.

More information can be found in the various soil survey reports. Contact the local USDA Service Center for soil survey reports that include more detail specific to your location.

**Table 4. Representative soil features**

Surface texture	(1) Silt loam (2) Loam (3) Very fine sandy loam
Family particle size	(1) Clayey
Drainage class	Moderately well drained to well drained
Permeability class	Very slow
Soil depth	51–203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	10.16–15.24 cm
Calcium carbonate equivalent (0-101.6cm)	0–25%
Electrical conductivity (0-101.6cm)	0–30 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–30
Soil reaction (1:1 water) (0-101.6cm)	5.6–9.6
Subsurface fragment volume <=3" (Depth not specified)	0–20%
Subsurface fragment volume >3" (Depth not specified)	0–10%

## Ecological dynamics

This site developed under Northern Great Plains climatic conditions, light to severe grazing by bison and other large herbivores, sporadic natural or man-caused wildfire (often of light intensities), and other biotic and abiotic factors that typically influence soil/site development. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plant and animal species, and management actions. While the following plant community descriptions describe more typical transitions between communities that will occur, severe disturbances, such as periods of well-below average precipitation, can cause significant shifts in plant communities


and/or species composition.

In association with this site are also areas of slick spots that usually have considerably more bare ground, and are typically dominated by cactus. Slick spots are bare ground areas that are affected by high sodium concentrations. The soil factors are the dominant influence and grazing management is not necessarily the primary influence of these areas. These areas can occur as a complex with this site, sometimes being difficult to differentiate between the two.

Interpretations are primarily based on the Western Wheatgrass/Blue Grama Plant Community. It has been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been used. Plant communities, states, transitional pathways, and thresholds have been determined through similar studies and experience.

The following diagram illustrates the common plant communities and vegetation states commonly occurring on the site and the transition pathways between communities and states. The ecological processes will be discussed in more detail in the plant community descriptions following the diagram.

## **State and transition model**



**CHG** – Continuous heavy grazing (heavy levels of grazing of a unit during most or all of the growing season); **PG** – Prescribed grazing (planned, controlled harvest of vegetation with grazing or browsing animals – see FOTG, Section IV, 528).

## **State 1**

### **Western Wheatgrass/Blue Grama**

## **State 2**

### **Blue Grama/Cactus**

#### **Hydrological functions**

Water is the principal factor limiting herbage production on this site. The site is dominated by soils in hydrologic group D. Infiltration varies from moderate to very slow and runoff potential varies from medium to very high depending on soil hydrologic group, slope and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An exception would be where a dense sod of short grasses dominates the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook).

#### **Recreational uses**

This site provides hunting opportunities for upland game species. The wide varieties of plants, which bloom from spring until fall, have an esthetic value that appeals to visitors.

#### **Wood products**

No appreciable wood products are present on the site.

#### **Other products**

Seed harvest of native plant species can provide additional income on this site.

#### **Inventory data references**

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS; Jill Epley, Range Management Specialist, NRCS; Rick Peterson, Range Management Specialist, NRCS; David Steffen, Range Management Specialist, NRCS; Jeff Vander Wilt, Range Management Specialist, NRCS; Phil Young, Soil Scientist, NRCS.

#### **Other references**

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://hpccsun.unl.edu>)

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224. (<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS, Various Published Soil Surveys.

#### **Contributors**

SCB

#### **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem

condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Boltz
Contact for lead author	Stan Boltz, stanley.boltz@sd.usda.gov, 605-352-1236
Date	01/05/2010
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** None.

- 
2. **Presence of water flow patterns:** Broken or irregular in appearance or discontinuous with numerous debris dams.

- 
3. **Number and height of erosional pedestals or terracettes:** Pedestals are somewhat common, but few exposed roots would occur.

- 
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 5 to 20 percent is typical; this does not include associated slickspots that are not a soil/ecological site.

- 
5. **Number of gullies and erosion associated with gullies:** None should be present.

- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.

- 
7. **Amount of litter movement (describe size and distance expected to travel):** Small size litter classes will generally move short distances, some medium size class litter will move very short distances. Litter debris dams are occasionally present.

- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil aggregate stability ratings should typically be 3 or greater. Surface organic matter adheres to the soil surface in most cases. Soil surface fragments will typically retain structure for short periods when dipped in distilled water. Some fragments will dissolve in less than 1 minute.

- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon not present at the surface, but has light colored E-horizon 1 to 4 inches thick. Structure is thin platy parting to fine

granular.

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool-season grasses) with fine and coarse roots positively influences infiltration.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None – natural pan appears at roughly 1 to 4 inches with “biscuit-top” appearance at top of pan.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Mid cool-season rhizomatous grasses >
- Sub-dominant: Short warm-season grasses > Cool-season bunchgrasses >
- Other: Forbs = Shrubs = Grass-like species
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little evidence of decadence or mortality. Bunch grasses have strong, healthy centers and shrubs are vigorous.
- 
14. **Average percent litter cover (%) and depth ( in):** Litter cover typically 20 to 50 percent, with the depth roughly 0.25 inches.
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Total annual production ranges from 500 to 1,500 pounds/acre, with the reference value being 1,000 pounds/acre (air-dry basis).
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious weeds.
- 
17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.



---