


# Ecological site R064XY050NE Thin Breaks

Accessed: 05/06/2024

## General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



**Figure 1. Mapped extent**

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

## Classification relationships

Level IV Ecoregions of the Conterminous United States: 25a – Pine Ridge Escarpment, 43h – White River Badlands, and 43i – Keya Paha Tablelands.

## Associated sites

R064XY026NE	<b>Loamy Overflow</b>
R064XY027NE	<b>Clayey Overflow</b>
R064XY037NE	<b>Thin Upland</b>
R064XY047NE	<b>Very Shallow</b>

## Similar sites

R064XY026NE	<b>Loamy Overflow</b> [Lesser shrub component; green muhly not dominant.]
-------------	--

**Table 1. Dominant plant species**

Tree	Not specified
------	---------------

Shrub	Not specified
Herbaceous	(1) <i>Muhlenbergia racemosa</i> (2) <i>Andropogon gerardii</i>

## Physiographic features

This site occurs on steep side slopes of hills.

**Table 2. Representative physiographic features**

Landforms	(1) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	884–1,219 m
Slope	30–80%
Aspect	Aspect is not a significant factor

## Climatic features

MLRA 64 is considered to have a continental climate – cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the northern Great Plains and air masses move freely across the plains and account for rapid changes in temperature.

Annual precipitation ranges from 14 to 20 inches per year. The normal average annual temperature is about 47° F. January is the coldest month with average temperatures ranging from about 21° F (Wood, SD) to about 25° F (Hemingford, NE). July is the warmest month with temperatures averaging from about 70° F (Keeline 3 W, WY) to about 76° F (Wood, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 55° F. This large annual range attests to the continental nature of this area's climate. Hourly winds average about 11 miles per hour annually, ranging from about 13 miles per hour during the spring to about 10 miles per hour during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour.

Growth of cool season plants begins in early to mid March, slowing or ceasing in late June. Warm season plants begin growth about mid May and continue to early or mid September. Green up of cool season plants may occur in September and October when adequate soil moisture is present.

**Table 3. Representative climatic features**

Frost-free period (average)	143 days
Freeze-free period (average)	163 days
Precipitation total (average)	508 mm

## Influencing water features

No riparian areas or wetland features are directly associated with this site.

## Soil features

The soils in this have very fine sandy loam to silt loam textured surface soils and slopes of 30 to 80 percent. The soils in this site are well to somewhat excessively drained and formed in soft siltstone or sandstone. The very fine sandy loam to silt loam surface layer is 2 to 10 inches thick. The soils have a moderate infiltration rate. This site can show slight to moderate evidence of rills and pedestalled plants. Water flow paths are broken, irregular in appearance or discontinuous with few debris dams or vegetative barriers. The soil surface is relatively stable,

however natural erosion is not uncommon on steeper slopes and in areas with sparser vegetation. Sub-surface soil layers are variably restrictive to water movement and root penetration.

These soils are mainly susceptible to water erosion. The hazard of water erosion increases on slopes greater than about 40 percent and on areas with sparse vegetation. The soil-water-plant relationship is strongly influenced by the cooler exposures, and occasional additional moisture supplied by water seepage and springs emanating from rock fissures and soil/rock exposures.

More information can be found in the various soil survey reports. Contact the local USDA Service Center for soil survey reports that include more detail specific to your location.

**Table 4. Representative soil features**

Surface texture	(1) Loam (2) Silt loam (3) Very fine sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderate to moderately rapid
Soil depth	25–102 cm
Surface fragment cover <=3"	0–10%
Surface fragment cover >3"	0–20%
Available water capacity (0-101.6cm)	5.08–12.7 cm
Calcium carbonate equivalent (0-101.6cm)	0–25%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	6.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	5–45%
Subsurface fragment volume >3" (Depth not specified)	5–25%

## Ecological dynamics

This site developed under Northern Great Plains climatic conditions, light to severe grazing/browsing by cervids and other large herbivores, sporadic natural or man-caused wildfire (often of light intensities), and other biotic and abiotic factors that typically influence soil/site development. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plant and animal species, and management actions. While the following plant community description describes more typical situations, severe disturbances, such as periods of well-below average precipitation, can cause significant shifts in species composition.

The plant community upon which interpretations are primarily based is the Green Muhly/Bluestem, Shrubs, Trees Plant Community. It has been determined by study of rangeland relic areas which is apparently the current condition of most of where this site occurs.


This site evidently played an important role in the pre-European development of associated woody draw plant communities. During favorable climatic conditions, and the lack of fire in a given area, the woody species which dominate this site would tend to expand into the more favorable soils of the associated overflow sites. With extended dry periods or increased fire activity, the woody species would tend to be eliminated or greatly reduced on

overflow sites, but the Thin Breaks site often acts as a refugium for many woody species. It is thought that the current extent of woody draws in the associated overflow sites is largely due to fire suppression efforts post-settlement.

Apparently, this site continues to develop largely through natural climatic cycles, and as a result of plant species that can tolerate cooler conditions and natural plant decadence and mortality. As this site was not previously described, and little information has been collected, only the Green Muhly/Bluestem, Shrubs, Trees Plant Community is characterized at this time. Therefore, no plant community diagram is included here. As more information is collected, and the site studied further, more plant communities may be identified and described.

## State and transition model

### Ecosystem states



**State 1**  
**Green Muhly/Bluestem, Shrubs, Trees**

**State 2**  
**Clayey PCs**

## Hydrological functions

Water is the principal factor limiting herbage production on this site. The site is dominated by soils in hydrologic group D. Infiltration varies from moderately to high and runoff varies from low to medium depending on slope and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

## Recreational uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

## Wood products

Local or individual firewood can be utilized from this site. This site has low potential for timber harvest due to steep slopes and variability in site production.

## Other products

Seed harvest of native plant species can provide additional income on this site.

## Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS; Jill Epley, Range Management Specialist, NRCS; Rick Peterson, Range Management Specialist, NRCS; David Steffen, Range Management Specialist, NRCS; Jeff Vander Wilt; Range Management Specialist, NRCS; Phil Young, Soil Scientist, NRCS.

Data Source Number of Records Sample Period State County

## Other references

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728.  
(<http://hpccsun.unl.edu>)

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224.  
(<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS, Various Published Soil Surveys.

## Contributors

SCB

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Boltz
Contact for lead author	Stan Boltz, <a href="mailto:stanley.boltz@sd.usda.gov">stanley.boltz@sd.usda.gov</a> , 605-352-1236
Date	01/05/2010
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** Slight to none, typically on steeper slopes and discontinuous.
- 

2. **Presence of water flow patterns:** None, or barely visible and discontinuous with numerous debris dams when present.
- 

3. **Number and height of erosional pedestals or terracettes:** Some pedestalled plants typically on steeper slopes.
- 

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0 to 15 percent is typical.
- 

5. **Number of gullies and erosion associated with gullies:** None should be present.

- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Small size litter classes will generally move short distances, some medium size class litter will move very short distances. Litter debris dams are occasionally present.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil aggregate stability ratings should typically be 5 to 6, normally 6. Surface organic matter adheres to the soil surface. Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon should be 2 to 6 inches thick with light to dark brownish gray colors. Structure should typically be fine granular at least in the upper A-horizon. Some soils have subangular blocky structure parting to weak fine granular.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (shrubs, mid & tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None – when dry, subsoil can be hard and appear to be compacted, but no platy structure will be present.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Tall warm-season grasses >
- Sub-dominant: Tall cool-season bunchgrasses = Mid warm-season grasses = Shrubs = Trees >
- Other: Forbs > Short warm- and cool-season grasses
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little evidence of decadence or mortality. Bunch grasses have strong, healthy centers and shrubs are vigorous.
- 
14. **Average percent litter cover (%) and depth ( in):** Litter cover is typically 50 to 80 percent, with the depth roughly 0.25 to 0.5 inches.

- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Total annual production ranges from 1,800 to 3,200 pounds/acre, with the reference value being 2,500 pounds/acre (air-dry basis).
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious weeds, Eastern redcedar.
- 
17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.
-