

Ecological site R066XY026NE

Loamy Overflow

Accessed: 05/18/2024

General information

**Approved.** An approved ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model, enough information to identify the ecological site, and full documentation for all ecosystem states contained in the state and transition model.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Classification relationships

Level IV Ecoregions of the Conterminous United States: 43i – Keya Paha Tablelands.

Associated sites

R066XY036NE	<b>Loamy 18-22 P.Z.</b> Loamy 18-22" P.Z.
R066XY058NE	<b>Loamy 22-25 P.Z.</b> Loamy 22-25" P.Z.
R066XY066NE	<b>Loamy Terrace</b> Loamy Terrace

Similar sites

R066XY066NE	<b>Loamy Terrace</b> Loamy Terrace [less big bluestem, lower production]
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

## Physiographic features

This site occurs on nearly level areas along drainageways that receive additional water from overflow of intermittent streams or runoff from adjacent slopes.

**Table 2. Representative physiographic features**

Landforms	(1) Flood plain (2) Stream terrace (3) Swale
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Rare to frequent
Ponding frequency	None
Elevation	579–914 m
Slope	0–2%
Water table depth	107–203 cm
Aspect	Aspect is not a significant factor

## Climatic features

MLRA 66 is considered to have a continental climate – cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the northern Great Plains and the winds move freely across the plains and account for rapid changes in temperature.

Annual precipitation ranges from 18 to 25 inches per year. The normal average annual temperature is about 48° F. January is the coldest month with average temperatures ranging from about 19° F (Bonesteel, SD) to about 23° F (Ainsworth, NE). July is the warmest month with temperatures averaging from about 73° F (Harrington, SD) to about 75° F (Gregory, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 54° F. This large annual range attests to the continental nature of this area's climate. Hourly winds average about 10 miles per hour annually, ranging from about 11 miles per hour during the spring to about 9 miles per hour during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour.

Growth of native cool season plants begins mid to late March and continues to late June. Native warm season plants begin growth in early May and continue to late August. Green up of cool season plants may occur in September and October when adequate soil moisture is present.

**Table 3. Representative climatic features**

Frost-free period (average)	140 days
Freeze-free period (average)	158 days
Precipitation total (average)	559 mm

## Influencing water features

Soil features

The soils of this site are very deep, well drained soils that formed in alluvium. These soils have moderate to rapid permeability. The surface layer will vary from 3 to 15 inches deep and have one of the following textures: silt loam, loamy fine sand, fine sandy loam, and silty clay loam. These areas receive additional water from overflow of intermittent streams or runoff from adjacent slopes. Available water capacity is typically high. The general fertility level and organic content of these soils is medium to high. This site should show slight to no evidence of rills, wind scoured areas or pedestalled plants. Water flow paths are broken, irregular in appearance or discontinuous with numerous debris dams or vegetative barriers. The soil surface is stable and intact. Sub-surface soil layers are not restrictive to water movement and root penetration.

These soils are mainly susceptible to water erosion. Headcuts may develop if adequate vegetative cover is not maintained. A drastic loss of the soil surface layer on this site can result in a shift in species composition and/or production.

More information can be found in the various soil survey reports. Contact the local USDA Service Center for soil survey reports that include more detail specific to your location.

Table 4. Representative soil features

Surface texture	(1) Silty clay loam (2) Fine sandy loam (3) Loamy fine sand
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderate to rapid
Soil depth	203 cm
Available water capacity (0-101.6cm)	15.24–20.32 cm
Calcium carbonate equivalent (0-101.6cm)	0–15%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	5.6–8.4

Ecological dynamics


This site developed under Northern Great Plains climatic conditions, light to severe grazing by bison and other large herbivores, sporadic natural or man-caused wildfire (often of light intensities), and other biotic and abiotic factors that typically influence soil/site development. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plant and animal species, and management actions. While the following descriptions describe more typical transitions between communities that will occur, severe disturbances, such as periods of well-below average precipitation, can cause significant shifts in plant communities and/or species composition.

Continuous season-long grazing (during the typical growing season of May-October) and/or repeated seasonal grazing (e.g., every spring, every summer) without adequate recovery periods following each grazing occurrence causes this site to depart from the Big Bluestem/Western Wheatgrass Plant Community. Western wheatgrass increases initially and will eventually decrease with continuous grazing. Grasses such as big bluestem, prairie cordgrass and switchgrass will decrease in frequency and production. Introduced species such as Kentucky bluegrass, cheatgrass and smooth brome grass invade the site as a result of inadequate recovery periods between grazing events and overstocking. Where trees dominate the site, woody regeneration will decline and grasses and

forbs will become dominant in the understory. It is thought that the climax is an herbaceous dominated site on higher landscape positions where trees encroach from the adjacent landscapes, and flooding events are infrequent. Loamy Overflow sites occupying lower landscape positions or plant communities adjacent to riparian areas will typically be dominated by a mixed hardwood overstory.

Interpretations are primarily based on the Big Bluestem/Needlegrass/Western Wheatgrass Plant Community. It has been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been used. Plant communities, states, transitional pathways, and thresholds have been determined through similar studies and experience.

### State and transition model



**BM** – Brush management (fire, chemical, mechanical); **CSLG** – Continuous season-long grazing (grazing a unit for an entire growing season); **F** – Fire; **LTPG** – Long-term prescribed grazing; **NF** – No fire; **NU** – Non use; **PG** – Prescribed grazing.

Figure 6. Plant Communities and Transitional Pathways

## Inventory data references

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Wayne Bachman, Soil Scientist, NRCS; Stan Boltz, Range Management Specialist, NRCS; Anna Ferguson, Soil Conservationist, NRCS; Roger Hammer, Soil Scientist, NRCS; Dana Larsen, Range Management Specialist, NRCS; Dave Schmidt, Rangeland Management Specialist, NRCS; Kim Stine, Rangeland Management Specialist, NRCS.

## Other references

High Plains Regional Climate Center,  
University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://www.hprcc.unl.edu/>)

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224.  
(<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>).  
National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS, Various Published Soil Surveys.

## Contributors

Stan Boltz

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Boltz
Contact for lead author	Stan Boltz, <a href="mailto:stanley.boltz@sd.usda.gov">stanley.boltz@sd.usda.gov</a> , 605-352-1236
Date	08/01/2006
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** None.

- 
2. **Presence of water flow patterns:** Typically none or barely visible. Evidence of water flow may be present after high overland flow events or flooding from adjacent streams, but vegetation normally remains intact.

- 
3. **Number and height of erosional pedestals or terracettes:** None.
- 
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0 to 5 percent is typical.
- 
5. **Number of gullies and erosion associated with gullies:** None typical, however limited headcutting may form after high runoff or flooding events. Existing gullies should be stabilized with good vegetative cover.
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None typical, but limited deposition may occur after major runoff or flooding events.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Litter of small and medium size classes will move after average to high rainfall events. Litter does not travel far, typically being trapped in small bunches by the extensive vegetative cover. Litter movement may be fairly extensive after major runoff or flooding events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil aggregate stability ratings should typically be 5 to 6, normally 6. Surface organic matter adheres to the soil surface. Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon should be 8 to 15 inches thick or greater with mollic (dark) colors when moist. Structure typically is medium to fine granular in the upper A-horizon.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None – when dry, B horizons can be hard and appear to be compacted, but no platy structure will be present, or depth to these layers would not correspond to compaction by grazing animals.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Tall, warm-season rhizomatous grasses >>

Sub-dominant: Mid, cool-season rhizomatous grasses > mid, warm-season grasses = mid and tall, cool-season bunchgrasses >

Other: Forbs > shrubs > grass-like species > trees > short, cool-season bunchgrasses

Additional: In the woody draw areas, deciduous trees may dominate the site. In these areas, the tall, warm-season rhizomatous functional group may be the third most dominant in the herbaceous understory.

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little evidence of decadence or mortality. Bunch grasses have strong, healthy centers and shrubs are vigorous.
  14. **Average percent litter cover (%) and depth ( in):** Litter cover is typically 70 to 90 percent, and depth of litter ranges from 0.25 to 0.5 inches.
  15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Total annual production ranges from 2,500 to 4,300 pounds/acre, with the reference values being 3,400 pounds/acre (air-dry basis).
  16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** State and local noxious weeds, Kentucky bluegrass, smooth brome grass, snowberry.
  17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.
-