Ecological site R066XY066NE
Loamy Terrace

Accessed: 10/12/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Boltz
Contact for lead author	Stan Boltz, stanley.boltz@sd.usda.gov, 605-352-1236
Date	08/01/2006
Approved by	Stan Boltz
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

2. **Presence of water flow patterns:** None, or barely visible and discontinuous.

3. **Number and height of erosional pedestals or terracettes:** None.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0 to 5 percent is typical.

5. **Number of gullies and erosion associated with gullies:** None should be present.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None.

7. **Amount of litter movement (describe size and distance expected to travel):** Litter should fall in place. Slight amount of movement of smallest size class litter is possible, but not normal.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil aggregate stability ratings should typically be 5 to 6, normally 6. Surface organic matter adheres to the soil surface. Soil surface fragments will typically retain structure indefinitely when dipped in distilled water.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon should be 8 to 15 inches thick with mollic (dark) colors when moist. Structure typically is medium to fine granular in the upper A-horizon.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool- and warm-season grasses) with fine and coarse roots positively influences infiltration.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None – when dry, B horizons can be hard and appear to be compacted, but no platy structure will be present.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

    Dominant: Mid, cool-season rhizomatous grasses > mid and tall, cool-season bunchgrasses >

    Sub-dominant: Tall, warm-season rhizomatous grasses > shrubs >

    Other: Forbs > trees > grass-like species > short, warm-season grasses

    Additional: Other grasses in other functional groups occur in minor amounts.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very little evidence of decadence or mortality. Bunch grasses have strong, healthy centers and shrubs are vigorous.

14. **Average percent litter cover (%) and depth (in):** Litter cover is roughly 60 to 80 percent, and the depth is 0.25 to 0.5 inches.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Total annual production ranges from 2,000 to 3,600 pounds/acre, with the reference value being 2,800 pounds/acre (air-dry basis).

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that**

become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: State and local noxious weeds; also Kentucky bluegrass and smooth bromegrass.

---

17. **Perennial plant reproductive capability:** All species exhibit high vigor relative to climatic conditions. Do not rate based solely on seed production. Perennial grasses should have vigorous rhizomes or tillers.