

Ecological site R067AY106WY Closed Depression (Cd)

Last updated: 9/07/2023 Accessed: 05/07/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| Author(s)/participant(s) | Dave Cook, Kristin Dickinson, George Gamblin, John Hartung, Andy Steinert, Nadine Bishop |
|---|--|
| Contact for lead author | |
| Date | 11/19/2020 |
| Approved by | Kirt Walstad |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

present on the site.

| | indicators | | |
|----|---|--|--|
| 1. | Number and extent of rills: None. Rills are not expected on the site. | | |
| 2. | Presence of water flow patterns: None. Water flow patterns are not expected on this site. | | |
| 3. | Number and height of erosional pedestals or terracettes: None. Erosional pedestals or terracettes are not expected on this site. | | |
| 4. | Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Bare ground is typically less than 10 percent. After prolonged ponding, bare ground will increase for relatively short periods of time. | | |
| 5. | Number of gullies and erosion associated with gullies: None. Gullies should not be present on this site. | | |

6. Extent of wind scoured, blowouts and/or depositional areas: None. Wind-scoured and/or depositional areas are not

- 7. Amount of litter movement (describe size and distance expected to travel): Litter should fall in place. Slight amount of movement of fine litter from water is possible, but not normal. Litter movement from wind is not expected.
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Soil aggregate stability ratings typically 5 to 6, normally 6. Surface organic matter adheres to the soil surface. Soil surface peds will typically retain structure indefinitely when dipped in distilled water.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): The surface layer ranges from a depth of 2 to 8 inches (5-20.3 cm). Soil colors are gray (5/1) dry and very dark gray (3/1) moist. Soil surface structure is typically granular. Soils tend to crack when dry.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: The functional/structural groups provide a combination of rooting depths and structure which positively influences infiltration. Combination of shallow and deep rooted species (mid & tall rhizomatous and tufted perennial cool season grasses) with fine and coarse roots positively influences infiltration. The expected composition of the plant community is about 85 to 95 percent perennial grasses and grass-likes, 5 to 10 percent forbs, and 0 to 5 percent shrubs.

The grass and grass-like component is made up cool-season, rhizomatous grasses (45-50%); cool-season bunch grasses (20-30%); warm-season, tall and mid-grasses (0-15%), warm-season short grasses (0-5%), and grass-likes (0-5%).

- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): A compaction layer is not expected on this site.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Community 1.1 – 12-14" PZ:

1. Native, C3, rhizomatous grasses - 810-900 #/ac (45-50%), 1 species minimum

Community 1.1 – 15-17" PZ:

1. Native, C3, rhizomatous grasses – 900-1000 #/ac (45-50%), 1 species minimum

Sub-dominant: Community 1.1 – 12-14" PZ:

- 2. Native, C3, bunch grasses 360-540 #/ac (20-30%), 1 species minimum
- 3. Native, Perennial and Annual Forbs 100-270 #/ac (5-15%), 5 species minimum

Community 1.1 – 15-17" PZ:

- 2. Native, C3, bunch grasses 400-600 #/ac (20-30%), 1 species minimum
- 3. Native, Perennial and Annual Forbs 100-300 #/ac (5-15%), 5 species minimum

Other: Community 1.1 – 12-14" PZ:

4. Minor: Native, C4, tall and mid-grasses – 0-270 #/ac (0-15%)

- 5. Minor: Grass-likes 0-90 #/ac (0-5%)
- 6. Minor: Native, C4, short grasses 0-90#/ac (0-5%)
- 7. Minor: Shrubs, vines, cacti 0-90 #/ac (0-5%)

Community 1.1 – 15-17" PZ:

- 4. Minor: Native, C4, tall and mid-grasses 0-300 #/ac (0-15%)
- 5. Minor: Shrubs, vines, cacti 0-200 #/ac (0-10%)
- 6. Minor: Grass-likes 0-100 #/ac (0-5%)
- 7. Minor: Native, C4, short grasses 0-100#/ac (0-5%)

Additional: 12a. Relative Dominance:

12-14" PZ: Community 1.1: Native, C3, rhizomatous grasses > Native, C3, bunch grasses > Native, Perennial and Annual Forbs > Native, C4, tall and mid-grasses > Grass-likes = Native, C4, short grasses = Shrubs, vines, cacti

15-17" PZ: Community 1.1:

Community 1.1: Native, C3, rhizomatous grasses > Native, C3, bunch grasses > Native, Perennial and Annual Forbs > Native, C4, tall and mid-grasses > Shrubs, vines, cacti > Grass-likes = Native, C4, short grasses

12b. F/S Groups not expected for the site: Introduced annual grasses, perennial introduced and naturalized grasses, trees.

12c. Number of F/S Groups: 7

12d. Species number in Dominant and Sub-dominant F/S Groups: 7

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very little evidence of decadence or mortality. Bunch grasses have strong, healthy centers with less than 3 percent mortality and shrubs have few dead stems.
- 14. Average percent litter cover (%) and depth (in): Plant litter cover is evenly distributed throughout the site and is expected to be 50 to 70 percent. Litter depth is expected to be 0.50 to 1.50 inch (1.3 to 3.9 cm). Plant litter cover will vary with the length and duration of ponding.
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): In the 12-14 inch Precipitation Zone, annual production ranges from 900 to 2,700 pounds per acre on an air dry basis. Average annual production is 1,800 pounds per acre under normal precipitation and weather conditions.

In the 15-17 inch precipitation zone, annual production ranges from 1,000 to 3,000 pounds per acre on an air dry basis. Average annual production is 2000 pounds per acre under normal precipitation and weather conditions.

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Annual bromes, Canada thistle, Russian thistle, kochia, cottonwood trees (primarily in the eastern portion of the MLRA) and others as they become known.

See:

Colorado Department of Agriculture Invasive Species Website:

https://www.colorado.gov/pacific/agconservation/noxious-weed-species

Wyoming Weed and Pest Council Website: https://wyoweed.org/ Nebraska Invasive Species website: https://neinvasives.com/plants

17. **Perennial plant reproductive capability:** All perennial species exhibit high vigor relative to recent weather conditions. Perennial grasses should have vigorous rhizomes or tillers; vegetative and reproductive structures are not stunted. All perennial species should be capable of reproducing annually.