

Ecological site R072XY102KS Saline Subirrigated

Accessed: 04/18/2024

General information

Approved. An approved ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model, enough information to identify the ecological site, and full documentation for all ecosystem states contained in the state and transition model.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 072X--Central High Tableland

Major Land Resource Area (MLRA) 72--Central High Tableland. This area is in Kansas (54 percent), Nebraska (25 percent), and Colorado (21 percent). A very small part of the area is in Wyoming. The area makes up about 34,550 square miles (89,535 square kilometers). It includes the towns of Garden City, Goodland, and Colby, Kansas; Imperial, North Platte, Ogallala, and Sidney, Nebraska; and Holyoke and Wray, Colorado. Interstate 70 bisects the area, and Interstates 76 and 80 follow the south side of the South and North Platte Rivers, respectively. The Cimarron National Grasslands occur in the southwest corner of the MLRA.

Classification relationships

Major Land Resource Area (MLRA) 72--Central High Tableland.

Ecological site concept

The Saline Subirrigated ecological site occurs on nearly level areas, adjacent to streams having underground water movement. This site consists of alluvial soils that usually contain a high concentration of salts. A seasonal or perennial water table less than 6 feet from the surface characterizes this site. Due to the landform position this site is subject to flooding.

This ESD was developed using the existing ESD's developed in 2001: Saline Subirrigated North R072XA020KS, Saline Subirrigated South R072XB020KS and Salt Meadow R072XY035CO. Changes were made in accordance with policy using the National Ecological Site Handbook. Text changes were made in regards to clarity and style. Content was changed reflecting the range of variability of the site.

Associated sites

R072XY103KS	Subirrigated Subirrigated site is located adjacent to Saline Subirrigated and has a seasonal or perennial high water table (<6 feet from surface).
R072XY104KS	Saline Lowland Saline lowland is located adjacent to Saline Subirrigated and does not have a seasonal or perennial high water table (>6 feet from surface).

Similar sites

R072XY104KS	Saline Lowland Saline Lowland and Saline Subirrigated will have similar vegetation and have soils with high concentration of salts. The significant differences between the sites is the depth to the seasonal or perennial high water table. The water table is greater than 6 feet from the surface on a Saline Lowland site and less than 6 feet from the surface on a Saline Subirrigated site.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Sporobolus airoides</i> (2) <i>Distichlis spicata</i>

Physiographic features

The Saline Subirrigated site occurs on nearly level floodplains adjacent to streams and rivers. These soils are seasonally wet, have an accumulation of salts within the root zone, and are subject to flooding. These soils receive runoff from areas higher on the landscape.

The extent of the major Hydrologic Unit Areas (identified by four-digit numbers) that make up MLRA 72 is as follows: Republican (1025), 38 percent; Middle Arkansas (1103), 20 percent; Smoky Hill (1026), 15 percent; South Platte (1019), 13 percent; Upper Cimarron (1104), 11 percent; North Platte (1018), 2 percent; and Upper Arkansas (1102), 1 percent. The North Platte River forms the northern boundary of this MLRA. The South Platte River joins the North Platte River at the town of North Platte, Nebraska. The Arkansas River bisects the southern part of the MLRA. Other large rivers between the North Platte and Arkansas Rivers in the area include the Republican, Sappa, Prairie Dog, Solomon, Saline, and Smoky Hill Rivers. The Cimarron River is the southern boundary of MLRA 72 in the western part of this area.




Figure 2. MLRA 72 block diagram

Table 2. Representative physiographic features

Landforms	(1) Flood plain
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Rare to frequent
Ponding frequency	None
Elevation	2,500–5,000 ft
Slope	0–3%
Water table depth	12–48 in

Climatic features

The average annual precipitation in this area ranges from 12 to 23 inches (305 to 584 millimeters). It fluctuates widely from year to year. Most of the rainfall occurs as high-intensity, convective thunderstorms during the growing season. The maximum precipitation occurs from late spring through early autumn. Precipitation in winter occurs as snow. The annual snowfall ranges from about 16 inches (40 centimeters) in the southern part of the area to 35 inches (90 centimeters) in the northern part. The average annual temperature is 46 to 57 degrees F (8 to 14 degrees C). The freeze-free period averages 162 days and ranges from 135 to 210 days, increasing in length from northwest to southeast. Climate data comes from the Natural Resources Conservation Service (NRCS) National Water and Climate Center. The data set is from 1981-2010.

Table 3. Representative climatic features

Frost-free period (average)	147 days
Freeze-free period (average)	162 days
Precipitation total (average)	19 in


Climate stations used

- (1) LAKIN [USC00144464], Lakin, KS
- (2) GARDEN CITY RGNL AP [USW00023064], Garden City, KS
- (3) SHARON SPRINGS [USC00147397], Sharon Springs, KS
- (4) IDALIA [USC00054242], Idalia, CO
- (5) SYRACUSE 1NE [USC00148038], Syracuse, KS
- (6) CROOK [USC00051996], Crook, CO
- (7) JULESBURG [USC00054413], Julesburg, CO
- (8) BENKELMAN [USC00250760], Benkelman, NE

Influencing water features

A seasonal water table and adjacent perennial water table have influence on the kinds and amounts of vegetation on this site.

Figure 7-1 The hydrologic cycle with factors that affect hydrologic processes



7.1-4

(190-VI-NRPH, December 2003)

Figure 7. Fig. 7-1 from National Range and Pasture Handbook

Soil features

This site consists of deep to very deep saline and saline-alkali soils. The high concentration of salts affects both the kind and amount of vegetation present. Surface and subsoil layers have a seasonal high water table between 12 and 48 inches below the soil surface in normal to wet years. These soils are generally very deep, but some are shallow over gravelly coarse sand. The surface soil is generally darker colored and ranges from 7 to 24 inches thick. Less common are light colored soils with a surface soil less than 7 inches. Texture of the soil surface ranges widely from silt loam to fine sand. The underlying material is lighter colored than the surface soil, and commonly has redoximorphic concentrations in the upper part. It ranges widely in texture from loam to gravelly coarse sand. Some soils in this site are calcareous to the surface. Saline and alkali salts affect other soil properties such as available water capacity, soil structure, infiltration, and permeability. Soils in this site generally have a low to moderate content of organic matter.

Major soil series correlated to this ecological site include Las Animas (20%), Las (19%), Lawet (8%), Lebsack (6.5%), Caruso (6.2%), Marksbutte (5.7%), and Lesho (5.5%).




Figure 8. Las Animas OSD profile

Table 4. Representative soil features

Surface texture	(1) Loam (2) Sandy loam
-----------------	----------------------------

Family particle size	(1) Loamy
Drainage class	Poorly drained to moderately well drained
Permeability class	Moderately slow to rapid
Soil depth	60–80 in
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0–3%
Available water capacity (0-40in)	2.8–10 in
Calcium carbonate equivalent (0-40in)	0–40%
Electrical conductivity (0-40in)	4–20 mmhos/cm
Sodium adsorption ratio (0-40in)	2–30
Soil reaction (1:1 water) (0-40in)	7.4–9
Subsurface fragment volume <=3" (Depth not specified)	0–50%
Subsurface fragment volume >3" (Depth not specified)	0–10%

Ecological dynamics

The plant community for this site is dynamic due to the complex interaction of many ecological processes. The interpretive plant community for this site is the Reference Plant Community. This community has been determined by the study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing strategies. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

This site developed with occasional fires as part of the ecological processes. Historically, it is believed that the fires were infrequent, randomly distributed, and started by lightning at various times throughout the season when thunderstorms were likely to occur. It is also believed that pre-European inhabitants may have used fire as a management tool for attracting herds of large migratory herbivores (bison, elk, deer, and pronghorn). The impact of fire over the past 100 years has been relatively insignificant due to the human control of wildfires and the lack of acceptance of prescribed fire as a management tool in the semi-arid, High Plains area.

The Saline Subirrigated site occurs on nearly level areas adjacent to streams or rivers. The site is characterized as subirrigated as a result of groundwater ranging from 12 to 48 inches from the surface throughout the growing season. The availability of water has a major influence on the vegetation that will persist on this site. This site is subject to occasional flooding. Historically, the flooding potential on these sites was much greater. The control of river water over the past 80 years through the use of structures has greatly reduced the frequency of flooding.

Soils of this site are characterized by moderate to high salinity or alkaline conditions. There will often be white or gray deposits on the soil surface early in the spring or following wet periods. The degree of salinity or alkali condition has a major influence on the plant community that will persist on this site. Plants that can tolerate these conditions such as alkali sacaton, western wheatgrass, inland saltgrass, alkali cordgrass, slender wheatgrass, and salt tolerant rushes, and sedges will dominate these high pH sites. There may be areas with a moderate or lower pH interspersed that have plant communities similar to that of the Subirrigated ecological site or a combination of the two. Other soil inclusions may be present.

Continuous grazing without adequate recovery periods on this site will result in the loss of vigor and reproductive potential for such species as alkali sacaton, western wheatgrass, slender wheatgrass, and switchgrass on this site. The grazing management coupled with the high pH conditions has shifted many of these sites into a primarily inland saltgrass, with some western wheatgrass, plant community. Due to the lower stature and yield potential of these

grasses, most of the sites are used for grazing rather than hay production. Once the plant community has moved to inland saltgrass, it is very difficult to reverse the shift.

The degree of herbivory (feeding on herbaceous plants) has a significant impact on the dynamics of the site. Historically, periodic grazing by herds of large migratory herbivores was a primary influence. Secondary influences of herbivory by species such as prairie dogs, grasshoppers, gophers, and root feeding organisms impacted the vegetation historically, and continue to this day.

The management of herbivory by humans through grazing of domestic livestock and/or manipulation of wildlife populations has been a major influence on the ecological dynamics of the site. This management coupled with the High Plains climate largely dictates the plant communities for the site.

Typically, drought cycles have not had a major impact upon the vegetation due to the presence of a water table. Depletion of ground water reserves may change species composition and possibly cause a shift in ecological sites.


Growth of native cool season plants begins about April 15, and continues to about June 15. Native warm season plants begin growth about May 15, and continue to about August 15. Green up of cool season plants may occur in September and October if adequate moisture is available.

The following state and transition model (STM) illustrates the common plant communities that can occur on the site and the pathways among communities.


Following the state and transition model are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition table shown below has been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed and new ones may be added. None of these plant communities should necessarily be thought of as "Desired Plant Communities". According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities will be determined by the decision-makers and will meet minimum quality criteria established by NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

State and transition model

Ecosystem states



State 1 submodel, plant communities



State 2 submodel, plant communities

2.1. Warm season sod forming

State 3 submodel, plant communities

3.1. Reseed Community

3.2. Go-back Community

**State 1
Grassland state**

The grassland state is supported by empirical data, historical data and local expertise. This state is defined by two native plant communities that are a result of periodic fire, drought and grazing. These events are part of the natural disturbance regime and climatic process. The reference plant community consist of warm season dominant and cool season sub-dominant grasses. The at-risk plant community is the other plant community within the grassland state. This plant community is made up of warm and cool season grasses with decreasing amounts of alkali sacaton.

**Community 1.1
Reference Plant Community**

The interpretive plant community for this site is the Reference Plant Community. The natural potential vegetation of this community is a mixed grass prairie. Grass and grass-like plants make up approximately 90-98% of the potential production and forbs make up 2-5%. Alkali sacaton, inland saltgrass, western wheatgrass and switchgrass are the dominant species in this community. Secondary species include alkali cordgrass, slender wheatgrass, and little bluestem. The plant community has a limited forb and shrub diversity due to the high pH of the soil and the adaptability of the plants to these conditions. The stability and diversity of this plant community are influenced by the frequency of flooding, fluctuation of groundwater, and salinity of the soil. Although this plant community will fluctuate due to natural conditions it is not resistant to heavy, continuous grazing without periodic rest. With proper management and adequate recovery periods this plant community will remain diverse, productive, and functioning. Total annual production ranges from 1,500 to 4,200 pounds of air-dried vegetation per acre per year and will average 3,000 pounds.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1365	2820	3850
Forb	125	150	300
Shrub/Vine	10	30	50
Total	1500	3000	4200

Figure 10. Plant community growth curve (percent production by month). KS7215, Warm season dominant, cool season sub-dominant. The growth curve represents a dominance (40-100% composition) of warm season grasses followed by a sub-dominance (10-40% composition) of cool season grasses..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	20	35	25	10	5	0	0	0

Community 1.2

At-risk community

The At-risk plant community is most vulnerable to exceeding the resilience limits of the Grassland State and transitioning to an alternative state. This plant community is a result of repetitive heavy use, not providing adequate rest, and no forage and animal balance. As a result, alkali sacaton, switchgrass, and other preferred grasses rapidly lose productive capacity through loss of vigor and reproductive potential. Western wheatgrass and inland saltgrass increase to fill the voids left as the preferred species decrease. This plant community is at risk of losing the tall warm season grass species and developing into an inland saltgrass sod community. When this occurs, it will require considerable time and expense to return this community to the Reference State. Total annual production ranges from 1,400 to 3,000 pounds of air-dried vegetation per acre per year and will average 2,200 pounds.

Pathway 1.1 to 1.2

Community 1.1 to 1.2

No forage and animal balance and repetitive heavy use without adequate recovery periods between grazing events will convert the Reference Plant Community to western wheatgrass, inland saltgrass, and alkali sacaton remnants. Annual haying during mid-summer or multiple hay harvests during a single growing season will accelerate the movement to western wheatgrass, inland saltgrass and alkali sacaton remnants.

Pathway 1.2 to 1.1

Community 1.2 to 1.1

This community pathway will require long term management (>20 years). Management will need to provide adequate rest and recovery to existing vegetation combined with a forage and animal balance. It will be necessary to have remnant Reference Plant Community species present to shift the At-risk Community to the Reference Plant Community.

State 2

Sod bound state

The Sod-bound state is supported by empirical data, historical data, and local expertise. This state is defined by one plant community that is a result of repetitive heavy use, no rest and recovery of the grasses, and no forage and animal balance. The Grassland State ecosystem has been driven beyond the limits of ecological resilience and has crossed a threshold into the Sod-bound State. The designation of the Sod-bound State denotes changes in plant community composition. This change in species composition affects hydrologic function and biotic integrity of the ecosystem. This alternative state should be treated as a hypothesis that will be tested through long-term observation of ecosystem behavior and repeated application of conservation and restoration practices. This state should be re-evaluated and refined continually.

Community 2.1

Warm season sod forming

Repetitive heavy use, no rest and recovery, and no forage and animal balance will favor inland saltgrass to dominate and become sod bound. Foxtail barley can increase in heavily used areas. Western wheatgrass, alkali sacaton, sedges, rushes, and other grasses or grass-likes may exist only in remnant populations scattered throughout the community. This plant community is resistant to change due to the grazing tolerance of inland saltgrass. A significant amount of production and diversity has been lost when compared to the Reference Plant Community. Loss of key warm and cool season grasses have negatively impacted energy flow and nutrient cycling. Water infiltration is reduced significantly due to the massive shallow root system, characteristic of inland saltgrass. The restoration pathway has not been documented and needs further study before it is reflected in the state and transition model. It could take generations to bring this plant community back to the Reference State with management alone. Renovation (mechanical and/or chemical inputs) is not recommended due to high salt content of the soil and saltgrass persistence. Populations of woody species can occur on this site. Russian olive and/or eastern red cedar will most likely be the invading species. Without management, an increase of these species can reduce the areas available for forage production. The open areas between the trees will be dominated by inland saltgrass. Remnant populations of other grasses may exist among the trees where access by livestock is limited.

Initial control of the woody invasion where the trees have reached trunk calipers of greater than one inch will require mechanical removal. The removal of Russian olive trees often results in a flush of eastern red cedar seedlings. When this occurs, additional brush control measures will be needed. More information will be necessary to determine if this is an alternative state in the state and transition model. Other species to monitor encroachment potential is salt cedar. Total annual production ranges from 900 to 2,100 pounds of air-dried vegetation per acre per year and will average 1,500 pounds.

State 3

Tillage state

The Grassland State ecosystem has been driven beyond the limits of ecological resilience and has crossed a threshold into the Tillage State. The designation of the Tillage State denotes changes in land use which has affected the plant community, soil inherent, and dynamic properties. Changes in soil structure, aggregate stability, content of organic matter, and species composition has negatively affected the hydrologic function, biotic integrity, and soil site stability of the site. This alternative state should be tested through long-term observation of ecosystem behavior and repeated application of conservation and restoration practices. This state should be re-evaluated and refined continually. The Tillage State includes two community phases that are highly variable. They are derived through two distinct management scenarios, and are not related successional. Infiltration, runoff, and soil erosion varies depending on the vegetation present.

Community 3.1

Reseed Community

This plant community can vary considerably depending on how eroded the soil was, the species seeded, the stand that was established, how long ago the stand was established, and the management of the stand since establishment. A forage and animal balance with adequate recovery periods will be necessary to maintain productivity and desirable species. There are several factors that make seeded rangeland a different grazing resource than native rangeland. Factors such as species selected, stand density, varieties, and harvest efficiency all impact the production level and palatability. This results in uneven grazing distribution when both seeded and native rangelands are in the same grazing unit. Therefore, the seeded rangeland should be managed as a separate grazing unit if possible. Seeded rangeland back to native adapted grass species can provide a productive haying resource. Species diversity on seeded rangeland is often lower and native forb species generally take longer to re-establish.

Community 3.2

Go-back Community

The Go-back Plant Community is created when the soil is tilled or farmed (sodbusted), and abandoned. All of the native plants are killed, soil organic matter/carbon reserves are reduced, soil structure is changed, and a plowpan or compacted layer can be formed. This compaction layer will decrease water infiltration. Synthetic chemicals may remain as a residual from farming operations. In early successional stages, this community is not stable. Erosion is a concern in this plant community. An annual plant community such as Russian thistle, kochia, foxtail barley, and other introduced annuals occupy the community initially. These plants give some protection from erosion. This plant community is gradually replaced by early perennial species such as inland saltgrass and western wheatgrass. Alkali sacaton and switchgrass can become established depending upon whether a remnant seed source is available. Eventually other perennial warm and cool season species can establish. This successional process can take generations as the soil begins to repair its dynamic and inherent properties (aggregate stability, structure, and content of organic matter).

Transition 1 to 2

State 1 to 2

Long term heavy grazing without adequate recovery periods will cause this plant community to change to an inland saltgrass sod, foxtail barley plant community.

Transition 2 to 3

State 1 to 3

This transition is triggered by a management action as opposed to a natural event. Tillage or breaking the ground with machinery for crop production will move the grassland state to a tillage state.

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Warm season grass (salts)			800–1300	
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	450–750	–
	saltgrass	DISP	<i>Distichlis spicata</i>	450–700	–
	alkali cordgrass	SPGR	<i>Spartina gracilis</i>	0–150	–
2	Cool season grass			400–800	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	400–600	–
	slender wheatgrass	ELTRT	<i>Elymus trachycaulus ssp. trachycaulus</i>	0–150	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	0–40	–
3	Warm season tallgrass			100–450	
	switchgrass	PAVI2	<i>Panicum virgatum</i>	150–400	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	0–270	–
	prairie cordgrass	SPPE	<i>Spartina pectinata</i>	0–70	–
4	Grasslike			15–130	
	sedge	CAREX	<i>Carex</i>	15–70	–
	rush	JUNCU	<i>Juncus</i>	15–70	–
5	Warm season midgrass			0–125	
	vine mesquite	PAOB	<i>Panicum obtusum</i>	0–100	–
	scratchgrass	MUAS	<i>Muhlenbergia asperifolia</i>	0–70	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–40	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	0–40	–
6	Warm season shortgrass			0–70	
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	0–40	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0–40	–
Forb					
7	Forbs			60–300	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	5–30	–
	pussytoes	ANTEN	<i>Antennaria</i>	0–30	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0–30	–
	Illinois bundleflower	DEIL	<i>Desmanthus illinoensis</i>	5–30	–
	American licorice	GLLE3	<i>Glycyrrhiza lepidota</i>	5–30	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	0–30	–
	swamp smartweed	POHY2	<i>Polygonum hydropiperoides</i>	5–30	–
	Pursh seepweed	SUCA2	<i>Suaeda calceoliformis</i>	0–30	–
	white heath aster	SYER	<i>Symphotrichum ericoides</i>	5–30	–
Shrub/Vine					
8	Shrubs			0–30	
	willow baccharis	BASA	<i>Baccharis salicina</i>	0–30	–

Animal community

Grazing Interpretations

The following table lists relative production values based on plant communities. These are conservative estimates

that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production.

Plant Community

Alkali Sacaton, Western Wheatgrass (Reference plant community)
Production (lbs/acre) - 3,000

Western Wheatgrass, Inland Saltgrass w/Remnant A. Sacaton
Production (lbs/acre) - 2,200

Inland Saltgrass Sod, Foxtail Barley
Production (lbs/acre) - 1,500

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangelands in this area provide yearlong forage under prescribed grazing for cattle, sheep, horses, and other herbivores. During the dormant period, livestock may need supplementation based on reliable forage analysis.

Wildlife: Saltgrass is a larval foodplant for the Wandering Skipper (*Panoquina panoquinoides errans*) butterfly. It is also an important food in the diet of waterfowl and the Florida salt marsh vole (*Microtus pennsylvanicus dukecampbelli*). Ducks are reported to occasionally eat the dried seeds and controlled burning provides tender forages for wild geese. *Distichlis spicata* (inland saltgrass) is significant in the salt marshes, which provide nesting grounds for birds, fish, and larvae of many species of marine invertebrate animals. As salt marsh plants decompose, their stored nutrients provide a steady source of food for clams, crabs, and fish.

Hydrological functions

Water is the principal factor limiting forage production on this site. Infiltration is moderate. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff.

Recreational uses

The wide variety of plants which bloom from spring until fall have an aesthetic value that appeals to visitors.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Other information

Site Development and Testing Plan.

Future work (for approved ESD) includes field visits to verify ES site concepts with field staff. Field staff include but are not limited to project office leader, area soil scientist, state soil scientist, ecological site specialist, state rangeland conservationist, area rangeland management specialist, and local field personnel. This site includes collaboration between Kansas, Colorado and Nebraska. Field visits are to be determined by spatial extent of the site as well as personal knowledge of the site. Activity during field visits will include but are not limited to identifying the soil, landform, plant community, and verifying existing site concepts.

Inventory data references

Information presented for Saline Subirrigated North has been derived from NRCS clipping data, numerous ocular estimates, and other inventory data. Field observations from experienced range personnel was used extensively to develop this ecological site description.

Those involved in developing the ESD Saline Subirrigated North include, Harvey Sprock from Colorado. Carol Eakins, Chuck Markley, Jeff Nichols, and Mary Schrader from Nebraska. Joan Gienger and Ted Houser from Kansas.

Information presented for Saline Subirrigated South has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel was also used. Those involved in developing this site include: Tim Watson, Amanda Shaw, Susan Francis, Jon Deege, Harvey Sprock, Robert Schiffner, and Josh Saunders.

All of this ESD information was developed using the 2001 ESD's (Saline Subirrigated North and South and Salt Meadow site in Colorado). Updates and data entry were completed by Chris Tecklenburg in accordance with National Ecological Site Handbook.

Other references

David J. Kraft, State Rangeland Management Specialist, NRCS, Kansas. Site Author circ. early 2000s.

High Plains Regional Climate Center, University of Nebraska, Lincoln(<http://hpcc.unl.edu>)

N. C. Brady and R. R. Weill, The Nature and Properties of Soils, 14th Edition, 2008, pp. 504–517.

USDA, NRCS. National Water and Climate Center, Portland, OR. (<http://wcc.nrcs.usda.gov>)

USDA, NRCS. National Range and Pasture Handbook, September 1997

USDA, NRCS. National Soil Information System, Information Technology Center, Fort Collins, CO. (<http://nasis.nrcs.usda.gov>)

USDA, NRCS. 2002. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA USA.

Contributors

Chris Tecklenburg

Acknowledgments

The ecological site development process is a collaborative effort, conceptual in nature, dynamic, and is never considered complete. I thank all those who set the foundational work in the early 2000s in regards to this ESD. I thank all those who contributed to the development of this site. In advance, I thank those who would provide insight, comments, and questions about this ESD in the future.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Original reference sheet created by David Kraft, John Henry, Doug Spencer, and Dwayne Rice in February 2005. Entered information and made minor revisions on #5, #12, #15 by Chris Tecklenburg 1-8-2016.
Contact for lead author	Chris Tecklenburg chris.tecklenburg@ks.usda.gov David Kraft david.kraft@ks.usda.gov
Date	01/08/2016
Approved by	David Kraft
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

2. **Presence of water flow patterns:** There is little, if any, evidence of soil deposition or erosion. Water generally flows evenly over the entire landscape.

3. **Number and height of erosional pedestals or terracettes:** There is no evidence of pedestaled plants or terracettes on the site.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Less than 3% bare ground is found on this site. Cover can be defined as live plants, litter, rocks, moss, lichens, etc.

5. **Number of gullies and erosion associated with gullies:** None

6. **Extent of wind scoured, blowouts and/or depositional areas:** There is no evidence of wind erosion creating bare areas or denuding vegetation.

7. **Amount of litter movement (describe size and distance expected to travel):** Plant litter is distributed evenly throughout the site.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Plant canopy is large enough to intercept the majority of raindrops. A soil fragment will not "melt" or lose its structure when immersed in water for 30 seconds. There is no evidence of pedestaled plants or terracettes. Soil stability scores will range from 5-6.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** The

topsoil layer has not been disturbed or eroded. Site is consistent with published soil survey description. A horizon 0 to 6 inches; dark gray (N 4/0) moist, moderate fine granular structure; soft, very friable; strongly effervescent, moderately alkaline (pH 8.2).

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** There is no negative effect on water infiltration and/or runoff due to plant composition or distribution. Plant composition and distribution are adequate to prevent any rill formation and/or pedestalling. Inter-spatial distribution is consistent with expectation for the site.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** There is no evidence of compacted soil layers due to cultural practices. Soil structure is conducive to water movement and root penetration.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm season grass (salts): inland saltgrass = alkali sacaton >> alkali cordgrass
- Sub-dominant: Cool season grasses: western wheatgrass >> slender wheatgrass >> Canada wildrye
- Other: Warm season tallgrasses (switchgrass, indiangrass, prairie cordgrass) > Grasslike (sedge, rush) = warm season midgrass > warm season shortgrass
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** The majority of plants are alive and vigorous. Some mortality and decadence is expected for the site. This in part is due to drought, unexpected wildfire or a combination of the two events. This would be expected for both dominant and sub-dominant groups.
-
14. **Average percent litter cover (%) and depth (in):** Plant litter is distributed evenly throughout the site. There is no restriction to plant regeneration due to depth of litter. When prescribed burning is practiced there will be little litter the first half of the growing season. 50-70% litter cover at 1.0 - 1.5 inch depth.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 3000 lbs. Vegetative production is 95-100% of normal based upon the range site description and the weather the past year. (refer to ecological site description for favorable or unfavorable growing conditions)
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site: There are no noxious weeds present. Invasive plants make up a small percentage of plant community, and invasive brush species are < 5% canopy.

17. **Perennial plant reproductive capability:** Plants on site exhibit the required vigor and growth to be able to reproduce vegetatively or by seed.
-