

Ecological site R077BY725TX Draw 12-17" PZ

Last updated: 9/11/2023 Accessed: 04/28/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Bradbury, Zone RMS, NRCS, Lubbock, Texas
Contact for lead author	806-791-0581
Date	09/04/2007
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

IIC	idicators							
1.	Number and extent of rills: None.							
2.	Presence of water flow patterns: None.							
3.	Number and height of erosional pedestals or terracettes: None.							
4.	Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 20-25%							
5.	Number of gullies and erosion associated with gullies: None.							
6.	Extent of wind scoured, blowouts and/or depositional areas: None.							
7.	Amount of litter movement (describe size and distance expected to travel): None to slight.							

8.	Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Very resistant to erosion.				
9.	Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Loam to clay loam; friable surface; high SOM.				
10.	Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Basal cover and density with small interspaces should make rainfall impact minimal. This site has moderately permeable soil, runoff is slow to medium and available water holding capacity is high.				
11.	Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None				
12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):				
	Dominant: Cool-season grasses >				
	Sub-dominant: Warm-season midgrasses > Warm-season tallgrasses > Warm-season shortgrasses >				
	Other: Forbs > Shrubs/Vines > Trees				
	Additional:				
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Grasses due to their growth habit will exhibit some mortality and decadence, though minimal.				
14.	Average percent litter cover (%) and depth (in): Dominant litter is herbaceous.				
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): 2,200 - 2,800 lbs/ac				
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Yucca and baccharis. Broom snakeweed can become invasive.				
17.	Perennial plant reproductive capability: All plant species should be capable of reproduction except during periods of				

prolonged drought conditions, heavy natural herbivory or intense wildfires.