

Ecological site R077CY036TX Sandy Loam 16-21" PZ

Last updated: 9/11/2023
Accessed: 05/06/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 077C–Southern High Plains, Southern Part

MLRA 77C is characterized by nearly level plains with numerous playa depressions, moderately sloping breaks along drainageways, and a steep escarpment along the eastern margin. From southwest to northeast, soils grade from coarse-textured to fine-textured. Soils are generally deep and occur in a thermic soil temperature regime and ustic soil moisture regime bordering on aridic. Current land use is dominantly cropland.

Classification relationships

This ecological site is correlated to soil components at the Major Land Resource Area (MLRA) level which is further described in USDA Ag Handbook 296.

Ecological site concept

This site occurs on gently sloping sandy loam soils on uplands. The reference vegetation consists of midgrasses with some shortgrasses and forbs and few woody plants. Abusive grazing practices can lead to a decrease in palatable species and a shift in the plant community. Without fire or other brush management, woody species may increase across the site.

Associated sites

R077CY027TX	<p>Playa 16-21" PZ</p> <p>The Playa site occurs on fine textured soils of closed depressions or playa basins. Water is ponded for various lengths of time. These sites are on lower landscape positions and water is ponded for various lengths of time. Vegetation is variable and may fluctuate because of the duration and depth of ponding.</p>
R077CY028TX	<p>Limy Upland 16-21" PZ</p> <p>The Limy Upland is on plains, interdunes, playa steps, and playa slopes. This site is associated with Sandy Loam sites that are on similar or slightly lower landscape positions. Midgrasses and shortgrasses dominate these sites.</p>
R077CY037TX	<p>Very Shallow 16-21" PZ</p> <p>The shallow site is on uplands and has calcareous, very gravelly soils or soils that are shallow to an indurated petrocalcic horizon. These sites are on landscape positions similar to the Sandy Loam site. The reference plant community consists of midgrasses, shortgrasses and forbs. Overall production is limited due to the shallow depth of the soils.</p>

Similar sites

R077CY035TX	<p>Sandy 16-21" PZ</p> <p>This site occurs on very deep, loamy fine sand soils on gently rolling uplands. The reference vegetation includes tall and midgrasses with forbs and few shrubs. This site is less productive than the Sandy Loam site in MLRA 77C.</p>
R077DY047TX	<p>Sandy Loam 12-17" PZ</p> <p>This site occurs on very deep, loamy fine sand soils on gently rolling uplands. The reference vegetation includes tall and midgrasses with forbs and few shrubs. Mean annual precipitation is lower (15 to 17 inches). This site is less productive than the Sandy Loam site in MLRA 77C.</p>
R077BY021TX	<p>Sandy Loam 12-17" PZ</p> <p>This site occurs on very deep, loamy fine sand soils on gently rolling uplands. The reference vegetation includes tall and midgrasses with forbs and few shrubs.</p>

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Yucca</i> (2) <i>Prosopis glandulosa var. torreyana</i>
Herbaceous	(1) <i>Bouteloua curtipendula</i> (2) <i>Schizachyrium scoparium</i>

Physiographic features

This upland site is moderately deep to very deep, moderately well to well drained and composed of nearly level to gently sloping plains, concave plains in broad, shallow basins and swales, and adjacent to natural drains. Elevations generally range from 2,500 to 4,700 feet. Surface textures are fine sandy loams with slopes ranging from 0 to 5 percent and runoff is negligible to medium.

Table 2. Representative physiographic features

Landforms	(1) Plateau > Plain (2) Plateau > Playa slope (3) Plateau > Playa floor
Runoff class	Negligible to low
Flooding frequency	None
Ponding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Ponding frequency	Occasional to rare
Elevation	792–1,554 m

Slope	0–5%
Ponding depth	5–30 cm
Water table depth	13–61 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

Climate is semi-arid dry steppe. Summers are hot with winters being generally mild with numerous cold fronts that drop temperatures into the single digits for 24 to 48 hours. Temperature extremes are the rule rather than the exception. Humidity is generally low and evaporation high. Wind speeds are highest in the spring and are generally southwesterly. Canadian and Pacific cold fronts come through the region in fall, winter and spring with predictability and temperature changes can be rapid. Most of the precipitation comes in the form of rain and during the period from May through October. Snowfall averages around 15 inches but may be as little as 8 inches or as much as 36 inches. Rainfall in the growing season often comes as intense showers of relatively short duration. Long-term droughts occur on the average of once every 20 years and may last as long as five to six years (during these drought years, moisture during the growing season is from 50 to 60 percent of the mean). Based on long-term records, approximately 60 percent of the years are below the mean rainfall and approximately 40 percent are above the mean. May, June and July are the main growth months for perennial warm-season grasses. Forbs make their growth somewhat earlier.

Table 3. Representative climatic features

Frost-free period (characteristic range)	155-189 days
Freeze-free period (characteristic range)	187-205 days
Precipitation total (characteristic range)	457-508 mm
Frost-free period (actual range)	148-198 days
Freeze-free period (actual range)	184-210 days
Precipitation total (actual range)	457-559 mm
Frost-free period (average)	174 days
Freeze-free period (average)	198 days
Precipitation total (average)	483 mm

Climate stations used

- (1) BIG SPRING [USW00023041], Big Spring, TX
- (2) PORTALES [USC00297008], Portales, NM
- (3) DENVER CITY [USC00412408], Denver City, TX
- (4) CROSBYTON [USC00412121], Crosbyton, TX
- (5) CLOVIS 13 N [USC00291963], Clovis, NM
- (6) RAGLAND 3 SSW [USC00297226], McAlister, NM
- (7) MULESHOE #1 [USC00416135], Muleshoe, TX
- (8) LEVELLAND [USC00415183], Levelland, TX
- (9) PLAINS [USC00417074], Plains, TX
- (10) LAMESA 1 SSE [USC00415013], Lamesa, TX

Influencing water features

Water features are generally not an influencing factor on these sites. Some areas may receive runoff from heavy rainfall events and concentrate in low areas for a short period of time.

Wetland description

N/A

Soil features

The Sandy Loam site is made up of moderately deep to very deep, nearly level to gently sloping, noncalcareous fine sandy loams on uplands and in shallow playas or depressions and swales. This site was formed from loamy eolian sediments or wind-altered lacustrine sediments. Slopes range from 0 to 5 percent and elevation ranges from 2,600 to 5,100 feet. These soils are moderately well or well drained and permeability is moderate. Some sites are moderately deep to a restrictive petrocalcic horizon between 20 and 40 inches and have moderate permeability above a very slowly permeable layer. Due to the sandy loam soil texture, wind and water erosion is moderate without vegetative cover.

Major Soil Taxonomic Units correlated to this site include: Amarillo fine sandy loam, Arvana fine sandy loam, Seagraves fine sandy loam, and Tokio fine sandy loam, Zita fine sandy loam.

Table 4. Representative soil features

Parent material	(1) Eolian deposits—metamorphic and sedimentary rock
Surface texture	(1) Fine sandy loam
Family particle size	(1) Fine-loamy
Drainage class	Moderately well drained to well drained
Permeability class	Moderate
Depth to restrictive layer	51–203 cm
Soil depth	51–203 cm
Surface fragment cover ≤3"	0–3%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	11.68–17.53 cm
Calcium carbonate equivalent (0-101.6cm)	0–50%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	6.6–8.8
Subsurface fragment volume ≤3" (0-101.6cm)	0–8%
Subsurface fragment volume >3" (0-101.6cm)	0%

Ecological dynamics

The soils, topographic location, climate, periodic droughts and fire influenced the stabilization of the reference plant community on this site as was the case on most all High Plains ecological sites. This plant community as found by European settlers in the early 1800's developed under the prevailing climate over time along with the soils in their topographic location. Grazing and/or browsing by local and nomadic wildlife influenced the plant community as well. The resulting plant community is a Midgrass/Shortgrass Community (1.1). The sandy loam site is quite possibly the most productive site in the Texas Panhandle. Midgrasses tend to dominate over most of the site with sideoats grama (*Bouteloua curtipendula*) being the overall dominant species. Lesser amounts of vine-mesquite (*Panicum obtusum*), Arizona cottontop (*Digitaria californica*), plains bristlegrass (*Setaria leucopila*), hooded windmillgrass (*Chloris cucullata*), sand dropseed (*Sporobolus cryptandrus*), tumble windmillgrass (*Chloris verticillata*), silver bluestem (*Bothriochloa laguroides*), hairy grama (*Bouteloua hirsuta*), fall witchgrass (*Digitaria cognata*), gummy lovegrass (*Eragrostis curtipedicellata*), black grama (*Bouteloua eriopoda*), and sand muhly (*Muhlenbergia*

arenicola) are present. Little bluestem (*Schizachyrium scoparium*) is present with lesser amounts of Indiangrass (*Sorghastrum nutans*) and sand bluestem (*Andropogon hallii*). In areas where tighter soils occur, the shortgrasses are blue grama (*Bouteloua gracilis*) and buffalograss (*Bouteloua dactyloides*). Some cool-season grasses occur in small amounts such as western wheatgrass (*Pascopyrum smithii*), Canada wildrye (*Elymus canadensis*) and needle & thread (*Hesperostipa comata*). Typically associated forbs include dotted gayfeather (*Liatris punctata*), prairie clover (*Dalea purpurea*), catclaw sensitivebriar (*Mimosa microphylla*), golden dalea (*Dalea aurea*), gaura (*Gaura* spp.), rushpea (*Hoffmanseggia glauca*), Engelmann's daisy (*Engelmannia peristenia*), lyreleaf greeneyes (*Berlandiera lyrata*), sagewort (*Artemisia ludoviciana*), scarlet globemallow (*Sphaeralcea coccinea*), Fendler's penstemon (*Penstemon fendleri*), wild alfalfa (*Psoraleidum tenuiflorum*) and numerous annual forbs. Woody species include sand sagebrush (*Artemisia filifolia*), yucca (*Yucca glauca*), and catclaw mimosa (*Mimosa aculeaticarpa* var. *biuncifera*). Nutrient cycling, the water cycle, watershed protection and biological functions were functioning at their peak.

Fire plays an important role in the function of most plains sites, especially the tallgrass communities. Tallgrasses such as sand bluestem and little bluestem were dependent upon fire to stimulate them and remove old growth that would accumulate on the soil surface. Fire also keeps shrubs from getting too thick. Fire helps to keep a balance between the grasses, forbs, and shrubs. Wildlife habitat is improved by opening up canopies and stimulating forb growth. The deep-rooted species that grow on the site are not easily damaged by fire. Yucca and associated shrubs will usually resprout, but are suppressed for a time allowing grasses to dominate. If periodic fire does not occur, then the yucca and woody plants will slowly increase and with grazing pressure can begin to dominate the site. Since fire is not always available to be applied, practices such as brush management may be necessary from time to time to help keep the community in balance.

Periodic grazing and trampling by migrating herds of bison and resident herds of pronghorn antelope probably occurred during drought periods. However, long rest periods followed once the large herds of bison and antelope moved out of the area, allowing the resilient grassland to re-establish itself and maintain its climax community structure.

The major forces influencing the transition from the reference community are continued overgrazing by livestock and a decrease in the frequency and intensity of fire. As livestock and wildlife numbers increase and grazing use exceeds a plant's ability to sustain defoliation, the more palatable and generally more productive species decline in stature, productivity and density.

Under good management this is one of the most productive sites in the Texas Panhandle and will give good animal performance. Little bluestem and sideoats grama are fairly resistant to grazing pressure but will decline if continuous heavy grazing persists. The tallgrasses are fairly sensitive to overgrazing and will begin to decrease more quickly if continuous abusive grazing occurs for long periods. If excessive grazing pressure continues, ecological retrogression occurs. The tendency of this site, as excessive grazing pressure occurs, is to become a shortgrass dominant site, allowing sideoats grama and little bluestem to give way to blue grama. There will be an increase in perennial and annual forbs, and perennial three-awn will increase. Yucca and sand sagebrush tend to increase. Invasion of mesquite, broom snakeweed and pricklypear may occur on some sites. The decrease in density and stature of the midgrasses and tallgrasses and an increase in shortgrasses and the density of the yucca and woody vegetation bring about a new plant community, the Shortgrass Community (2.1).

In the Shortgrass Community (2.1), the transition back to the reference community requires proper grazing management, brush management, and pest management. Prescribed burning could be used if the conditions allow. The production of vegetation has shifted from mostly herbaceous vegetation to more yucca and woody, although the herbaceous vegetation biomass is still the largest amount. Nutrient cycling, the water cycle, watershed protection and biological functions have changed little.

If long-term abusive grazing continues, a threshold will be crossed to a Degraded Shortgrass Community (3.1). In this degraded state, typical vegetation will be low vigor blue grama with bare areas opening up with annuals filling the voids. Perennial three-awn will increase when the more desirable grasses are weakened and/or removed. Yucca and sand sagebrush will increase dramatically. On sites where mesquite, pricklypear and broom snakeweed have invaded, these woody invaders can increase to the point of dominating the woody canopy. The loss of herbaceous cover and increased bare ground encourages accelerated erosion. Nutrient cycling, the water cycle, watershed protection and biological functions have been severely reduced. The plant community is so degraded that it cannot reverse retrogression without extensive energy and management inputs. Restoration will require

prescribed grazing with rest periods during the growing season, re-seeding bare areas with adapted native grass species, and chemical and/or mechanical brush management and some form of pest management. With the reduced amounts of grass fuel, prescribed burning is usually not an option in this phase.

Recovery can occur fairly rapidly if the competitive plants are controlled and proper grazing management is applied. Full recovery and maintenance of the reference community requires continued proper grazing management as well as occasional brush and pest management.

NOTE: Rangeland Health Reference Worksheets have been posted for this site on the Texas NRCS website (www.tx.nrcs.usda.gov) in Section II of the eFOTG under (F) Ecological Site Descriptions.

STATE AND TRANSITIONAL PATHWAYS


Narrative:

The following diagram suggests some pathways that the vegetation on this site might take. There may be other states not shown on the diagram. This information is intended to show what might happen in a given set of circumstances; it does not mean that this would happen the same way in every instance. Local professional guidance should always be sought before pursuing a treatment scenario.

Changes in the structure and composition of the plant community may be due to management and/or natural occurrences. At some point thresholds are crossed as indicated by the State and Transition Diagram. This suggests that once changes have progressed to a certain point, the plant community has been altered to the extent that a return to the former state is not possible unless some form of energy is applied. These changes take place on all ecological sites. Some sites support communities that are more resistant to change than others. Also, some sites are more resilient and can heal or restore themselves more easily. Usually, changes in management practices alone, such as grazing techniques, will not be sufficient to restore former plant communities. An example of energy input might be the implementation of chemical brush management to decrease the amount of woody/cacti shrubs and increase the amount of grasses and forbs. This shift in community balance could not be brought about with grazing alone. The amount of energy required to bring about a change in plant community balance may vary a great deal depending on the present state and the desired result.

State and transition model

Ecosystem states



T1A - Absence of disturbance and natural regeneration over time, may be coupled with excessive grazing pressure

T2A - Absence of disturbance and natural regeneration over time, may be coupled with excessive grazing pressure

R3A - Adequate rest from defoliation and removal of woody canopy, followed by range land seeding

State 1 submodel, plant communities

1.1.
Midgrass/Shortgrass
Dominant Community

State 2 submodel, plant communities

2.1. Shortgrass
Community

State 3 submodel, plant communities

3.1. Degraded
Shortgrass Community

State 1 Midgrass State

The Midgrass/Shortgrass Community has a good mixture of midgrasses, shortgrasses and lesser amounts of tallgrasses. Grasses make up approximately 90 percent of the plant community. Midgrasses tend to dominate over most of the site with sideoats grama being the overall dominant species. Little bluestem is the dominant tallgrass and blue grama is the dominant shortgrass. The cool-season grasses include western wheatgrass and Canada wildrye and lesser amounts of needle & thread. Perennial forbs make up 3 – 5 percent of the community and shrubs such as sand sagebrush, yucca and catclaw acacia compose of remaining 3 – 5 percent of the overall annual production for the plant community.

Dominant plant species

- sideoats grama (*Bouteloua curtipendula*), grass

Community 1.1 Midgrass/Shortgrass Dominant Community



Figure 8. 1.1 Midgrass/Shortgrass Community

The interpretive or "reference" plant community for this site is a good mixture of midgrasses, shortgrasses and lesser amounts of tallgrasses. Grasses make up approximately 90 percent of the plant community. Midgrasses tend to dominate over most of the site with sideoats grama being the overall dominant species. Little bluestem is the dominant tallgrass and blue grama is the dominant shortgrass. The cool-season grasses include western wheatgrass and Canada wildrye and lesser amounts of needle & thread. Perennial forbs make up 3 – 5 percent of the community and shrubs such as sand sagebrush, yucca and catclaw acacia compose of 3 – 5 percent of the overall plant community. The sites ecological processes are in balance with the environment. Most energy and nutrient cycling is contained in the narrow grass/soil interface and evapo-transpiration is minimal. Maintenance of this community requires continued proper grazing management as well as occasional brush and pest management.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1513	1765	2018
Forb	84	101	112
Shrub/Vine	84	95	112
Tree	–	–	–
Microbiotic Crusts	–	–	–
Total	1681	1961	2242

Figure 10. Plant community growth curve (percent production by month). TX1031, Midgrass/Shortgrass Dominant Community. Growth is predominately mid and shortgrasses with few forbs and shrubs from April through October with peak growth from May to July..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	2	6	9	22	25	12	6	11	4	2	1

State 2 Shortgrass State

Shortgrasses dominate with some midgrasses present. There is an increase in annual forbs and grasses. There are also increasing amounts of yucca, mesquite and broom snakeweed. Percent bare ground is increasing and production is decreasing.

Dominant plant species

- yucca (*Yucca*), shrub
- broom snakeweed (*Gutierrezia sarothrae*), shrub
- buffalograss (*Bouteloua dactyloides*), grass

Community 2.1 Shortgrass Community



Figure 11. 2.1 Shortgrass Community

As retrogression occurs, the tendency of this site is to become a Shortgrass Community (2.1) site. The sideoats grama and little bluestem will give way to blue grama. There will be an increase in perennial and annual forbs, with increasing amounts of yucca and sand sagebrush. Some sites may see an invasion of broom snakeweed, pricklypear and mesquite. The production of vegetation has shifted from mostly herbaceous vegetation to more yucca and woody species, although the herbaceous vegetation biomass is still the largest amount. Nutrient cycling, the water cycle, watershed protection and biological functions have changed little. The transition back to the reference community is reversible with proper grazing management, brush and pest management.

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1132	1216	1351
Shrub/Vine	404	465	532
Forb	146	224	275
Microbiotic Crusts	–	–	–
Tree	–	–	–
Total	1682	1905	2158

Figure 13. Plant community growth curve (percent production by month). TX1009, Shortgrass Dominant. Shortgrass dominant species include blue grama and buffalograss..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	2	3	5	30	30	8	4	8	6	3	1

State 3 Degraded Shortgrass State

The Degraded Shortgrass State for a Sandy Loam site has low vigor shortgrasses and increased amounts of perennial three-awn. Mesquite, yucca, sand sagebrush and broom snakeweed along with other woody species make up greater than 40 percent canopy. The bare areas have filled with annuals and the annual production potential is low.

Dominant plant species

- honey mesquite (*Prosopis glandulosa*), shrub
- yucca (*Yucca*), shrub
- threeawn (*Aristida*), grass

Community 3.1

Degraded Shortgrass Community



Figure 14. 3.1 Degraded Shortgrass Community

In this phase of retrogression a threshold has been crossed to a Degraded Shortgrass Community (3.1). In this degraded state, typical vegetation will be low vigor blue grama; bare areas will open up with annuals filling the voids. Perennial three-awn will invade this site when the more desirable grasses are weakened and/or removed. Yucca, sand sagebrush mesquite, pricklypear and broom snakeweed will increase dramatically. The loss of herbaceous cover and increased bare ground encourages accelerated erosion. Nutrient cycling, the water cycle, watershed protection and biological functions have been severely reduced. The plant community is so degraded that it cannot reverse retrogression without extensive energy and management inputs. Restoration requires prescribed grazing with rest periods during the growing season, re-seeding bare areas with adapted native grass species, and chemical and/or mechanical brush management and some form of pest management. With the reduced amounts of grass fuel, prescribed burning is usually not an option in this phase.

Table 7. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	370	566	740
Shrub/Vine	370	432	493
Forb	62	95	112
Microbiotic Crusts	–	–	–
Tree	–	–	–
Total	802	1093	1345

Figure 16. Plant community growth curve (percent production by month). TX1027, Degraded shortgrasses with annual forbs/grasses. Growth is predominately shortgrasses with forbs and shrubs from April through October with peak growth from May to July..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	2	5	15	30	25	5	3	6	6	2	1

Transition T1A

State 1 to 2

With heavy continuous grazing pressure, no fires, and no brush management practices implemented, the Midgrass/Shortgrass Community will shift to the Shortgrass Community.

Transition T2A

State 2 to 3

With heavy continuous grazing, no fires, no pest management, and no brush management, the Shortgrass

Community will transition to the Degraded Shortgrass Community.

Restoration pathway R3A State 3 to 2

Restoration will require prescribed grazing with rest periods during the growing season, re-seeding bare areas with adapted native grass species, and chemical and/or mechanical brush management and some form of pest management. With the reduced amounts of grass fuel, prescribed burning is usually not an option in this phase.

Conservation practices

Brush Management
Prescribed Grazing
Range Planting
Integrated Pest Management (IPM)

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Tall/Mid/Shortgrasses			841–1121	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	336–448	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	336–448	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	168–224	–
2	Tallgrasses			84–112	
	sand bluestem	ANHA	<i>Andropogon hallii</i>	56–112	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	56–112	–
3	Cool-season grasses			84–112	
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	56–112	–
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	56–112	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	56–112	–
4	Midgrasses			252–336	
	vine mesquite	PAOB	<i>Panicum obtusum</i>	112–280	–
	large-spike bristlegrass	SEMA5	<i>Setaria macrostachya</i>	112–280	–
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	112–224	–
5	Shortgrasses			168–224	
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	112–168	–
	hooded windmill grass	CHCU2	<i>Chloris cucullata</i>	112–168	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	112–168	–
6	Mid/Shortgrasses			84–112	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	22–56	–
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	22–56	–
	silver beardgrass	BOLAT	<i>Bothriochloa laguroides ssp. torreyana</i>	22–56	–
	tumble windmill grass	CHVE2	<i>Chloris verticillata</i>	22–56	–

	fall witchgrass	DICO6	<i>Digitaria cognata</i>	22–56	–
	gummy lovegrass	ERCU	<i>Eragrostis curtipedicellata</i>	22–56	–
	ear muhly	MUAR	<i>Muhlenbergia arenacea</i>	22–56	–
Forb					
7	Forbs			84–112	
	Forb, annual	2FA	<i>Forb, annual</i>	0–56	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	22–56	–
	lyreleaf greeneyes	BELY	<i>Berlandiera lyrata</i>	22–56	–
	golden prairie clover	DAAU	<i>Dalea aurea</i>	22–56	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	22–56	–
	Engelmann's daisy	ENPE4	<i>Engelmannia peristenia</i>	22–56	–
	beeblossom	GAURA	<i>Gaura</i>	22–56	–
	Indian rushpea	HOGL2	<i>Hoffmannseggia glauca</i>	22–56	–
	dotted blazing star	LIPU	<i>Liatris punctata</i>	22–56	–
	Nuttall's sensitive-briar	MINU6	<i>Mimosa nuttallii</i>	22–56	–
	Fendler's penstemon	PEFE	<i>Penstemon fendleri</i>	22–56	–
	elephant grass	PEPU2	<i>Pennisetum purpureum</i>	22–56	–
	slimflower scurfpea	PSTE5	<i>Psoralidium tenuiflorum</i>	22–56	–
	scarlet globemallow	SPCOE	<i>Sphaeralcea coccinea ssp. elata</i>	22–56	–
Shrub/Vine					
8	Shrubs/Vines			84–112	
	sand sagebrush	ARFI2	<i>Artemisia filifolia</i>	56–112	–
	catclaw mimosa	MIACB	<i>Mimosa aculeaticarpa var. biuncifera</i>	56–112	–
	yucca	YUCCA	<i>Yucca</i>	56–112	–

Animal community

Native animals that occupy this site include scaled quail, pronghorn, lesser prairie chicken, and various small mammals and grassland birds. The site lacks woody cover for deer and turkey. Predators such as coyotes and bobcats may utilize the site for hunting prey and to hide during the day. If the site is a Shortgrass Dominant Community (2), then diversity is increased and the wildlife habitat may become more desirable.

Hydrological functions

This site captures nearly most of the water that falls on it, provided the vegetation is in good condition. The sandy loam soil infiltrates water fairly rapidly. There is not significant runoff if cover is good. With poor cover, runoff is significant and small gullies can develop.

Recreational uses

Hunting, Camping, Hiking, Bird watching, Photography, Horseback Riding.

Wood products

None.

Other products

None.

Other information

None.

Inventory data references

NRCS FOTG – Section II of the FOTG Range Site Descriptions and numerous historical accounts of vegetative conditions at the time of early settlement in the area were used in the development of this site description. Vegetative inventories were made at several site locations for support documentation.

Inventory Data References (documents):

NRCS FOTG – Section II - Range Site Descriptions

NRCS Clipping Data summaries over a 20 year period

Other references

1. Archer S. 1994. Woody plant encroachment into southwestern grasslands and savannas: rates, patterns and proximate causes. In Ecological implications of livestock herbivory in the West, Ed M Vavra, W Laycock, R Pieper, pp13-68, Denver, CO: society for Range Management
2. Gould F. 1978. Common Texas Grasses: an illustrated guide. College Station, TX: Texas A & M Press.
3. Hatch, Brown and Ghandi, Vascular Plants of Texas (An Ecological Checklist)
4. Heischmidt RK, Stuth, Eds. 1991 Grazing Management: an ecological perspective. Portland OR: Timberline Press
5. North Rolling Plains RC&D, NRCS, and GLCI. 2006 edition. Common Rangeland Plants of the Texas Panhandle.
6. Scifres CJ, Hamilton WT. 1993. Prescribed burning for Brushland management: the South Texas example. College Station, TX: Texas A & M Press.
7. Natural Resources Conservation Service - Range Site Descriptions
8. USDA-Natural Resources Conservation Service - Soil Surveys & Website soil database

The following individuals assisted with the development of this site description:

Clint Rollins – Rangeland Management Specialist- NRCS; Amarillo, Texas

Justin Clary – Rangeland Management Specialist – NRCS; Temple, Texas

Stan Bradbury – Rangeland Management Specialist - NRCS; Lubbock, Texas

Kelly Attebury – Soil Scientist- NRCS; Lubbock, Texas

Charles Coffman – Biologist - NRCS; Lubbock, Texas

Contributors

Brent Conlin, Area Conservationist, Amarillo

Duckworth-Cole, Inc., Bryan, Texas

T. Craig Byrd

Approval

Bryan Christensen, 9/11/2023

Acknowledgments

Site Development and Testing Plan

Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document.

Annual reviews of the Project Plan are to be conducted by the Ecological Site Technical Team.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Bradbury, Zone RMS, NRCS, Lubbock, Texas
Contact for lead author	806-791-0581
Date	09/04/2007
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None to slight.

2. **Presence of water flow patterns:** None to slight.

3. **Number and height of erosional pedestals or terracettes:** None to slight.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 20-25%.

5. **Number of gullies and erosion associated with gullies:** None to slight.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None to slight.

7. **Amount of litter movement (describe size and distance expected to travel):** None to slight.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Moderate resistance to surface erosion.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Fine sandy loam, friable surface and medium SOM.

-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Basal cover and density with moderate interspaces should make rainfall impact minimal. This site has moderate permeability, runoff is slow and available water holding capacity is high.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm-season midgrasses = Warm-season shortgrasses >
- Sub-dominant: Warm-season tallgrasses > Cool-season grasses >
- Other: Shrubs/Vines > Forbs
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Grasses due to their growth habit will exhibit some mortality and decadence, though minimal.
-
14. **Average percent litter cover (%) and depth (in):** Litter is dominantly herbaceous.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1,400 to 1,800 pounds per acre.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Sand Sagebrush and Yucca can become invasive.
-
17. **Perennial plant reproductive capability:** All plant species should be capable of reproduction except during prolonged drought conditions, heavy natural herbivory or intense wildfires.
-