

Ecological site R078AY117TX Clayey Upland 25-28" PZ

Last updated: 9/12/2023 Accessed: 05/09/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Lem Creswell, Zone RMS, NRCS, Weatherford, Texas
Contact for lead author	817-596-2865
Date	11/19/2008
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills: None.
- 2. Presence of water flow patterns: Deposition or erosion is uncommon during normal rainfall events, but may occur in limited areas during intense rainfall events.
- 3. Number and height of erosional pedestals or terracettes: Pedestals or terracettes are uncommon for this site.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): Expect no more than 10% bare ground scattered randomly throughout the site.
- 5. Number of gullies and erosion associated with gullies: None.
- 6. Extent of wind scoured, blowouts and/or depositional areas: None.
- 7. Amount of litter movement (describe size and distance expected to travel): Little or no litter movement or

deposition during normal rainfall events; however, litter of all sizes may move long distances depending on obstructions under intense storm events.

- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Soil surface in HCPC is resistant to erosion. Stability range is expected to be 5-6.
- Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Up to 48" thick with dark grayish brown clay with generally moderate fine granular and subangular blocky structure. SOM approximately 1-6%. See Soil Survey for specific soil.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Areas of dense, low growing, sod forming grasses may contribute to moderately high runoff rate and reduced infiltration rates.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Warm-season midgrasses >>

Sub-dominant: Warm-season shortgrasses >

Other: Cool-season grasses > Forbs > Shrubs/Vines > Trees

Additional:

- Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Perennial grasses will naturally exhibit a minor amount (less than 5%) of senescence and some mortality every year.
- 14. Average percent litter cover (%) and depth (in): Litter is primarily herbaceous.
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): 1800-3400 pounds per acre.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not

invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Mesquite, pricklypear, bermudagrass, Johnsongrass, King Ranch bluestem.

17. **Perennial plant reproductive capability:** All perennial species should be capable of reproducing every year unless disrupted by extended drought, overgrazing, wildfire, insect damage, or other events occuring immediately prior to, or during the reproductive phase.