

Ecological site R078BY090TX Shallow Clay 19-26" PZ

Last updated: 9/15/2023 Accessed: 05/07/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Stan Bradbury, Zone RMS, NRCS, Lubbock, Texas			
Contact for lead author	806-791-0581			
Date	09/04/2007			
Approved by	Bryan Christensen			
Approval date				
Composition (Indicators 10 and 12) based on	Annual Production			

nc	ndicators					
1.	Number and extent of rills: Slight to moderate.					
2.	Presence of water flow patterns: Slight to moderate.					
3.	Number and height of erosional pedestals or terracettes: Slight to moderate.					
4.	Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 35 to 40% bare ground.					
5.	Number of gullies and erosion associated with gullies: Slight to moderate.					
6.	Extent of wind scoured, blowouts and/or depositional areas: None to slight.					
7.	Amount of litter movement (describe size and distance expected to travel): Slight to moderate.					

Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Water erosion hazards are moderate to severe.
Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Clay to clay loam surfaces; moderate fine subangular blocky structure; very hard; firm; very sticky, plastic; many fine and common medium roots; moderate SOM.
Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Low vegetative cover and percent slopes makes this site susceptible to erosion. This site is a very slowly permeable soil, runoff is medium to high depending on slope and available water holding capacity is moderate to high.
Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):
Dominant: Warm-season shortgrasses >
Sub-dominant: Warm-season midgrasses >
Other: Forbs > Warm-season tallgrasses > Shrubs/Vines > Trees
Additional:
Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Plant mortality and decadence is minimal.
Average percent litter cover (%) and depth (in): Litter is dominantly herbaceous.
Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): 650 to 1,400 pounds per acre.
Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Mesquite, lotebush and pricklypear can be invasive.
