

Ecological site R078BY701TX Shallow Sandstone 19-26" PZ

Last updated: 9/15/2023 Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 078B-Central Rolling Red Plains, Western Part

MLRA 78B is characterized by strongly dissected, rolling plains with prominent ridges and valleys and rolling to steep irregular topography. Loamy soils are generally well drained, range from shallow to deep, and developed in sediments of Triassic and Permian age.

LRU notes

NA

Classification relationships

This ecological site is correlated to soil components at the Major Land Resource Area (MLRA) level which is further described in USDA Ag Handbook 296.

Ecological site concept

These sites occur on shallow soils on uplands. Reference vegetation includes midgrasses and shortgrasses with forbs and scattered woody plants. Abusive grazing practices can lead to a shift in the plant community. Without fire

or alternative brush management, woody species may increase on the site.

Associated sites

| R078BY076TX | Gyp 19-26" PZ Shallow soils over gypsum |
|-------------|--|
| R078BY084TX | Rough Breaks 19-26" PZ Shallow soils on breaks |
| R078BY090TX | Shallow Clay 19-26" PZ Shallow clay soils on uplands |
| R078BY092TX | Very Shallow Clay 19-26" PZ Very shallow soils on uplands |

Table 1. Dominant plant species

| Tree | (1) Juniperus pinchotii |
|------------|--|
| Shrub | Not specified |
| Herbaceous | (1) Schizachyrium scoparium(2) Bouteloua curtipendula |

Physiographic features

This is an upland site with very shallow soils mixed with outcrops of Permian and Triassic sandstone. It occurs on ridges, low hills, side slopes along drainageways and along escarpments. Slopes are moderate to moderately steep. There are often sandstone fragments mixed with shale and siltstone on the soil surface and often sandstone bedrock will be showing. The site may be small to relatively large, varying from 50 to over 1000 acres in size. The actual soils are so intermingled with rock outcrops that they cannot be separated.

This site may be encountered at any aspect of the landscape.

| Landforms | (1) Plains > Ridge (2) Plains > Escarpment (3) Plains > Hill |
|--------------------|---|
| Runoff class | Low to medium |
| Flooding frequency | None |
| Ponding frequency | None |
| Elevation | 411–975 m |
| Slope | 1–20% |
| Water table depth | 152–203 cm |
| Aspect | Aspect is not a significant factor |

Table 2. Representative physiographic features

Climatic features

The climate of the western rolling plains is dry, sub-humid with hot summers and mild winters. Temperatures often reach 100 degrees F for several consecutive days during summer. Cold spells with temperatures less than 20 degrees F only last short periods of time. The soil is not frozen below the 3-inch depth for more than 2 to 3 days. Humidity is low during the winter and early spring months. Sometimes relative humidity is high enough to make summer days seem uncomfortable. Most of the precipitation comes in the form of rain and that in the spring and early summer principally. May is the wettest month followed by June. July and August are dryer and much hotter. Rainfall often comes as intense showers of relatively short duration. Rainfall rate per hour is often high and runoff is significant. Infiltration is diminished due to lack of opportunity time. The growing season begins in April and ends with the first killing frost in November. There is little snowfall with the average being about 10 inches. Rainfall

averages about 22 inches.

There is a 70% chance that yearly precipitation will fall between 16 and 24 inches. About 55% of the time, the yearly rainfall is below the mean. Dry spells during the growing season are common and long-term droughts occur in cycles of about 20 years. Native vegetation is principally warm season.

| Table 3. Representative climatic features |
|---|
|---|

| Frost-free period (characteristic range) | 189-199 days |
|--|--------------|
| Freeze-free period (characteristic range) | 202-222 days |
| Precipitation total (characteristic range) | 584 mm |
| Frost-free period (actual range) | 184-203 days |
| Freeze-free period (actual range) | 201-223 days |
| Precipitation total (actual range) | 559-584 mm |
| Frost-free period (average) | 193 days |
| Freeze-free period (average) | 213 days |
| Precipitation total (average) | 584 mm |

Climate stations used

- (1) WELLINGTON [USC00419565], Wellington, TX
- (2) MATADOR [USC00415658], Matador, TX
- (3) SPUR [USC00418566], Spur, TX
- (4) SNYDER [USC00418433], Snyder, TX
- (5) ROBERT LEE [USC00417669], Robert Lee, TX

Influencing water features

Rapid runoff due to impermeable soil material and steep slopes.

There are no streams or wetlands associated with this ecological site.

Wetland description

NA

Soil features

Soils are very shallow, well drained and calcareous with limited soil development. Rock outcrops are intermingled with soil material. Geologic erosion has stripped away soil material exposing bedrock in many places. Layers of sandstone may be quite thick and are interlaid with sandy to clayey redbeds. The soil is generally loam to fine sandy loam in texture with stony or gravelly modifier. Underlying material is pale brown strongly cemented sandstone. Available water holding capacity is low and fertility is low. Plant growth and production is limited due to shallow depth. Cracks in underlying bedrock can allow water to penetrate. Shrubs and grasses with deep root systems may be able to access some of this moisture.

Major Soil Taxonomic Units correlated to this site include: Latom soils and Latom Rock Outcrop.

Table 4. Representative soil features

| Parent material | (1) Residuum-sandstone |
|-----------------|---|
| Surface texture | (1) Fine sandy loam (2) Gravelly fine sandy loam |

| Drainage class | Well drained |
|--|-----------------------------|
| Permeability class | Moderately slow to moderate |
| Depth to restrictive layer | 10–51 cm |
| Soil depth | 10–51 cm |
| Surface fragment cover <=3" | 5–40% |
| Surface fragment cover >3" | 1–5% |
| Available water capacity (0-101.6cm) | 1.02–3.05 cm |
| Calcium carbonate equivalent (0-101.6cm) | 0–10% |
| Electrical conductivity (0-101.6cm) | 0–2 mmhos/cm |
| Sodium adsorption ratio (0-101.6cm) | 0 |
| Soil reaction (1:1 water) (0-101.6cm) | 7.9–8.4 |
| Subsurface fragment volume <=3" (Depth not specified) | 3–22% |
| Subsurface fragment volume >3" (Depth not specified) | 1–3% |

Ecological dynamics

This reference plant community is dominated by tall and midgrasses with a good perennial forb population and a variety of short shrubs. Some juniper occurs on the site along with occasional mesquite. The site occurs in rougher terrain and is found along escarpments and drainageways within the rolling red plains. The parent material is sandstone and rock outcrops are mixed with soil material. Production potential is limited due to shallow depth of soil and by steep terrain.

Slopes vary from moderate to steep but usually fall into the 5 to 20 % slope range. Runoff is rapid and infiltration limited. Evaporation is high. Plant basal cover is generally sparse but varies. Cracks and fissures in the underlying sandstone trap water in some locations and plants can sometimes access this water. Lack of inherent fertility may affect palatability of plants. This site is not as prone to over utilization as some sites with deeper, more fertile soils. Because of it's diversity of plant species, the site is utilized by a variety of wildlife. There are several desirable forb and shrub species that afford good quality browse.

Natural fire shaped the ecology of this site as with most plains grassland sites. Grasses such as little bluestem (*Schizachyrium scoparium*), sand bluestem (*Andropogon hallii*), and sideoats grama (*Bouteloua curtipendula*) are stimulated by periodic fire. Redberry juniper (*Juniperus pinchotii*) is suppressed but usually resprouts and attains pre-burn status within a few years. But in general, shrubs are suppressed sufficiently to allow for a grassland aspect to exist. Cacti species (Opuntia spp) are damaged by fire provided sufficient fuel exists to insure an intense burn. Wildfires are thought to have occurred perhaps every 8 to 15 years in the region depending on fuel accumulation and random lightning strikes. Native Americans sometimes purposely set fires to influence grazing by large herbivores. Due to relatively thin vegetative cover, limited fuel loads and protective terrain, some susceptible shrub species have been naturally protected from wildfire damage on some portions of the site. If not naturally burned, juniper will increase on the site and can form a significant canopy cover in some areas. Overgrazing by livestock will reduce the more palatable grass and forb species. Cattle will generally not spend as much time on this site but will selectively graze when there.

The main influences that drive the ecology of the site are: climatic variability, occurrence of / or lack of fire, shallow soils, and selective grazing or browsing by livestock and wildlife. Geologic erosion can have an effect on portions of the site. The site can deteriorate through mismanagement and is sensitive to abuse.

Plant Communities and Transitional Pathways (diagram):

The following diagram suggests some pathways that the vegetation on this site might take. There may be other states not shown on the diagram. This information is intended to show what might happen in a given set of circumstances; it does not mean that this would happen the same way in every instance. Local professional guidance should always be sought before pursuing a treatment scenario.

State and transition model

Ecosystem states



- T1A Absence of disturbance and natural regeneration over time, may be coupled with excessive grazing pressure
- R2A Adequate rest from defoliation and removal of woody canopy, followed by reintroduction of historic disturbance regimes
- T2A Absence of disturbance and natural regeneration over time, may be coupled with excessive grazing pressure

State 1 submodel, plant communities



State 2 submodel, plant communities



State 3 submodel, plant communities



State 1 Mixed-grass/Shrubland State

The interpretive or "reference" plant community for the Tall/Midgrass/Shrubs Community (1.1) is a balanced mixture of tallgrasses such as little bluestem and sand bluestem; midgrasses such as sideoats grama and a few shortgrasses such as blue grama, buffalograss, and hairy grama. There is also a variety of perennial forbs including

dotted gayfeather, plains blackfoot daisy, heath aster, western ragweed, gaura, and catclaw sensitivebriar. There are also scattered short woody shrubs in this community including feather dalea, skunkbush sumac, and vine ephedra. A few scattered redberry juniper, mesquite, and hackberry trees also occur. The Mid/Tallgrass/Shrubs Community (1.2) occurs when tallgrasses such as little bluestem and sand bluestem decrease and midgrasses such as sideoats grama, sand dropseed, and silver bluestem dominate. There is an increase in perennial three-awns. Some low growing shrubs such as feather dalea, pricklypear, elbowbush, and yucca also occur. Redberry juniper increases to ten to fifteen percent woody canopy.

Dominant plant species

- Pinchot's juniper (Juniperus pinchotii), tree
- little bluestem (Schizachyrium scoparium), grass
- sideoats grama (Bouteloua curtipendula), grass

Community 1.1 Tall/Midgrass/Shrubs Community



Figure 8. 1.1 Tall/Midgrass/Shrubs Community

The interpretive or "reference" plant community for this site is a balanced mixture of tallgrasses - little bluestem (*Schizachyrium scoparium*), sand bluestem (*Andropogon hallii*); midgrasses - sideoats grama (*Bouteloua curtipendula*) and a few short grasses – blue grama (*Bouteloua gracilis*), buffalograss (Buchloe dactyloides) and hairy grama (*Bouteloua hirsuta*), a variety of perennial forbs – dotted gayfeather, (*Liatris punctata*), plains blackfoot daisy (*Melampodium leucanthum*), heath aster (Aster ericoides), western ragweed (*Ambrosia psilostachya*), gaura (Gaura spp), catclaw sensitivebriar (*Mimosa nuttallii*), and scattered short woody shrubs – feather dalea (*Dalea formosa*), skunkbush sumac (*Rhus trilobata*), and vine epedra (*Ephedra pedunculata*). A few scattered redberry juniper (*Juniperus pinchotii*), mesquite (*Prosopis glandulosa*) and hackberry (Celtis spp.) trees also occur.

Table 5. Annual production by plant type

| Plant Type | Low (Kg/Hectare) | Representative Value (Kg/Hectare) | High (Kg/Hectare) |
|--------------------|---------------------|--------------------------------------|----------------------|
| Grass/Grasslike | 560 | 897 | 1121 |
| Tree | 28 | 67 | 112 |
| Shrub/Vine | 56 | 67 | 101 |
| Forb | 56 | 112 | 101 |
| Microbiotic Crusts | 11 | 22 | 28 |
| Total | 711 | 1165 | 1463 |

Figure 10. Plant community growth curve (percent production by month). TX2047, Tall & Midgrass with Shrubs community. mid and tallgrasses, forbs and shrubs..

| Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1 | 4 | 6 | 10 | 20 | 24 | 10 | 5 | 10 | 7 | 2 | 1 |

Community 1.2 Mid/Tallgrass/Shrubs Community



Figure 11. 1.2 Mid/Tallgrass/Shrubs Community

This community represents the first phase in the transition of the Tall/Midgrass/Shrubs Community (1.1) toward the Mid/Shortgrass/Shrubs Community (1.2). Tallgrasses such as little bluestem and sand bluestem decrease and midgrasses such as sideoats grama, sand dropseed (*Sporobolus cryptandrus*) and silver bluestem (*Bothriochloa laguroides*) dominate with some increase in perennial three-awns (Aristida spp.) with some low growing shrubs such as feather dalea, pricklypear (Opuntia spp.), elbowbush (*Forestiera pubescens*) and yucca (*Yucca glauca*). Redberry juniper increases to 10-15% woody canopy. Proper grazing and the use of prescribed burning can maintain or possibly restore this community to the reference community and prevent the transition toward the Shortgrass/Shrubland State. With further heavy continued grazing and fire suppression, the community will degrade toward the Short/Midgrass/Shrubs Community (2.1).

| Plant Type | Low (Kg/Hectare) | Representative Value (Kg/Hectare) | High (Kg/Hectare) |
|--------------------|---------------------|--------------------------------------|----------------------|
| Grass/Grasslike | 448 | 673 | 897 |
| Tree | 39 | 84 | 168 |
| Shrub/Vine | 84 | 95 | 112 |
| Forb | 45 | 56 | 67 |
| Microbiotic Crusts | 6 | 11 | 22 |
| Total | 622 | 919 | 1266 |

Table 6. Annual production by plant type

Figure 13. Plant community growth curve (percent production by month). TX2048, Midgrasses and forbs. warm-season midgrasses and forbs..

| Jan | Feb | Mar | Apr | Мау | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 0 | 1 | 3 | 5 | 20 | 25 | 17 | 8 | 15 | 4 | 1 | 1 |

Pathway 1.1A Community 1.1 to 1.2



Tall/Midgrass/Shrubs Community Mid/Tallgrass/Shrubs

Community

With heavy continuous grazing and no fires, the Tall/Midgrass/Shrubs Community will shift to the Mid/Tallgrass/Shrubs Community.

Pathway 1.2A Community 1.2 to 1.1



Mid/Tallgrass/Shrubs Community Tall/Midgrass/Shrubs Community

With Prescribed Grazing and Prescribed Burning, the Mid/Tallgrass/Shrubs Community can be reverted back to the Tall/Midgrass/Shrubs Community.

Conservation practices

Prescribed Burning

Prescribed Grazing

State 2 Shortgrass/Shrubland State

The Short/Midgrass/Shrubs Community (2.1) occurs when little and sand bluestem decrease significantly, allowing midgrasses such as perennial three-awn, silver bluestem, and slim tridens to increase. Sideoats grama decreases in dominance and vigor. Shortgrasses such as buffalograss, hairy grama, and hairy tridens increase in density. Climax forbs decrease in frequency and diversity. Woody shrubs such as feather dalea, elbowbush and skunkbush sumac increase. Redberry juniper increases to 20% woody canopy.

Dominant plant species

- juniper (Juniperus), tree
- featherplume (Dalea formosa), shrub
- skunkbush sumac (*Rhus trilobata*), shrub
- buffalograss (Bouteloua dactyloides), grass

Community 2.1 Short/Midgrass/Shrubs Community



Figure 14. 2.1 Short/Midgrass/Shrubs Community

The Short/Midgrass/Shrubs Community (2.1) occurs when little and sand bluestem decrease significantly, allowing midgrasses such as perennial three-awn, silver bluestem, and slim tridens (Tridens mutica) to increase. Sideoats grama decreases in dominance and vigor. Shortgrasses such as buffalograss, hairy grama, and hairy tridens (*Erioneuron pilosum*) increase in density. Climax forbs decrease in frequency and diversity. Woody shrubs such as feather dalea, elbowbush and skunkbush sumac increase. Redberry juniper increases to 20% woody canopy. More intense management using Prescribed Burning and Prescribed Grazing will be necessary to prevent further degradation to the Shrubs/Shortgrass Community (3.1). Brush management may be necessary to prevent competition from juniper.

Table 7. Annual production by plant type

| Plant Type | Low (Kg/Hectare) | Representative Value (Kg/Hectare) | High (Kg/Hectare) |
|--------------------|---------------------|--------------------------------------|----------------------|
| Grass/Grasslike | 336 | 504 | 673 |
| Tree | 56 | 146 | 224 |
| Shrub/Vine | 90 | 112 | 135 |
| Forb | 22 | 34 | 45 |
| Microbiotic Crusts | _ | 6 | 11 |
| Total | 504 | 802 | 1088 |

Figure 16. Plant community growth curve (percent production by month). TX2049, shortgrass, shrubs & forbs community. warm- and cool-season forbs, shrubs and shortgrasses..

| Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 0 | 1 | 3 | 8 | 16 | 25 | 5 | 5 | 10 | 16 | 8 | 3 |

State 3 Shrubland State

The Shrubs/Shortgrass Community (3.1) occurs when tallgrasses and midgrasses decrease to the extent that shortgrasses such as hairy grama, hairy tridens, and buffalograss dominate. Forbs decrease in abundance, contributing to increased bare ground, accelerated soil erosion and an increase of juniper more than 25% canopy. Broom snakeweed may increase, further deteriorating the ecology of the site.

Dominant plant species

- Pinchot's juniper (Juniperus pinchotii), tree
- broom snakeweed (*Gutierrezia sarothrae*), shrub
- hairy grama (Bouteloua hirsuta), grass
- buffalograss (Bouteloua dactyloides), grass

Community 3.1 Shrubs/Shortgrass Community



Figure 17. 3.1 Shrubs/Shortgrass Community

The Shrubs/Shortgrass Community (3.1) occurs when tallgrasses and midgrasses decrease to the extent that shortgrasses such as hairy grama, hairy tridens, and buffalograss dominate. Forbs decrease in abundance, contributing to increased bare ground, accelerated soil erosion and an increase of juniper more than 25% canopy. Broom snakeweed (*Gutierrezia sarothrae*) may increase, further deteriorating the ecology of the site. Production of forage available for grazing is less than 25% of the reference plant community. Mechanical Brush Management, Range Planting, and Prescribed Burning followed by prescribed grazing will be mandatory for any restoration of this site to the Mixed-grass/Shrubland State.

| Plant Type | Low (Kg/Hectare) | Representative Value (Kg/Hectare) | High (Kg/Hectare) |
|--------------------|---------------------|--------------------------------------|----------------------|
| Grass/Grasslike | 168 | 280 | 392 |
| Tree | 78 | 168 | 280 |
| Shrub/Vine | 101 | 135 | 168 |
| Forb | 11 | 22 | 34 |
| Microbiotic Crusts | _ | 2 | 6 |
| Total | 358 | 607 | 880 |

Table 8. Annual production by plant type

Figure 19. Plant community growth curve (percent production by month). TX2050, Shrubs and Shortgrass community. shortgrasses, warm- and cool-season forbs, and short shrubs..

| Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 0 | 1 | 3 | 8 | 16 | 25 | 5 | 5 | 10 | 16 | 8 | 3 |

Transition T1A State 1 to 2

With Heavy Continuous Grazing, No Fires, and No Brush Management, the Mixed-grass/Shrubland State can transition into the Shortgrass/Shrubland State.

Restoration pathway R2A State 2 to 1

With Prescribed Grazing, Brush Management and Prescribed Burning conservation practices, the Shortgrass/Shrubland State can be restored back to the Mixed-grass/Shrubland State.

Conservation practices

Brush Management

Prescribed Burning

Prescribed Grazing

Transition T2A State 2 to 3

With Heavy Continuous Grazing, No Fires, No Brush Management, and Brush Invasion, the Mixed-grass/Shrubland State can transition into the Shrubland State.

Additional community tables

Table 9. Community 1.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production (Kg/Hectare) | Foliar Cover (%) |
|-------|---------------------|--------|---|-----------------------------------|---------------------|
| Grass | /Grasslike | - | | | |
| 1 | tall/midgrasses | | | 448–588 | |
| | sideoats grama | BOCUC | Bouteloua curtipendula var. caespitosa | 224–588 | _ |
| | little bluestem | SCSC | Schizachyrium scoparium | 224–588 | _ |
| 2 | mid/shortgrasses | _ | | 112–224 | |
| | Wright's threeawn | ARPUW | Aristida purpurea var. wrightii | 0–224 | _ |
| | black grama | BOER4 | Bouteloua eriopoda | 0–224 | _ |
| | blue grama | BOGR2 | Bouteloua gracilis | 0–224 | _ |
| | hairy grama | BOHI2 | Bouteloua hirsuta | 0–224 | _ |
| | silver beardgrass | BOLAT | Bothriochloa laguroides ssp. torreyana | 0–224 | _ |
| | hairy woollygrass | ERPI5 | Erioneuron pilosum | 0–224 | _ |
| | sand dropseed | SPCR | Sporobolus cryptandrus | 0–224 | _ |
| | slim tridens | TRMU | Tridens muticus | 0–224 | _ |
| 3 | tallgrass | - | | 140–224 | |
| | sand bluestem | ANHA | Andropogon hallii | 0–224 | _ |
| | Indiangrass | SONU2 | Sorghastrum nutans | 0–224 | _ |
| 4 | mid/shortgrass | _ | | 39–84 | |
| | buffalograss | BODA2 | Bouteloua dactyloides | 0–84 | _ |
| | fall witchgrass | DICO6 | Digitaria cognata | 0–84 | _ |
| | tobosagrass | PLMU3 | Pleuraphis mutica | 0–84 | _ |
| | plains bristlegrass | SEVU2 | Setaria vulpiseta | 0–84 | _ |
| Forb | | - | | | |
| 5 | forbs | | | 56–112 | |
| | Forb, annual | 2FA | Forb, annual | 0–112 | _ |
| | Cuman ragweed | AMPS | Ambrosia psilostachya | 0–112 | _ |
| | white sagebrush | ARLUM2 | Artemisia ludoviciana ssp. mexicana | 0–112 | _ |
| | woollypod milkweed | ASER | Asclepias eriocarpa | 0–112 | _ |
| | yellow sundrops | CASE12 | Calylophus serrulatus | 0–112 | |

| rose heath CHER2 Chaetopapa ericoides 0-112 blacksamson echinaces ECAN2 Echinacea angustifolia 0-112 spearleaf buckwheat ERLO4 Eriogonum ionchophyllum 0-112 warty spurge EUSP Euphorbis spathulata 0-112 shaggy dwarf morning- glory EVNU Evolvulus nuttallianus 0-112 western blanketflower GASP Gaillardia spathulata 0-112 hoary false goldenaster HECA8 Heterotheca canescens 0-112 collegeflower HYFL Hymenopappus flavescens 0-112 dotted blazing star LIPU Liaris punctala 0-112 dotted blazing star LIPU Liaris punctala 0-112 hoary blackfoot MEE12 Melampodium cinereum 0-112 blazingstar MENZ Menzelia 0-112 huttall's sensitive-briar MINU6 Mimosa nuttalii 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 white milkwort PAV4 Paronychia viginica 0-112 white milkwort POALA Polygala alba 0 | | | | | | |
|---|-------|--------------------------------|-------|---|--------|---|
| blacksamson echinacea ECAN2 Echinacea angustifolia 0-112 warty spurge EUSP Euphorbia spathulata 0-112 warty spurge EUSP Euphorbia spathulata 0-112 warty spurge EUSP Euphorbia spathulata 0-112 warty spurge GASP Gallardia spathulata 0-112 warty spurge HSCA8 Heterotheca canescens 0-112 hoary false goldenaster HECA8 Heterotheca canescens 0-112 collegeflower HYEL Hymenopappus flavescens 0-112 dotted blazing star LIPU Listis punctata 0-112 hoary blackfoot MELE2 Melampodium cincerum 0-112 blazingstar MENTZ Mentzelia 0-112 hoary blackfoot MELE2 Melampodium cincerum 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nalwort POAL4 Paronychia virginica 0-112 yel | | rose heath | CHER2 | Chaetopappa ericoides | 0–112 | _ |
| spearleaf buckwheat ERLO4 Eriogonum lonchophyllum 0-112 warty spurge EUSP Euphobia spathulata 0-112 shaggy dwarf morning- glory GNU Evolvulus nuttallianus 0-112 western blanketflower GASP Gaillardia spathulata 0-112 hoary false goldenaster HECA8 Heterotheca carescens 0-112 collegeflowor HYL Hymenopappus flavescens 0-112 Gordon's bladderpod LEGO Legourolla gordonii 0-112 dotted blazing star LIPU Liafris punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MECI Melampodium cinereum 0-112 vellains blackfoot MECI Melampodium cinereum 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAV4 Paronychia virginica 0-112 yellow nailwort POL4 Polygala alba 0-112 white milkwort POA4 Polygala alba 0-112 | | blacksamson echinacea | ECAN2 | Echinacea angustifolia | 0–112 | _ |
| warty spurge EUSP Euphorbia spathulata 0-112 shaggy dwarf morning- glory EVNU Evolvulus nuttallianus 0-112 western blanketflower GASP Gaillardia spathulata 0-112 hoary false goldenaster HECA8 Heterotheca canescens 0-112 trailing krameria KRLA Krameria lanceolata 0-112 Gordon's bladderpod LEGO Lesquerelle gordonii 0-112 dotted blazing star LIPU Liatris punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MELZ Melampodium cinereum 0-112 blazingstar MENTZ Menzolia 0-112 vening primose OENOT Oenothera 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 white milkwort POAL4 Polygal alba 0-112 simflower scurfpea PSTE5 Psoralidum tenuiflorum 0-112 white milkwort POAL4 Polygal alba 0-112 start sragitolia 0-112 0-112 0-112 start scurfpea PSTE5 Psoralidum tenuiflorum 0-11 | | spearleaf buckwheat | ERLO4 | Eriogonum lonchophyllum | 0–112 | _ |
| shaggy dwarf morning- glory EVNU Evolvulus nuttallianus 0–112 western blanketflower GASP Gallardia spathulata 0–112 hoary false goldenaster HECA8 Heterotheca canescens 0–112 collegeflower HYFL Hymenopappus flavescens 0–112 trailing krameria KRLA Krameria lanceolata 0–112 dotted blazing star LIPU Listris punctate 0–112 hoary blackfoot MECI Melampodium cinereum 0–112 blazingstar MENTZ Mentzolia 0–112 vening primrose OENOT Oenothera 0–112 upriple locoweed OXLA3 Oxytropis lambertii 0–112 yellow nailwort PAVI4 Paronychia virginica 0–112 yellow nailwort PAVI4 Paronychia virginica 0–112 white milkwort PASES Posralidum tenuifforum 0–112 gellow nailwort PAVI4 Paronychia virginica 0–112 gellow nailwort PAVI4 Paronychia virginica 0–112 simflower scurfpea PSTE5 Psoralidum tenuifforum 0– | | warty spurge | EUSP | Euphorbia spathulata | 0–112 | - |
| western blanketflower GASP Gaillardia spathulata 0-112 hoary false goldenaster HECAB Heterotheca canescens 0-112 collegeflower HYFL Hymenopappus flavescens 0-112 trailing krameria KRLA Krameria lanceolata 0-112 dotted blazing star LIPU Listur punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MELE2 Melampodium cinereum 0-112 blazingstar MENTZ Menzelia 0-112 vening primose OENOT Oenothera 0-112 evening primose OENOT Oonthera 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 yellow nailwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 stemmy four-nerve daisy TESC Ternauris scaposa 0-112 stemmy four-nerve daisy TESC2 Ternauris scaposa 0-112 stemturburu | | shaggy dwarf morning- glory | EVNU | Evolvulus nuttallianus | 0–112 | - |
| Image: Section of the sectin of the section of the | | western blanketflower | GASP | Gaillardia spathulata | 0–112 | - |
| collegeflower HYFL Hymenopappus flavescens 0-112 Itailing krameria KRLA Krameria lanceolata 0-112 Gordon's bladderpod LEGO Lesquerella gordonii 0-112 dotted blazing star LIPU Liatris punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MELE2 Melampodium leucanthum 0-112 blazingstar MINUG Minzelia 0-112 Nuttall's sensitive-briar MINUG Minusea nuttallii 0-112 evening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpaa PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragifolia 0-112 stermty four-nerve daisy TESC2 Tetraeuris scaposa 0-112 | | hoary false goldenaster | HECA8 | Heterotheca canescens | 0–112 | - |
| trailing krameria KRLA Krameria lanceolata 0-112 Gordon's bladderpod LEGO Lesquerella gordonii 0-112 dotted blazing star LIPU Liatris punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MELE2 Melampodium leucanthum 0-112 blazingstar MENTZ Mentzelia 0-112 vening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort POAL4 Pornychia virginica 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 skemmy four-nerve daisy TESC2 Teraneuris scaposa 0-112 Shrubs/vines S6–101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 0-101 0-101 vine jointfir <t< td=""><td></td><td>collegeflower</td><td>HYFL</td><td>Hymenopappus flavescens</td><td>0–112</td><td>-</td></t<> | | collegeflower | HYFL | Hymenopappus flavescens | 0–112 | - |
| Gordon's bladderpod LEGO Lesquerella gordonii 0-112 dotted blazing star LIPU Liatris punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MELE2 Melampodium leucanthum 0-112 blazingstar MENTZ Mentzelia 0-112 Nuttall's sensitive-briar MINU6 Mirosa nuttallii 0-112 evening primrose OENOT Oenothera 0-112 yellow naliwort PAVI4 Paronychia virginica 0-112 yellow naliwort PAVI4 Paronychia virginica 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuifforum 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Shrubs/vines S66-101 112 112 Shrubs/vines CYLE8 Cylindropuntia leptocaulis 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 vine | | trailing krameria | KRLA | Krameria lanceolata | 0–112 | - |
| dotted blazing star LIPU Liatris punctata 0-112 hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MEL22 Melampodium leucanthum 0-112 blazingstar MENTZ Mentzelia 0-112 Nuttall's sensitive-briar MINU6 Mimosa nuttallii 0-112 evening primose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 earleaf fanpetals SITR Sida tragifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Shrubs/vines S6-101 S6-101 Shrubs/vine 0-101 f faatherplume DAFO Dalea formosa 0-101 | | Gordon's bladderpod | LEGO | Lesquerella gordonii | 0–112 | - |
| hoary blackfoot MECI Melampodium cinereum 0-112 plains blackfoot MEL2 Melampodium leucanthum 0-112 blazingstar MENTZ Mentzelia 0-112 Nuttall's sensitive-briar MINU6 Mimosa nuttallii 0-112 evening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAV14 Paronychia virginica 0-112 Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 earleaf fanpetals SITR Sida tragitolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Shrub/Vine 6 shrubs/vines 56-101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 i featherplume DAFO Dalea formosa 0-101 vine jointfir E | | dotted blazing star | LIPU | Liatris punctata | 0–112 | _ |
| plains blackfoot MELE2 Melampodium leucanthum 0-112 blazingstar MENTZ Mentzelia 0-112 Nuttall's sensitive-briar MINU6 Mimosa nuttallii 0-112 evening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuifforum 0-112 earleaf fanpetals SITR Sida tragifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiffora 0-112 ShrubN/ine 56-101 1 1 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 if eatherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberr | | hoary blackfoot | MECI | Melampodium cinereum | 0–112 | _ |
| blazingstar MENTZ Mentzelia 0-112 Nuttall's sensitive-briar MINU6 Mimosa nuttallii 0-112 evening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragiifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiffora 0-112 Shrubs/vines 56-101 56-101 56-101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 if eatherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 algerita< | | plains blackfoot | MELE2 | Melampodium leucanthum | 0–112 | _ |
| Nuttall's sensitive-briar MINU6 Mimosa nuttallii 0-112 evening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragiifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vines 56-101 101 101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 featherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberry FOPU2 Forestiera pubescens 0-101 algerit | | blazingstar | MENTZ | Mentzelia | 0–112 | _ |
| evening primrose OENOT Oenothera 0-112 purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vine 56-101 C Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 featherplume DAFO Dalea formosa 0-101 10 10 vine jointfir EPPE Ephedra pedunculata 0-101 10 stretchberry FOPU2 Forestiera pubescens 0-101 algerita MATR3 Mahonia trifoliolata 0-101 gaurita | | Nuttall's sensitive-briar | MINU6 | Mimosa nuttallii | 0–112 | _ |
| purple locoweed OXLA3 Oxytropis lambertii 0-112 yellow nailwort PAVI4 Paronychia virginica 0-112 Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vine 56-101 112 112 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 featherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberry FOPU2 Forestiera pubescens 0-101 algerita MATR3 Mahonia trifoliolata 0-101 plains pricklypear OPPO Opuntia polyacantha 0-101 skunkbush sumac RHTR Rhus trilobata 0-101 soapweed | | evening primrose | OENOT | Oenothera | 0–112 | _ |
| yellow nailwort PAVI4 Paronychia virginica 0-112 Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 ShrubVine 56-101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 featherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberry FOPU2 Forestiera pubescens 0-101 algerita MATR3 Mahonia trifoliolata 0-101 plains pricklypear OPPO Opuntia polyacantha 0-101 skunkbush sumac RHTR Rhus trilobata 0-101 skunkbush sumac RHTR Rhus trilobata 0-101 Tree | | purple locoweed | OXLA3 | Oxytropis lambertii | 0–112 | - |
| Fendler's penstemon PEFE Penstemon fendleri 0-112 white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragiifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vine 56-101 56-101 6 shrubs/vines 56-101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 if eatherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberry FOPU2 Forestiera pubescens 0-101 algerita MATR3 Mahonia trifoliolata 0-101 plains pricklypear OPPO Opuntia polyacantha 0-101 skunkbush sumac RHTR Rhus trilobata 0-101 soapweed yucca YUGL Yucca glauca 0-101 Tree 28-112 0 0 <td></td> <td>yellow nailwort</td> <td>PAVI4</td> <td>Paronychia virginica</td> <td>0–112</td> <td>_</td> | | yellow nailwort | PAVI4 | Paronychia virginica | 0–112 | _ |
| white milkwort POAL4 Polygala alba 0-112 slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragiifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vine 56-101 0 6 shrubs/vines 56-101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0-101 featherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberry FOPU2 Forestiera pubescens 0-101 algerita MATR3 Mahonia trifoliolata 0-101 glains pricklypear OPPO Opuntia polyacantha 0-101 stunkbush sumac RHTR Rhus trilobata 0-101 soapweed yucca YUGL Yucca glauca 0-101 Tree 7 trees 28-112 | | Fendler's penstemon | PEFE | Penstemon fendleri | 0–112 | _ |
| slimflower scurfpea PSTE5 Psoralidium tenuiflorum 0-112 earleaf fanpetals SITR Sida tragiifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vine 6 shrubs/vines 56-101 6 shrubs/vines 0-101 0-101 featherplume DAFO Dalea formosa 0-101 vine jointfir EPPE Ephedra pedunculata 0-101 stretchberry FOPU2 Forestiera pubescens 0-101 algerita MATR3 Mahonia trifoliolata 0-101 plains pricklypear OPPO Opuntia polyacantha 0-101 skunkbush sumac RHTR Rhus trilobata 0-101 soapweed yucca YUGL Yucca glauca 0-101 Tree 7 trees 28-112 | | white milkwort | POAL4 | Polygala alba | 0–112 | - |
| earleaf fanpetals SITR Sida tragiifolia 0-112 stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0-112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0-112 Shrub/Vine 0 0 0 0 6 shrubs/vines 56-101 0 0 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0 0 0 featherplume DAFO Dalea formosa 0 | | slimflower scurfpea | PSTE5 | Psoralidium tenuiflorum | 0–112 | - |
| stemmy four-nerve daisy TESC2 Tetraneuris scaposa 0–112 Rocky Mountain zinnia ZIGR Zinnia grandiflora 0–112 Shrub/Vine 56–101 56–101 6 shrubs/vines 56–101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0–101 featherplume DAFO Dalea formosa 0–101 vine jointfir EPPE Ephedra pedunculata 0–101 stretchberry FOPU2 Forestiera pubescens 0–101 algerita MATR3 Mahonia trifoliolata 0–101 catclaw mimosa MIACB Mimosa aculeaticarpa var. biuncifera 0–101 plains pricklypear OPPO Opuntia polyacantha 0–101 soapweed yucca YUGL Yucca glauca 0–101 Tree 7 trees 28–112 | | earleaf fanpetals | SITR | Sida tragiifolia | 0–112 | _ |
| Rocky Mountain zinniaZIGRZinnia grandiflora0–112Shrub/Vine6shrubs/vines56–1016christmas cactusCYLE8Cylindropuntia leptocaulis0–1011featherplumeDAFODalea formosa0–1011vine jointfirEPPEEphedra pedunculata0–1011stretchberryFOPU2Forestiera pubescens0–1011algeritaMATR3Mahonia trifoliolata0–1011catclaw mimosaMIACBMimosa aculeaticarpa var. biuncifera0–1011skunkbush sumacRHTRRhus trilobata0–1011soapweed yuccaYUGLYucca glauca0–1017trees28–112 | | stemmy four-nerve daisy | TESC2 | Tetraneuris scaposa | 0–112 | - |
| Shrub/Vine 6 shrubs/vines 56–101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0–101 featherplume DAFO Dalea formosa 0–101 vine jointfir EPPE Ephedra pedunculata 0–101 stretchberry FOPU2 Forestiera pubescens 0–101 algerita MATR3 Mahonia trifoliolata 0–101 catclaw mimosa MIACB Mimosa aculeaticarpa var. biuncifera 0–101 plains pricklypear OPPO Opuntia polyacantha 0–101 skunkbush sumac RHTR Rhus trilobata 0–101 Tree 7 trees 28–112 | | Rocky Mountain zinnia | ZIGR | Zinnia grandiflora | 0–112 | _ |
| 6 shrubs/vines 56–101 Christmas cactus CYLE8 Cylindropuntia leptocaulis 0–101 featherplume DAFO Dalea formosa 0–101 vine jointfir EPPE Ephedra pedunculata 0–101 stretchberry FOPU2 Forestiera pubescens 0–101 algerita MATR3 Mahonia trifoliolata 0–101 catclaw mimosa MIACB Mimosa aculeaticarpa var. biuncifera 0–101 plains pricklypear OPPO Opuntia polyacantha 0–101 skunkbush sumac RHTR Rhus trilobata 0–101 Tree 7 trees 28–112 | Shrub | /Vine | • | • | | |
| Christmas cactusCYLE8Cylindropuntia leptocaulis0–101featherplumeDAFODalea formosa0–101vine jointfirEPPEEphedra pedunculata0–101stretchberryFOPU2Forestiera pubescens0–101algeritaMATR3Mahonia trifoliolata0–101catclaw mimosaMIACBMimosa aculeaticarpa var. biuncifera0–101plains pricklypearOPPOOpuntia polyacantha0–101skunkbush sumacRHTRRhus trilobata0–101soapweed yuccaYUGLYucca glauca0–101Tree7trees28–112 | 6 | shrubs/vines | | | 56–101 | |
| featherplumeDAFODalea formosa0-101vine jointfirEPPEEphedra pedunculata0-101stretchberryFOPU2Forestiera pubescens0-101algeritaMATR3Mahonia trifoliolata0-101catclaw mimosaMIACBMimosa aculeaticarpa var. biuncifera0-101plains pricklypearOPPOOpuntia polyacantha0-101skunkbush sumacRHTRRhus trilobata0-101soapweed yuccaYUGLYucca glauca0-101Tree28-1120 | | Christmas cactus | CYLE8 | Cylindropuntia leptocaulis | 0–101 | _ |
| vine jointfirEPPEEphedra pedunculata0–101stretchberryFOPU2Forestiera pubescens0–101algeritaMATR3Mahonia trifoliolata0–101catclaw mimosaMIACBMimosa aculeaticarpa var. biuncifera0–101plains pricklypearOPPOOpuntia polyacantha0–101skunkbush sumacRHTRRhus trilobata0–101soapweed yuccaYUGLYucca glauca0–101Tree28–11228–112 | | featherplume | DAFO | Dalea formosa | 0–101 | _ |
| stretchberry FOPU2 Forestiera pubescens 0–101 algerita MATR3 Mahonia trifoliolata 0–101 catclaw mimosa MIACB Mimosa aculeaticarpa var. biuncifera 0–101 plains pricklypear OPPO Opuntia polyacantha 0–101 skunkbush sumac RHTR Rhus trilobata 0–101 soapweed yucca YUGL Yucca glauca 0–101 Tree 28–112 28–112 | | vine jointfir | EPPE | Ephedra pedunculata | 0–101 | _ |
| algerita MATR3 Mahonia trifoliolata 0–101 catclaw mimosa MIACB Mimosa aculeaticarpa var. biuncifera 0–101 plains pricklypear OPPO Opuntia polyacantha 0–101 skunkbush sumac RHTR Rhus trilobata 0–101 soapweed yucca YUGL Yucca glauca 0–101 Tree 28–112 28–112 | | stretchberry | FOPU2 | Forestiera pubescens | 0–101 | _ |
| catclaw mimosa MIACB Mimosa aculeaticarpa var. biuncifera 0-101 plains pricklypear OPPO Opuntia polyacantha 0-101 skunkbush sumac RHTR Rhus trilobata 0-101 soapweed yucca YUGL Yucca glauca 0-101 Tree 28-112 | | algerita | MATR3 | Mahonia trifoliolata | 0–101 | _ |
| plains pricklypear OPPO Opuntia polyacantha 0–101 skunkbush sumac RHTR Rhus trilobata 0–101 soapweed yucca YUGL Yucca glauca 0–101 Tree 28–112 28–112 | | catclaw mimosa | MIACB | Mimosa aculeaticarpa var. biuncifera | 0–101 | _ |
| skunkbush sumac RHTR Rhus trilobata 0–101 soapweed yucca YUGL Yucca glauca 0–101 Tree 28–112 | | plains pricklypear | OPPO | Opuntia polyacantha | 0–101 | _ |
| soapweed yucca YUGL Yucca glauca 0–101 Tree 7 trees 28–112 | | skunkbush sumac | RHTR | Rhus trilobata | 0–101 | _ |
| Tree 28–112 | | soapweed yucca | YUGL | Yucca glauca | 0–101 | _ |
| 7 trees 28–112 | Tree | | | | | |
| | 7 | trees | | | 28–112 | |
| netleaf hackberry CELAR Celtis laevigata var. reticulata 0–112 | | netleaf hackberry | CELAR | Celtis laevigata var. reticulata | 0–112 | _ |
| Pinchot's juniper JUPI Juniperus pinchotii 0–112 | | Pinchot's juniper | JUPI | Juniperus pinchotii | 0–112 | _ |
| honey mesquite PRGL2 Prosopis glandulosa 0–112 | | honey mesquite | PRGL2 | Prosopis glandulosa | 0–112 | |

Animal community

White tail deer and mule deer utilize the site for browsing. The site has little cover for bobwhite quail but scaled quail frequent the site during feeding. The Texas horned lizard is often present along with several species of lizards and snakes. A few species of small mammals often find dens in rock crevices.

Hydrological functions

Runoff water from the site will enter streams lower on the landscape. The site yields considerable runoff due to steep slopes.

Recreational uses

Hunting, Camping, Hiking, Birdwatching, Photography, and Horseback Riding.

Wood products

None.

Other products

None.

Other information

None.

Inventory data references

NRCS FOTG – Section II of the FOTG Range Site Descriptions and numerous historical accounts of vegetative conditions at the time of early settlement in the area were used in the development of this site description. Vegetative inventories were made at several site locations for support documentation. Several years of clipping data were surveyed.

Inventory Data References (documents): NRCS FOTG – Section II - Range Site Descriptions and NRCS Clipping Data summaries over a 20 year period

Other references

Natural Resources Conservation Service - Range Site Descriptions USDA-Natural Resources Conservation Service - Soil Surveys & Website soil database Rathjen, Frederick W., The Texas Panhandle Frontier, Rev. 1998, Univ. of Texas Press Hatch, Brown and Ghandi, Vascular Plants of Texas (An Ecological Checklist) Texas A&M Exp. Station, College Station, Texas Texas Tech University – Range, Wildlife & Fisheries Dept.

Technical Reviewers: Tony Garcia, Zone RMS, NRCS, Lubbock, Texas Clint Rollins, RMS, NRCS, Amarillo, Texas Mark Moseley, Acting State RMS, NRCS, San Antonio, Texas Jack Eckroat, Oklahoma Ecologist, NRCS, Stillwater, Oklahoma Justin Clary, RMS, NRCS, Temple, Texas

Contributors

James Crownover PES Edits by Tyson Morley, MLRA Soil Scientist, Altus, Oklahoma

Approval

Acknowledgments

Site Development and Testing Plan:

Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document.

Annual reviews of the Project Plan are to be conducted by the Ecological Site Technical Team.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| Author(s)/participant(s) | Stan Bradbury, Zone RMS, NRCS, Lubbock, Texas |
|---|---|
| Contact for lead author | 806-791-0581 |
| Date | 09/04/2007 |
| Approved by | Bryan Christensen |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

- 1. Number and extent of rills: Due to the percent slopes, rills will be common.
- 2. Presence of water flow patterns: Due to the percent slopes, water flow patterns will be common.
- 3. Number and height of erosional pedestals or terracettes: Due to the percent slopes, pedestals/terracettes will be common.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 25-30% mineral soil; low percentage due to rock fragments scattered throughout the profile.
- 5. Number of gullies and erosion associated with gullies: Slight to moderate.

6. Extent of wind scoured, blowouts and/or depositional areas: None to slight.

- 7. Amount of litter movement (describe size and distance expected to travel): Slight to moderate.
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): If the soil surface is unprotected by vegetation, the sloping soil is highly susceptible to water erosion.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Calcareous loam to fine sandy loam; moderate SOM.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Low vegetative cover and percent slopes makes this site susceptible to erosion. This site is a slowly permeable soil, well-draiend and available water holding capacity is low. Plant growth and production is limited due to very shallow depth.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Warm-season tallgrasses >

Sub-dominant: Warm-season midgrasses > Warm-season shortgrasses >

Other: Forbs > Shrubs/Vines > Trees

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Plant mortality and decadence is minimal.

14. Average percent litter cover (%) and depth (in): Litter is dominantly herbaceous.

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): 650 - 1,300 pounds per acre.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that

become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Juniper and mesquite can be invasive.

17. **Perennial plant reproductive capability:** All plant species should be capable of reproduction, except during periods of prolonged drought conditions, heavy natural herbivory or intense wildfires.