

Ecological site R081CY355TX Adobe 29-35 PZ

Last updated: 9/19/2023
Accessed: 05/23/2025

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	Zone Rangeland Management Specialist, NRCS, San Angelo, Texas, 325-944-0147
Date	06/29/2005
Approved by	Colin Walden
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

2. **Presence of water flow patterns:** None, except following extremely high intensity storms where short flow patterns may appear.

3. **Number and height of erosional pedestals or terracettes:** None

-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Expect no more than 10% bare ground randomly distributed throughout in small and non-connected areas.
-
5. **Number of gullies and erosion associated with gullies:** None.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Under normal rainfall, little litter movement should be expected; however, litter of all sizes may move long distances. Minimal and short.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface is resistant to erosion. Stability class range is expected to be 5-6
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is light brownish gray gravelly clay loam with limestone moderately fine subangular blocky and moderately fine granular structure on the surface. SOM is approximately 1-3%. See soil survey for specific soils.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The savannah of tallgrasses, midgrasses, forbs and trees having adequate litter and little bare ground can provide for maximum infiltration and little runoff under normal rainfall events.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No evidence of compaction.

-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Warm-season midgrasses

Sub-dominant: Forbs Trees

Other: Warm-season tallgrasses Warm-season shortgrasses Shrubs

Additional: Forbs make up 10 percent species composition, shrubs has about 5 percent species composition and trees have 15 percent annual production.n

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** There should be little mortality or decadence for any functional groups.
-

14. **Average percent litter cover (%) and depth (in):** Litter is primarily herbaceous
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1800 to 4100 pounds per acre
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Ashe Juniper is dominant, Honey mesquite, Prickly pear, Bermudagrass, Johnsongrass, King Ranch bluestem
-

17. **Perennial plant reproductive capability:** All perennial plants should be capable of

reproducing, except during periods of prolonged drought conditions, heavy natural herbivory, and wildfires.
