

Ecological site R081CY359TX Gravelly Redland 29-35 PZ

Last updated: 9/19/2023
Accessed: 04/25/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 081C—Edwards Plateau, Eastern Part

This area represents the eastern part of the Edwards Plateau region. Limestone ridges and canyons and nearly level to gently sloping valley floors characterize the area. Elevation is 400 feet (120 meters) at the eastern end of the area and increases westward to 2,400 feet (730 meters) on ridges. This area is underlain primarily by limestones in the Glen Rose, Fort Terrett, and Edwards Formations of Cretaceous age. Quaternary alluvium is in river valleys.

Classification relationships

Major Land Resource Area (MLRA) and Land Resource Unit (LRU) (USDA-Natural Resources Conservation Service, 2006)

National Vegetation Classification/Shrubland & Grassland/2C Temperate & Boreal Shrubland and Grassland/M051 Great Plains Mixedgrass Prairie & Shrubland/ G133 Central Great Plains Mixedgrass Prairie Group.

Ecological site concept

These upland sites occur on gravelly clay loam soils over limestone. Gravel content ranges up to 25 percent at the surface to 80 percent in the subsoil. Reference vegetation includes mid and tallgrasses with numerous forbs and

scattered oaks. Without fire or other brush management, woody species are likely to increase on the site.

Associated sites

R081CY356TX	Blackland 29-35 PZ The Blackland ecological site has higher production and deeper soils.
R081CY355TX	Adobe 29-35 PZ The Adobe ecological site has less production, more slope, and more caliche type soils of a higher pH with no post oak or blackjack oak.
R081CY360TX	Low Stony Hill 29-35 PZ The Low Stony Hill ecological site is generally higher in the landscape and is the plateau above the Gravelly Redland ecological site with no post oak or blackjack oak.

Similar sites

R081CY361TX	Redland 29-35 PZ The Redland ecological site has deeper soils, less permeable and is more productive. There is typically post oak, blackjack oak, and live oak on the Redland ecological site as well as taller grasses.
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Quercus fusiformis</i> (2) <i>Acacia greggii</i>
Shrub	Not specified
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Bouteloua curtipendula</i>

Physiographic features

This site is located in the 81C, Eastern Edwards Plateau Major Land Resource Area (MLRA). Gravelly Redland is found on gently to moderately sloping plains and uplands. Slopes range from 1 to 8 percent. The site consists of moderately deep, well-drained, moderately slowly permeable soils formed in residuum and colluviums over indurated limestone bedrock. Due to the slopes, this site is susceptible to erosion. The elevation ranges from 800 feet to 2300 feet at this site. No flooding or ponding is expected. Low to very high run-off occurs on the Gravelly Redland.




Figure 2. Gravelly Redland 081CY359TX

Table 2. Representative physiographic features

Landforms	(1) Plateau > Ridge (2) Plateau > Plain
-----------	--

Runoff class	Low to very high
Flooding frequency	None
Ponding frequency	None
Elevation	800–2,300 ft
Slope	1–8%
Aspect	Aspect is not a significant factor

Climatic features

The climate is humid subtropical and is characterized by hot summers and relatively mild winters. The average first frost should occur around November 15 and the last freeze of the season should occur around March 19.

The average relative humidity in mid-afternoon is about 50 percent. Humidity is higher at night, and the average at dawn is about 80 percent. The sun shines 70 percent of the time possible during the summer and 50 percent in winter. The prevailing wind direction is southeast.

Drought is calculated as 75% below average rainfall. It should be noted that timing of rainfall may be more significant than average rainfall.

Approximately two-thirds of annual rainfall occurs during the April to September period. Rainfall during this period generally falls during thunderstorms, and fairly large amount of rain may fall in a short time. Hurricanes provide another source of extremely high rains in a short time. A review of the rainfall records suggest that rainfall is below “normal” at least 60 percent of the time. Therefore, the erratic nature of the rainfall should be considered when developing any land management plans.

The impact of droughts in the Edwards Plateau cannot be under-estimated. Not only are droughts devastating to the land but also to those that manage the land. Droughts occur roughly every 20 years but not always. A severe drought in 2012 coupled with extreme heat resulted in a die off of juniper over millions of acres as well as other native plants.

Table 3. Representative climatic features

Frost-free period (characteristic range)	220-260 days
Freeze-free period (characteristic range)	227-269 days
Precipitation total (characteristic range)	32-37 in
Frost-free period (actual range)	187-260 days
Freeze-free period (actual range)	224-332 days
Precipitation total (actual range)	31-37 in
Frost-free period (average)	240 days
Freeze-free period (average)	257 days
Precipitation total (average)	34 in

Climate stations used

- (1) MEDINA 1NE [USC00415742], Medina, TX
- (2) SAN ANTONIO/SEAWORLD [USC00418169], San Antonio, TX
- (3) KERRVILLE 3 NNE [USC00414782], Kerrville, TX
- (4) BLANCO [USC00410832], Blanco, TX
- (5) CANYON DAM [USC00411429], Canyon Lake, TX
- (6) BURNET MUNI AP [USW00003999], Burnet, TX
- (7) AUSTIN GREAT HILLS [USC00410433], Austin, TX

- (8) GEORGETOWN LAKE [USC00413507], Georgetown, TX
- (9) PRADE RCH [USC00417232], Leahey, TX

Influencing water features

This being an upland site, it is not influenced by water from a wetland or stream.

Wetland description

N/A

Figure 7-1 The hydrologic cycle with factors that affect hydrologic processes




Figure 9.

Soil features

In a representative profile for the Gravelly Redland ecological site, the surface layer is dark reddish-brown very gravelly clay loam about 5 inches thick. They are underlain by slightly fractured indurated limestone bedrock at depths of 20 to 40 inches. Rock fragments in the surface horizons are about 25 percent and up to 80 percent in the subsoil. There are gravelly to extremely gravelly texture modifiers throughout the soil profile. Because of the moderate slopes associated with Gravelly Redland, the site is well drained and has moderate runoff. The high gravel content in the subsurface horizons reduces the shrink-swell potential which causes the soil permeability to be moderately slow. When plant residues are inadequate, soil condition deteriorates and heavy surface crusts develop. In this condition water intake is very slow, runoff is rapid, erosion is a hazard, and grass recovery is slow. The mineral content and reaction of these soils enable the site to produce highly nutritious forage. In association with other sites, Gravelly Redland is usually the preferred grazing area. These sites occur on stable hillslopes on dissected plateaus.

Due to the scale of mapping, there are inclusions of minor components of other soils within these mapping units. Before performing any inventories, conduct a field evaluation to ensure the soils are correct for the site.

The representative soil series associated with the Gravelly Redland ecological site are Dina and Rumble.

Table 4. Representative soil features

Parent material	(1) Residuum–limestone
Surface texture	(1) Very gravelly clay loam (2) Cobbly silty clay loam
Drainage class	Well drained
Permeability class	Moderately slow to slow
Depth to restrictive layer	20–40 in
Soil depth	20–40 in

Surface fragment cover <=3"	5–20%
Surface fragment cover >3"	0–5%
Available water capacity (0-40in)	0.6–5.5 in
Calcium carbonate equivalent (0-40in)	0–10%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0
Soil reaction (1:1 water) (0-40in)	5.6–8.4
Subsurface fragment volume <=3" (4-40in)	30–60%
Subsurface fragment volume >3" (4-40in)	5–20%

Ecological dynamics

The information contained in the State and Transition Diagram (STD) and the Ecological Site Description was developed using archeological and historical data, professional experience, and scientific studies. The information presented is representative of a very complex set of plant communities. Not all scenarios or plants are included. Key indicator plants, animals and ecological processes are described to inform land management decisions.

The pre-settlement or reference plant community is perceived as a prairie with mostly mid-grasses, associated forbs and scattered mottes of woody species; the Mixed-grass Prairie Community (1.1). The dominant grasses were little bluestem (*Schizachyrium scoparium*), sideoats grama (*Bouteloua curtipendula*), and feathery bluestems (*Bothriochloa* spp.). Big bluestem (*Andropogon gerardii*) and Indiangrass (*Sorghastrum nutans*) were scattered in small amounts in the wetter areas and inclusions of low stony hill soils. Velvet bundleflower (*Desmanthus velutinus*), Engelmann's daisy (*Engelmannia peristeria*), gaura (*Gaura* spp.), western indigo (*Indigofera miniata* var. *leptosepala*), and blazing star (*Liatris* spp.) were some of many forbs found in the historic community. Shrubs included elbowbush (*Forestiera pubescens*) and catclaw acacia (*Acacia greggii*). The endemic woody plants historically shaded less than 10 percent of the soil surface, either as small mottes or individual trees that were either resistant to fire or occupied areas where fires were less frequent or intense. Texas live oak (*Quercus fusiformis*) was the most frequently occurring tree species.

Natural plant mortality is very low with the major species producing seeds and vegetative structure each year in normal years. Litter cover is 100 percent. Physical soil crust is largely absent.

A study of early photographs of this region reveals that today these sites are much denser with woody cover and less covered with grasslike vegetation. Early accounts consistently describe this region as a vast expanse of hills covered with "cedar" from San Antonio to Austin. Accounts also describe an abundance of clean, flowing water and abundant wildlife. These accounts seem to describe heavy wooded areas in mosaic patterns occurring along the highs and lows of the landscape.

The plant communities of this site are dynamic and vary in relation to grazing, fire, and rainfall. Studies of the pre-European vegetation of the general area suggested 47 percent of the area was wooded (Wills, 2006). Historical records are not specific on the Gravelly Redland site but do reflect area observations. From the Teran expedition in 1691, "great quantities of buffaloes" were noted in the area. By 1840 the Bonnell expedition reflected that "buffalo rarely range so far to the south" (Inglis, 1964). Another example is an early settler, Arnold Gugger, who wrote in his journal about the mid to late 1800s in the Helotes, Texas area, "in those days buffaloes were in droves by the hundreds.....and antelopes were three to four hundred in a bunch....and deer and turkeys at any amount" (Massey, 2009).

Many research studies document the interaction of bison grazing and fire (Fuhlendorf, et al., 2008.). Bison would

come into an area, graze it down, leave and then not come back for many months or even years. Many times this grazing scheme by buffalo was high impact and followed fire patterns and available natural water. This usually long deferment period allowed the taller grasses and forbs to recover from the high impact bison grazing. This relationship created a diverse landscape both in structure and composition.

Historical fire frequencies for the region are suggested to be 13 to 25 years (Frost, 1998). When fires did occur, they were set either by Native Americans or by lighting. Woody plant control would vary in accordance with the intensity and severity of the fire encountered, which resulted in a mosaic of vegetation types within the same site.

Ashe juniper (*Juniperus ashei*) will increase regardless of grazing. Juniper will establish with grazing and without unless goats are utilized. Goats and probably sheep will eat young juniper and when properly used, are an effective tool to maintain juniper (Taylor, 1997; Anderson, et al., 2013). The main role of excessive grazing relative to juniper is the removal of the fine fuel needed to carry an effective burn.

Ashe juniper, because of its dense low growing foliage, has the ability to retard grass and forb growth. Grass and forb growth can become non-existent under dense juniper canopies. Many times there is a resurgence of the better grasses such as little bluestem when Ashe juniper is controlled and followed by proper grazing management. Seeds and dormant rootstocks of many plant species are contained in the leaf mulch and duff under the junipers.

Currently, cattle, goats, white-tailed deer, sheep, and exotic animals are the primary large herbivores. At settlement, large numbers of deer occurred, but as human populations increased (with unregulated harvest) their numbers declined substantially. Eventually, laws and restrictions on deer harvest were put in place which assisted in the recovery of the species. Females were not harvested for several decades following the implementation of hunting laws, which allowed population booms. In addition, suppression of fire favored woody plants which provided additional browse and cover for the deer. Because of their impacts on livestock production, large predators such as red wolves (*Canis rufus*), mountain lions (*Felis concolor*), black bears (*Ursus americanus*), and eventually coyotes (*Canis latrans*) were reduced in numbers or eliminated (Schmidly, 2002).

The screwworm fly (*Cochilomyia hominivorax*) was essentially eradicated by the mid-1960s, and while this was immensely helpful to the livestock industry, this removed a significant control on deer populations (Teer, Thomas, and Walker, 1965; Bushland, 1985).

Progressive management of the deer herd, because of their economic importance through lease hunting, has the objective of improving individual deer quality and improving habitat. Managed harvest based on numbers, sex ratios, condition, and monitoring of habitat quality has been effective on individual properties. However, across the Edwards Plateau, excess numbers still exist which may lead to habitat degradation and significant die-offs during stress periods such as extended droughts.

The Edwards Plateau is home to a variety of exotic ungulates, mostly introduced for hunting (Schmidly, 2002). These animals are important sources of income to some landowners, but as with the white-tailed deer, their populations must be managed to prevent degradation of the habitat for themselves as well as for the diversity of native wildlife in the area. Many other species of medium- and small-sized mammals, birds, and insects can have significant influences on the plant communities in terms of pollination, herbivory, seed dispersal, and creation of local disturbance patches, all of which contribute to the plant species diversity.

The plants and topography aided in increasing the infiltration of rainfall into the moderately slowly permeable soil. Any loss of soil organic matter and plant cover has a negative effect on infiltration. More rainfall is directed to overland flow, which causes increased soil erosion and flooding. Soils are also more prone to drought stress since organic matter acts like a sponge aiding in moisture retention for plant growth. Mulch buildup under the Ashe juniper canopy, following brush management and incorporation into the soil, can have a positive effect on increasing infiltration.

This site contains a large diversity of plants and this document does not attempt to cover them all. The intent of this document is to describe ecological processes on representative plants.

European settlement occurred in the mid to late 1800s (Raunick, 2007). This time period also coincided with a stoppage of fire. It was during this time that large-scale fencing was initiated to help the introduction of livestock. Predators were also reduced to protect livestock. In many cases sheep and goats heavily utilized the site. Low

successional, unpalatable grasses, forbs, and shrubs have taken the place of the more desirable plant species. Non-preferred browse, such as juniper, fared well at the expense of the palatable browse. Juniper is undoubtedly the dominant woody plant over most of the site today.

Plant Communities and Transitional Pathways (diagram)

A State and Transition Diagram for the Gravelly Redland Ecological Site (R081CY359TX) is depicted in Figure 1. Descriptions of each state, transition, plant community, and pathway follow the model. Experts base this diagram on available experimental research, field observations, professional consensus, and interpretations. It is likely to change as knowledge increases.


Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The Reference Plant Community is not necessarily the management goal. Other vegetative states may be desired plant communities as long as the Range Health assessments are in the moderate and above category. The biological processes on this site are complex. Therefore, representative values are presented in a land management context. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

Both percent species composition by weight and percent canopy cover are described as are other metrics. Most observers find it easier to visualize or estimate percent canopy for woody species (trees and shrubs). Canopy cover can drive the transitions between communities and states because of the influence of shade and interception of rainfall. Species composition by dry weight is used for describing the herbaceous community and the community as a whole. Woody species are included in species composition for the site. Calculating similarity index requires the use of species composition by dry weight.

The following diagram suggests some pathways that the vegetation on this site might take. There may be other states not shown in the diagram. This information is intended to show what might happen in a given set of circumstances. It does not mean that this would happen the same way in every instance. Local professional guidance should always be sought before pursuing a treatment scenario.


State and transition model

Ecosystem states




- T1A** - Absence of disturbance and natural regeneration over time
- R2A** - Reintroduction of natural disturbance regimes
- T2A** - Ground disturbing brush management coupled with seeding non-native grasses


State 1 submodel, plant communities



State 2 submodel, plant communities



State 3 submodel, plant communities



State 1 Reference

The reference state is considered to be representative of the range of variability under pre-Euro settlement conditions. It is characterized by an open stand of mid and tallgrasses with scattered trees and shrubs. Woody canopy ranges from 10 to 20%. Community phase changes are primarily driven by grazing, periodic fire and variations in the subtropical climate. Wildfires set either by Native peoples or lightning, occurred at 7 to 12 year intervals (Frost, 1998).

Dominant plant species

- Texas live oak (*Quercus fusiformis*), tree
- little bluestem (*Schizachyrium scoparium*), other herbaceous
- sideoats grama (*Bouteloua curtipendula*), other herbaceous

Community 1.1 Mixed Grass Prairie Community

This is the diagnostic or reference plant community for the Gravelly Redland site. The metrics of the description of the reference plant community is derived from field observations, limited clipping data and professional consensus. The plant community is perceived as a prairie composed of mostly midgrasses with scattered tallgrasses, trees and shrubs. Vegetative composition by weight is estimated to be 80-87% grasses, 6-8% forbs, 4-6% shrubs, and 2-5% trees. Woody canopy cover is <10 percent. Woody plants, consisting of scattered trees or shrub mottes, shaded less than 10 percent of the site. Live oak, hackberry (*Celtis* spp.) and shrubs such as catclaw acacia, sumac (*Rhus* spp.), elbowbush (*Forestiera pubescens*), ephedra (*Ephedra* spp.), algerita, bumelia (*Sideroxylon lanuginosum*), and pricklypear (*Opuntia* spp.) were likely present but were kept suppressed by periodic fires and competition from the grasses. Above ground plant production ranged from 2,000 to 4,500 pounds per year. The grassland component accounted for 85 to 90 percent of the sites primary production, with little bluestem and sideoats grama the most abundant and productive species. Big bluestem and Indiangrass were confined to draws and wetter areas. Secondary midgrasses were Arizona cottontop (*Digitaria californica*), silver bluestem (*Bothriochloa laguroides* var. *torreyana*), green sprangletop (*Leptochloa dubia*), Texas cupgrass (*Eriochloa sericea*), and Texas wintergrass (*Nassella leucotricha*). Shortgrasses, like buffalograss and curlymesquite (*Hilaria belangeri*), were present in small amounts. Mexican sagewort (*Artemisia ludoviciana*), catclaw sensitivebriar (*Mimosa nuttallii*), bundleflower, western indigo, and orange zexmenia (*Zexmenia* spp.) were a few of the small (5 to 7 percent) but important forb

components of the plant community (See Plant Community Composition and Annual Production table below). Soil erosion, particularly on the upland plain areas, was very low because of the abundant plant cover, litter, good soil structure, and abundant chert outcrops. Runoff from the reference plant community was reduced because of grass cover and fissures in the limestone parent material. The vegetative ground cover helped disperse and slow down runoff, thus holding soil in place and enhancing infiltration. Concentrated water flow patterns were rare in this plant community. Without proper management that adjusts animal numbers based on annual forage production along with judicious prescribed burning and brush management, the Mixed-Grass Prairie Community will transition to: The combined effect of continued overgrazing (including deer), drought, no brush management, and the accompanying decrease in frequency and intensity of fires causes the reference plant community to shift toward the Midgrass Oak Savannah Plant Community (1.2).

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1700	2550	3825
Shrub/Vine	100	150	225
Tree	100	150	225
Forb	100	150	225
Total	2000	3000	4500

Table 6. Soil surface cover

Tree basal cover	1-3%
Shrub/vine/liana basal cover	1-3%
Grass/grasslike basal cover	10-15%
Forb basal cover	1-3%
Non-vascular plants	0%
Biological crusts	0%
Litter	80-100%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	5-20%
Bedrock	0%
Water	0%
Bare ground	0-5%

Table 7. Woody ground cover

Downed wood, fine-small (<0.40" diameter; 1-hour fuels)	—
Downed wood, fine-medium (0.40-0.99" diameter; 10-hour fuels)	—
Downed wood, fine-large (1.00-2.99" diameter; 100-hour fuels)	—
Downed wood, coarse-small (3.00-8.99" diameter; 1,000-hour fuels)	—
Downed wood, coarse-large (>9.00" diameter; 10,000-hour fuels)	—
Tree snags** (hard***)	—
Tree snags** (soft***)	—
Tree snag count** (hard***)	0-2 per acre
Tree snag count** (hard***)	

* **Decomposition Classes:** N - no or little integration with the soil surface; I - partial to nearly full integration with the soil surface.

** >10.16cm diameter at 1.3716m above ground and >1.8288m height--if less diameter OR height use applicable down wood type; for pinyon and juniper, use 0.3048m above ground.

*** Hard - tree is dead with most or all of bark intact; Soft - most of bark has sloughed off.

Table 8. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	1-3%	0-1%
>0.5 <= 1	—	1-3%	3-5%	1-3%
>1 <= 2	—	3-5%	10-15%	3-10%
>2 <= 4.5	—	3-5%	30-50%	—
>4.5 <= 13	—	—	—	—
>13 <= 40	5-10%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Figure 11. Plant community growth curve (percent production by month). TX3779, Mid and Tallgrass Prairie Community. Warm-season rangeland with most production April to October.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2	2	2	10	20	24	10	5	10	10	3	2

Community 1.2

Mid Grass Oak Community



Figure 12. Midgrass Oak Savannah Community, Gravelly Redland ecological site, Hays County, Texas.

The Midgrass Oak Savannah Community (1.2) is a midgrass dominated grassland being encroached by indigenous or invading woody species that had been held at low densities by repeated fires and competition from a vigorous grass component. Numerous woody species, including juniper and mesquite, are increasing in density because overgrazing by livestock has reduced grass cover, exposed some soil, and reduced fine fuel for fire. Less rainfall soaks into the soil except under the brush species where stem flow directs the rainfall to the soil beneath the brush. The woody canopy varies between 10 and 25 percent depending on time since last burned or the brush was treated and availability of invading species. Typically, oaks increase in size and mesquite and/or juniper increase in density. Less palatable brushy species such as catclaw acacia, bumelia, Texas persimmon (*Diospyros texana*), sumacs, condalia, elbowbush, and feather dalea (*Dalea* spp.) also increase. The preferred tall grasses are being replaced by the more grazing resistant midgrasses. Characteristic grasses are little bluestem, sideoats grama, tall (*Sporobolus compositus* var. *compositus*) and meadow dropseed (*Sporobolus asper* var. *asper*), vine mesquite (*Panicum obtusum*), plains lovegrass (*Eragrostis intermedia*), Texas cupgrass (*Eriochloa sericea*), and feathery bluestems (*Bothriochloa* spp.). Most of the perennial forbs found in the historic community remain in this plant community. In this phase, the increasing woody species are generally less than five feet tall and are subject to control by improved

grazing management, prescribed burning and individual plant treatments (IPT). Annual primary production still ranges from 2,000 to 4,500 pounds per acre but a greater percentage is now woody species. Forage production is still predominantly grass species although somewhat suppressed. Heavy continuous grazing will reduce plant cover, litter, and mulch and increase bare ground exposing the soil to water erosion. Because of gentle slopes and grass cover, some soil movement could take place in this vegetation type during rainstorms. The changes in species composition are small initially, but unless measures are taken to restore the health and vigor of the palatable plants and do some form of suppression, woody species will continue to increase in size and density. As the woody cover increases they capture more sunlight, soil moisture and nutrients at the expense of more desired plants. The midgrasses give way to curlymesquite, buffalograss, and Texas wintergrass. In the Midgrass Oak Savannah Community (1.2), ecological processes have changed somewhat because of the shift in energy, moisture, and nutrient capture. The pathway back to the reference plant community can be accomplished without costly acceleration of conservation practices. However, grazing management alone will not reverse retrogression; some form of woody plant control, such as prescribed burning or individual plant treatment (IPT), must be used. Some, but not all woody species can be managed with targeted grazing with sheep and goats. Mesquite is one that is not a preferred browsing species. When the canopy of the woody plants becomes dense enough (25 percent) and tall enough (greater than 5 feet) to suppress grass growth and resist damage from fire, a threshold in ecological succession is crossed. The Midgrass Oak Savannah Community (1.2) transitions into the Shrubland State (2). Once this threshold has been passed, low cost range management practices such as proper grazing and prescribed burning cannot reverse the transition to woody plant dominance.

Table 9. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1300	1950	2925
Shrub/Vine	300	450	675
Tree	200	300	450
Forb	200	300	450
Total	2000	3000	4500

Table 10. Ground cover

Tree foliar cover	5-10%
Shrub/vine/liana foliar cover	3-10%
Grass/grasslike foliar cover	15-30%
Forb foliar cover	2-15%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 11. Soil surface cover

Tree basal cover	1-3%
Shrub/vine/liana basal cover	1-5%
Grass/grasslike basal cover	10-15%
Forb basal cover	1-3%
Non-vascular plants	0%

Biological crusts	0%
Litter	80-100%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	5-20%
Bedrock	0%
Water	0%
Bare ground	0-10%

Table 12. Woody ground cover

Downed wood, fine-small (<0.40" diameter; 1-hour fuels)	—
Downed wood, fine-medium (0.40-0.99" diameter; 10-hour fuels)	—
Downed wood, fine-large (1.00-2.99" diameter; 100-hour fuels)	—
Downed wood, coarse-small (3.00-8.99" diameter; 1,000-hour fuels)	—
Downed wood, coarse-large (>9.00" diameter; 10,000-hour fuels)	—
Tree snags** (hard***)	—
Tree snags** (soft***)	—
Tree snag count** (hard***)	0-2 per acre
Tree snag count** (hard***)	

* **Decomposition Classes:** N - no or little integration with the soil surface; I - partial to nearly full integration with the soil surface.

** >10.16cm diameter at 1.3716m above ground and >1.8288m height--if less diameter OR height use applicable down wood type; for pinyon and juniper, use 0.3048m above ground.

*** **Hard** - tree is dead with most or all of bark intact; **Soft** - most of bark has sloughed off.

Table 13. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	1-3%	0-1%
>0.5 <= 1	—	1-3%	3-5%	1-3%
>1 <= 2	—	3-5%	10-15%	3-15%
>2 <= 4.5	—	3-10%	20-50%	—
>4.5 <= 13	—	—	—	—
>13 <= 40	5-10%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Figure 14. Plant community growth curve (percent production by month).
TX3779, Mid and Tallgrass Prairie Community. Warm-season rangeland with most production April to October.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2	2	2	10	20	24	10	5	10	10	3	2

Pathway 1.1A

Community 1.1 to 1.2

A shift in the composition of the plant community is primarily driven by the lack of managing woody plants, juniper in particular. Juniper and other woody species are introduced from the site primarily through wildlife fecal deposits.

Grazing that removes fuel loading for fire is a contributing factor. However juniper can increase regardless of grazing pressure unless sheep and goats are utilized. Long term droughts will hasten the change of any of the plant communities.

Pathway 1.2A

Community 1.2 to 1.1

This recovery pathway consist of some method of brush management such as fire, mechanical or hand cutting or targeted grazing with goats and/or possibly sheep. Prescribed grazing is essential.

State 2

Shrubland State

This state is the culmination of a loss of sunlight energy capture by historic plants, interruption of the hydrologic cycle because of interception, and stem flow and competition for nutrients. Mesquite was the early understory brush species, but Ashe juniper has increased tremendously resulting in further decline in midgrasses. Increase in shortgrasses and annuals. Increase in mesquite, catclaw, and pricklypear have also increased.

Dominant plant species

- oak (*Quercus*), tree
- Ashe's juniper (*Juniperus ashei*), tree
- honey mesquite (*Prosopis glandulosa*), shrub

Community 2.1

Mixed Brush Shrubland Community



Figure 15. Mixed-Brush Shrubland Community of juniper with little herbaceous understory.

When woody plant canopy reaches 30 to 35 percent and grasses provide less than 50 percent of the herbage production, the transition from the Midgrass Oak Savannah Community (1.2) to the Mixed Brush Shrubland Community (2.1) is complete. At this point, there is generally not enough fine fuel produced by the grassland component to carry a prescribed fire of the intensity needed to control the woody plants. Once this threshold is reached proper grazing management and prescribed burning cannot reverse the transition. Intensive and usually expensive brush control practices must be applied to reverse the transition to the dense woodland community. Oak, mesquite, and/or juniper dominate this community phase. Juniper can achieve dominance in as little as 30 years from the onset of seedlings. The only remnants of grassland vegetation remain in the interspaces. Estimated vegetative composition by weight is 30-35% grasses, 8-10% forbs, 30-35% shrubs, and 25-30% trees. Woody canopy cover exceeds 40%. Common understory shrubs for this plant community are pricklypear, algerita, condalia, yucca (*Yucca* spp.), Texas persimmon, elbowbush, prickly ash (*Zanthoxylum* spp), and catclaw acacia. Shortgrasses, cool-season grasses, and low quality annual and perennial forbs occupy the tree interspaces. Characteristic grasses are Texas wintergrass, curlymesquite, buffalograss, and cedar sedge (*Carex planostachys*). Common forbs include dotted gayfeather (*Liatris punctata* var. *punctata*), orange zexmenia, croton (*Croton* spp.), western ragweed (*Ambrosia psilostachya*), prairie coneflower (*Ratibida columnifera*), and broomweed (*Gutierrezia*

spp.). With continued overgrazing, either by livestock or deer, the brush canopy increases in density and thickens while shortgrasses, such as threeawns (*Aristida* spp.), red grama (*Bouteloua trifida*), Texas grama (*Bouteloua rigidiseta*), hairy tridens (*Erioneuron pilosum*), and broomweed (*Gutierrezia* spp.) replace the more palatable mid and short grasses. Annual primary production varies from 2,500 to 4,500 pounds per acre; most of it in the woody component. Grasses and forbs make up 25 percent or less of the annual herbage production. The oak/mixed-brush overstory can reach 80 to 90 percent ground cover and produce 70 percent or more of the annual production. Although this vegetative state provides cover for wildlife, only limited preferred forage or browse is available for livestock or wildlife. The tree and shrub canopy acts to intercept rainfall and increase evapotranspiration losses, creating a more xeric microclimate. Soil fauna and litter are reduced exposing more soil surface to erosion in the few interspaces. However, within the woody canopy, hydrologic processes stabilize and soil organic matter and mulch begin to increase and eventually stabilize in the woodland state. Without major brush control and management inputs, this plant community cannot be reversed. It will continue to thicken until it stabilizes with the climate and soil. Returning the Oak/Mixed Brush Woodland (2.1) Community back to Midgrass Oak Community (1.2) requires extensive and expensive reclamation practices. Mechanical and/or chemical brush control must be followed by prescribed grazing and prescribed burning practices. If ground disturbance methods are used, replanting can be done to speed up the recovery and increase plant diversity.

Table 14. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Tree	800	1200	1800
Shrub/Vine	600	900	1350
Grass/Grasslike	500	750	1125
Forb	100	150	225
Total	2000	3000	4500

Table 15. Ground cover

Tree foliar cover	5-10%
Shrub/vine/liana foliar cover	3-5%
Grass/grasslike foliar cover	15-40%
Forb foliar cover	2-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 16. Soil surface cover

Tree basal cover	1-10%
Shrub/vine/liana basal cover	0-5%
Grass/grasslike basal cover	2-8%
Forb basal cover	0-5%
Non-vascular plants	0%
Biological crusts	0%
Litter	90-100%

Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	5-20%
Bedrock	0%
Water	0%
Bare ground	0-5%

Table 17. Woody ground cover

Downed wood, fine-small (<0.40" diameter; 1-hour fuels)	—
Downed wood, fine-medium (0.40-0.99" diameter; 10-hour fuels)	—
Downed wood, fine-large (1.00-2.99" diameter; 100-hour fuels)	—
Downed wood, coarse-small (3.00-8.99" diameter; 1,000-hour fuels)	—
Downed wood, coarse-large (>9.00" diameter; 10,000-hour fuels)	—
Tree snags** (hard***)	—
Tree snags** (soft***)	—
Tree snag count** (hard***)	40-80 per acre
Tree snag count** (hard***)	

* **Decomposition Classes:** N - no or little integration with the soil surface; I - partial to nearly full integration with the soil surface.

** >10.16cm diameter at 1.3716m above ground and >1.8288m height--if less diameter OR height use applicable down wood type; for pinyon and juniper, use 0.3048m above ground.

*** **Hard** - tree is dead with most or all of bark intact; **Soft** - most of bark has sloughed off.

Table 18. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	1-3%	0-1%
>0.5 <= 1	—	1-3%	0-5%	1-3%
>1 <= 2	—	3-5%	0-10%	0-5%
>2 <= 4.5	—	3-5%	0-15%	—
>4.5 <= 13	20-40%	10-20%	—	—
>13 <= 40	50-70%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Figure 17. Plant community growth curve (percent production by month).
TX3771, Hardwood/Shrub Woodland Community. Hardwood and shrub woodland community..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2	3	10	15	20	18	5	4	10	7	4	2

State 3

Exotic Grass Savannah State

This state is a result of significant ground disturbing mechanical brush management. Depending upon the management goals, the site may have been seeded to native or non-native species or a combination. Non-native grasses can also invade without seeding being introduced through hay, livestock, or wildlife. Herbaceous species diversity and site resilience has been significantly reduced.

Dominant plant species

- Bermudagrass (*Cynodon dactylon*), grass
- Johnsongrass (*Sorghum halepense*), grass

Community 3.1

Exotic Grass Oak Savannah Community



Figure 18. Exotic Grass/Oak Savannah Community on Gravelly Redland Ecological Site.

This community phase is characterized by a mixture of exotic and native grasses with live oak canopy of about 10%. Seeded or non-native grasses include naturalized species such as King Ranch bluestem, bermudagrass, Johnsongrass (*Sorghum halepense*), silky bluestem (*Dichanthium sericeum*), kleingrass, and many others. The King Ranch bluestem has the most potential of the exotics to eventually dominate the site. When this occurs, there has been a dramatic reduction in the native forb and legume diversity. These monoculture type communities may be too dense for gallinaceous wildlife. Through the re-introduction of fire and prescribed grazing, plus reseeding of native forbs and grasses, this site can be restored to something resembling the historic plant. Utilizing native plants in the re-seeding will greatly benefit wildlife species such as deer, turkey, quail, and other birds. Total production for this site may be similar to the productive potential of this site in reference condition except the majority of the plant communities are exotics. Many times the exotic grasses can provide the same hydrologic functions as do the natives.

Table 19. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1700	2250	3825
Tree	200	300	450
Shrub/Vine	50	150	225
Forb	0	0	0
Total	1950	2700	4500

Table 20. Ground cover

Tree foliar cover	1-3%
Shrub/vine/liana foliar cover	0-5%
Grass/grasslike foliar cover	15-40%
Forb foliar cover	0-5%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%

Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 21. Soil surface cover

Tree basal cover	1-3%
Shrub/vine/liana basal cover	0-3%
Grass/grasslike basal cover	10-15%
Forb basal cover	0-3%
Non-vascular plants	0%
Biological crusts	0%
Litter	80-90%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	5-20%
Bedrock	0%
Water	0%
Bare ground	0-5%

Table 22. Woody ground cover

Downed wood, fine-small (<0.40" diameter; 1-hour fuels)	—
Downed wood, fine-medium (0.40-0.99" diameter; 10-hour fuels)	—
Downed wood, fine-large (1.00-2.99" diameter; 100-hour fuels)	—
Downed wood, coarse-small (3.00-8.99" diameter; 1,000-hour fuels)	—
Downed wood, coarse-large (>9.00" diameter; 10,000-hour fuels)	—
Tree snags** (hard***)	—
Tree snags** (soft***)	—
Tree snag count** (hard***)	40-80 per acre
Tree snag count** (hard***)	

* **Decomposition Classes:** N - no or little integration with the soil surface; I - partial to nearly full integration with the soil surface.

** >10.16cm diameter at 1.3716m above ground and >1.8288m height--if less diameter OR height use applicable down wood type; for pinyon and juniper, use 0.3048m above ground.

*** Hard - tree is dead with most or all of bark intact; Soft - most of bark has sloughed off.

Table 23. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	1-3%	0-1%
>0.5 <= 1	—	0-3%	3-5%	0-3%
>1 <= 2	—	0-8%	10-15%	0-10%
>2 <= 4.5	—	0-10%	30-50%	—
>4.5 <= 13	—	—	—	—
>13 <= 40	1-10%	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

Transition T1A

State 1 to 2

A transition occurs due to a lack of brush management with mechanical means, fire or targeted goat/sheep grazing. Grazing deferment alone will not halt the increase of woody species. Hydrologic characteristics are altered by increased woody species. Now, energy flows more through woody plants than herbaceous plants.

Restoration pathway R2A

State 2 to 1

R2A Restoration includes some form of brush management and many times an integrated approach utilizing several methods. In some cases of severe long-term overharvesting of the desired plants, replanting may be necessary. Prescribed burning is an option once the fine fuel load has recovered so prescribed grazing will be essential.

Transition T2A

State 2 to 3

This transition is driven by ground disturbance brush management with the replanting of exotic grasses, either as a mixture or single species. Exotic plants can also be introduced from hay brought to the site or from livestock and wildlife. Hydrologic characteristics are anticipated to be similar to the Mixed Grass Prairie Community.

Additional community tables

Table 24. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Tall/Midgrasses			700–1575	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	700–1200	—
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	500–1000	—
2	Tallgrasses			400–900	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	100–500	—
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	100–500	—
3	Midgrasses			400–900	
	composite dropseed	SPCO16	<i>Sporobolus compositus</i>	200–300	—
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	150–225	—
	plains lovegrass	ERIN	<i>Eragrostis intermedia</i>	100–150	—
	Texas cupgrass	ERSE5	<i>Eriochloa sericea</i>	100–150	—

	green sprangletop	LEDU	<i>Leptochloa dubia</i>	100–150	–
	vine mesquite	PAOB	<i>Panicum obtusum</i>	75–150	–
	silver bluestem	BOSA	<i>Bothriochloa saccharoides</i>	75–150	–
4	Shortgrasses			100–225	
	buffalograss	BODA2	<i>Bouteloua dactyloides</i>	25–75	–
	curly-mesquite	HIBE	<i>Hilaria belangeri</i>	25–75	–
	Hall's panicgrass	PAHA	<i>Panicum hallii</i>	25–50	–
	slim tridens	TRMU	<i>Tridens muticus</i>	25–50	–
	slim tridens	TRMUE	<i>Tridens muticus</i> var. <i>elongatus</i>	25–50	–
	Texas grama	BORI	<i>Bouteloua rigidiseta</i>	25–50	–
	red grama	BOTR2	<i>Bouteloua trifida</i>	25–50	–
	hairy woollygrass	ERPI5	<i>Erioneuron pilosum</i>	25–50	–
	Wright's threeawn	ARPUW	<i>Aristida purpurea</i> var. <i>wrightii</i>	25–50	–
5	Cool-season Grasses			100–225	
	cedar sedge	CAPL3	<i>Carex planostachys</i>	50–150	–
	wildrye	ELYMU	<i>Elymus</i>	50–150	–
	Texas wintergrass	NALE3	<i>Nassella leucotricha</i>	50–150	–
Forb					
6	Forbs			100–225	
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	25–75	–
	blazing star	LIATR	<i>Liatris</i>	25–75	–
	Nuttall's sensitive-briar	MINU6	<i>Mimosa nuttallii</i>	25–50	–
	slimflower scurfpea	PSTE5	<i>Psoraleidium tenuiflorum</i>	25–50	–
	snoutbean	RHYNC2	<i>Rhynchosia</i>	25–50	–
	zarzabacoa comun	DEIN3	<i>Desmodium incanum</i>	25–50	–
	bundleflower	DESMA	<i>Desmanthus</i>	25–50	–
	blacksamson echinacea	ECAN2	<i>Echinacea angustifolia</i>	0–50	–
	indigo	INDIG	<i>Indigofera</i>	25–50	–
	trailing krameria	KRLA	<i>Krameria lanceolata</i>	25–50	–
	Forb, annual	2FA	<i>Forb, annual</i>	0–50	–
	Indian mallow	ABUTI	<i>Abutilon</i>	25–50	–
Shrub/Vine					
7	Shrubs/Vines			100–225	
	catclaw acacia	ACGR	<i>Acacia greggii</i>	50–150	–
	littleleaf sumac	RHMI3	<i>Rhus microphylla</i>	25–100	–
	greenbrier	SMILA2	<i>Smilax</i>	25–100	–
	bully	SIDER2	<i>Sideroxylon</i>	25–75	–
	pricklypear	OPUNT	<i>Opuntia</i>	25–75	–
	jointfir	EPHED	<i>Ephedra</i>	25–75	–
	stretchberry	FOPU2	<i>Forestiera pubescens</i>	25–50	–
	algerita	MATR3	<i>Mahonia trifoliolata</i>	25–50	–
	Texas Hercules' club	ZAH12	<i>Zanthoxylum hirsutum</i>	0–25	–
Tree					

8	Trees			100–225	
	Texas live oak	QUFU	<i>Quercus fusiformis</i>	100–225	–
	hackberry	CELT1	<i>Celtis</i>	25–100	–
	western soapberry	SASAD	<i>Sapindus saponaria</i> var. <i>drummondii</i>	25–75	–
	elm	ULMUS	<i>Ulmus</i>	0–50	–
	honey mesquite	PRGL2	<i>Prosopis glandulosa</i>	0–25	–
	blackjack oak	QUMA3	<i>Quercus marilandica</i>	0	–
	post oak	QUST	<i>Quercus stellata</i>	0	–

Animal community

Many types of grassland insects, reptiles, birds, and mammals frequented the Mixed Grass Prairie Community (1.1) of the site, either as their base habitat or maneuvering from the adjacent sites. Small mammals included many kinds of rodents, jackrabbit, cottontail rabbit, raccoon, skunk, opossum, and armadillo. Predators included coyote, red fox, gray fox, bobcat, and occasionally mountain lion. Game birds, songbirds, and birds of prey were indigenous or frequent users. Most are still plentiful. Bison and pronghorn antelope, however, are no longer present. Native white-tailed deer and many species of exotic deer utilize the Gravelly Redland site in its various states. Several species of exotic wildlife have been introduced on the site including deer, such as axis, sika, and fallow; antelope, such as sable, oryx, blackbuck, and nilgai; and sheep, such as Barbados (mouflon) and aoudad. Their numbers must be included along with livestock and native wildlife, primarily white-tailed deer, in any conservation plan. Axis deer have the ability to shift their diet among plant groups giving them a competitive advantage over the native white-tailed deer. Feral hogs may also be present on the site. Hogs can be damaging to the plant community and ranch improvement structures if their numbers are not managed. Deer, turkey, and quail particularly favor the habitat provided by the Midgrass Oak Savannah Community (1.2).

The site is suitable for the production of livestock, including cattle, sheep, and goats. The site in reference condition was very suited to primary grass eaters such as cattle. As retrogression occurs and woody plants invade, the community becomes better habitat for sheep, goats, deer, and other wildlife because of the desirable browse and cool-season grasses. Cattle, sheep, and goats should be stocked in proportion to the available grass, forb, and browse forage keeping deer competition for forbs and browse in mind. Deer populations must also be kept within limits of the habitat sustainability even if the site is managed exclusively for deer. If the animal numbers are not kept in balance with herbage and browse production through prescribed grazing management and good wildlife population management, the late Midgrass Oak Savannah Community (1.2) will have little to offer as habitat except cover.

Plant Preference by Animal Kind:

This rating system provides general guidance as to animal forage preference for plant species. It also indicates possible competition and diet overlap between kinds of herbivores. Grazing preference changes from time to time, especially between seasons, and between animal kinds and classes. An animal's preference or avoidance of certain plants is learned over time through grazing experience and maternal learning (<http://extension.usu.edu/behave/Grazing> accessed 8/30/13). Preference does not necessarily reflect the ecological status of the plant within the plant community. For wildlife, plant preferences for food are rated. Refer to detailed habitat guides for a more complete description of a species habitat needs.

Legend: P=Preferred D=Desirable U=Undesirable N=Not Consumed T=Toxic X=Used, but not degree of utilization unknown

Preferred – Percentage of plant in animal diet is greater than it occurs on the land

Desirable – Percentage of plant in animal diet is similar to the percentage composition on the land

Undesirable – Percentage of plant in animal diet is less than it occurs on the land

Not Consumed – Plant would not be eaten under normal conditions. It is only consumed when other forages not available. This can also include plants that are unavailable during parts of the year.

Toxic – Rare occurrence in diet and, if consumed in any tangible amounts results in death or severe illness in animal (Hart, 2003). (Note: many plants can be good forage but toxic at certain doses or at certain times of the year. Animals in poor condition are most susceptible.)

Hydrological functions

The climate affecting the hydrology of the Gravelly Redland Ecological Site is humid subtropical with approximately two-thirds of annual rainfall occurring during the April to September period. Rainfall during this period generally falls during thunderstorms and fairly large amounts of rain may fall in a short time. Because of the topography and vegetation, quality of surface runoff is high with low erosion and sedimentation levels. Sinkholes and crevices facilitate water movement to deeper root zones and below, contributing some to the recharge of aquifers. The site is well drained. Runoff is moderate, with good cover. Erosion could be a hazard with excessive defoliation because of the slope.

Under reference conditions, the grassland vegetation intercepted and utilized much of the incoming rainfall in the soil solum. Only during extended rains or heavy thunderstorms was there much runoff. Litter and soil movement was slight. Standing plant cover, duff, and organic matter decrease and surface runoff increase as the Oak Savannah State (1) transitions to the Midgrass Oak Savannah State (2). These processes continue in the interstitial spaces in the Shrubland (2). The woody plants compete for moisture and sunlight with the remaining grasses and forbs; further reducing production and ground cover in openings. Decreased litter and more bare ground allow erosion from soils in openings between trees.

Once the Mixed Brush Shrubland Community (2.1) canopy surpasses 50 percent, the hydrology and ecological processes, nutrient cycling, and energy flow stabilize within the woody plant canopy; although a significant departure from the Oak Savannah State (1). Evaporation and interception losses are higher resulting in less moisture reaching the soil. The deeper-rooted woody plants are able to extract water from greater depths than most grasses so less water will be available for down-slope movement except through crevices.

Recreational uses

The Gravelly Redland Site is well suited for many outdoor recreational uses including hunting, hiking, camping, equestrian, and bird watching. This site along with adjacent upland sites provides diverse scenic beauty.

Wood products

Posts and specialty wood products are made from juniper, mesquite, oak and many shrubs. Mesquite and oak are used for firewood and charcoal.

Other products

Jams and jellies are made from many fruit-bearing species, such as algerita. Seeds are harvested from many plants for commercial sale. Grasses and forbs are harvested by the dried-plant industry for sale in dried flower arrangements. Honeybees are utilized to harvest honey from the many flowering plants, such as mesquite.

Other information

None.

Inventory data references

Information presented was derived from the revised Gravelly Redland Range Site description, literature, limited NRCS clipping data (417s), field observations, professional consensus, and personal contacts with range-trained personnel.

Other references

Anderson, J.R., C.A. Taylor, Jr., C.J. Owens, J.R. Jackson, D.K. Steele, and R. Brantley. 2013. Using experience and supplementation to increase juniper consumption by three different breeds of sheep. *Rangeland Ecol. Management.* 66:204-208. March.

Archer S. 1994. Woody plant encroachment into southwestern grasslands and savannas: rates, patterns and

- proximate causes. in: Ecological implications of livestock herbivory in the West, pp.13-68. Edited by M. Vavra, W. Laycock, R. Pieper, Society for Range Management Publication. , Denver, Colorado.
- Bestelmeyer, B.T., J.R. Brown, K.M. Havsted, R. Alexander, G. Chavez, and J.E. Hedrick. 2003. Development and use of state-and-transition models for rangelands. *Journal of Range Management*. 56(2): 114-126.
- Bushland, R.C. 1985. Eradication program in the southwestern United States. Symposium on eradication of the screwworm from the United States and Mexico. *Misc. Pub. Entomol. Soc. Am.*, 62:12-15.
- Foster, J.H. 1917. The spread of timbered areas in central Texas. *Journal of Forestry* 15:442-445.
- Frost, C. C. 1998. Presettlement fire frequency regimes of the United States: a First Approximation. Tall Timbers Fire Ecology Conference Proceedings. No. 20. Tall Timbers Research Station. Tallahassee, FL.
- Fuhlendorf, S. D., and Engle D.M., Kerby J., and Hamilton R. 2008. Pyric Herbivory: rewilding Landscapes through the Recoupling of Fire and Grazing. *Conservation Biology*. Volume 23, No. 3, 588-598.
- Hamilton W. and D. Ueckert. 2005. Rangeland Woody Plant Control--Past, Present, and Future. Chapter 1 in: *Brush Management-Past, Present, and Future*. Texas A & M University Press. Pp.3-16.
- Hanselka, W., R. Lyons, and M. Moseley. 2009. *Grazing Land Stewardship – A Manual for Texas Landowners*. Texas AgriLife Communications, <http://agrilifebookstore.org>.
- Hart, C., R.T. Garland, A.C. Barr, B.B. Carpenter, and J.C. Reagor. 2003. Toxic Plants of Texas. *Texas Cooperative Extension Bulletin B-6103* 11-03.
- Inglis, J. M. 1964. A History of Vegetation on the Rio Grande Plains. Texas Parks and Wildlife Department, Bulletin No. 45. Austin, Texas.
- Massey, C.L. 2009. The founding of a town – The Gugger and Benke families. *Helotes Echo*, July 1, 2009.
- Natural Resources Conservation Service. 1994. *The Use and Management of Browse in the Edwards Plateau of Texas*. Temple, Texas.
- Plant symbols, common names, and scientific names according to USDA/NRCS Texas Plant List (Unpublished)
- Pyne, S.J. 1982. *Fire in America*. Princeton University Press, Princeton, NJ.
- Roemer, Ferdinand Von. 1983. *Roemer's Texas*. Eakins Press.
- Schmidly, D.J. 2002. *Texas natural history: a century of change*. Texas Tech University Press, Lubbock.
- Scifres, C.J. and W.T. Hamilton. 1993. *Prescribed Burning for Brush Management: The South Texas Example*. Texas A & M University Press, 245 pp.
- Smeins, F., S. Fuhlendorf, and C. Taylor, Jr. 1997. Environmental and Land Use Changes: A Long Term Perspective. Chapter 1 in: *Juniper Symposium 1997*. Texas Agricultural Experiment Station. Pp 1-21.
- Taylor, C.A. (Ed.). 1997. *Texas Agriculture Experiment Station Technical Report 97-1 (Proceedings of the 1997 Juniper Symposium)*, Sonora Texas, pp. 9-22.
- Teer, J.G., J.W. Thomas, and E.A. Walker. 1965. Ecology and Management of White-tailed Deer in the Llano Basin of Texas. *Wildlife Monographs* 10: 1-62.
- Thurrow, T.O. and J.W. Hester. 1997. 1997 Juniper Symposium. Texas Agricultural Experiment Station, The Texas A&M University System. Tech. Rep. 97-1. January 9-10, 1997. San Angelo, Texas
- USDA-NRCS (Formerly Soil Conservation Service) Range Site Description (1972)

- Vines, R.A. 1984. Trees of Central Texas. University of Texas Press. Austin, Texas.
- Weninger, D. 1984. The Explorer's Texas. Eakin Press; Waco, Texas.
- Wilcox, B.P. and T.L. Thurow. 2006. Emerging Issues in Rangeland Ecohydrology: Vegetation Change and the Water Cycle. Rangeland Ecol. Management. 59:220-224, March.
- Wilcox, B.P., Y. Huang, and J.W. Walker. 2008. Long-term trends in stream flow from semiarid rangeland: uncovering drivers of change. Global Change Biology 14: 1676-1689, doi:10.1111/j.1365.2486.2008.01578.
- Wilcox, B.P., W.A. Dugas, M.K. Owens, D.N Ueckert, and C.R. Hart. 2005. Shrub Control and Water Yield on Texas Rangelands: Current State of Knowledge. Texas Agricultural Experiment Station Research Report 05-1.
- Wills, Frederick. 2006. Historic Vegetation of Camp Bullis and Camp Stanley, Southeastern Edwards, Plateau. Texas. Texas Journal of science. 58(3):219-230.
- Wright, H.A. and A.W. Bailey. 1982. Fire Ecology: United States and Southern Canada. John Wiley & Sons, Inc.
- Wu, B.X., E.J. Redeker, and T.L. Thurow. 2001. Vegetation and Water Yield Dynamics in an Edwards Plateau Watershed. Journal of Range Management. 54:98-105. March 2001.
<http://extension.usu.edu/behave/> (accessed 9/4/2013)

Contributors

Dr. Joseph Schuster
Mark Moseley
Travis Waiser, MLRA Leader, NRCS, Kerrville, TX

Approval

Bryan Christensen, 9/19/2023

Acknowledgments

Site Development and Testing Plan:

Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high-intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document. Annual reviews of the Project Plan are to be conducted by the Ecological Site Technical Team.

Reviewers and Technical Contributors:

Joe Franklin, Zone Range Management Specialist, NRCS, San Angelo Zone Office, Texas
Ryan McClintock, Biologist, NRCS, San Angelo Zone Office, Texas
Jessica Jobes, Project Leader, Kerrville Soil Survey Office, NRCS, Kerrville, Texas
Travis Waiser, Soil Scientist, NRCS, Kerrville Soil Survey Office, Texas
Bryan Hummel, Natural Resources Technician, Joint Base San Antonio-Camp Bullis, Texas
Ann Graham, Editor, NRCS, Temple, Texas
Charles Anderson, NRCS RMS, San Angelo, Texas
Homer Sanchez, NRCS, State RMS, Temple, Texas
Justin Clary, NRCS, RMS, Temple, Texas

QC/QA completed by:

Bryan Christensen, SRESS, NRCS, Temple, TX
Erin Hourihan, ESDQS, NRCS, Temple, TX

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	San Angelo NRCS Zone Office
Contact for lead author	325-944-0147
Date	04/08/2013
Approved by	Colin Walden
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

2. **Presence of water flow patterns:** Some water flow patterns are expected due to runoff from Adobe, Steep Adobe, and Low stony hill sites.

3. **Number and height of erosional pedestals or terracettes:** None.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0-5% bare ground.

5. **Number of gullies and erosion associated with gullies:** None.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None.

7. **Amount of litter movement (describe size and distance expected to travel):** Little or no litter movement or deposition during normal rainfall events.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Stability range is expected to be 5-6.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Reddish brown clay surface with medium angular blocky structure, 0 to 8 inches Thick A horizon. Gravelly modifier. 1 to 3 percent

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** High canopy, basal cover and density with small interspaces should make rainfall impact negligible. This site has well drained soils, very slowly permeable with 1 to 5 percent slopes which allow negligible runoff and erosion.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm-season tallgrasses Warm-season midgrasses
- Sub-dominant: Warm-season midgrasses Trees Shrubs Forbs
- Other: Warm-season shortgrasses Cool-season grasses
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Perennial grasses will naturally exhibit a minor amount (less than 5%) of senescence and some mortality every year.
-
14. **Average percent litter cover (%) and depth (in):** >90 percent litter, 0.5 to 1.0 inch depth.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 2000 to 4500 pounds per acre.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Ashe juniper, pricklypear, yucca, tasajillo, pricklyash, lotebush, mesquite, King Ranch bluestem, silky bluestem.
-
17. **Perennial plant reproductive capability:** All perennial species should be capable of reproducing every year unless disrupted by extended drought, overgrazing, wildfire, insect damage, or other events occurring immediately prior to, or during the reproductive phase.
