

Ecological site R081CY574TX Shallow 29-35 PZ

Last updated: 9/19/2023
Accessed: 04/26/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Zone RMS, San Angelo, Texas
Contact for lead author	325-944-0147
Date	02/14/2012
Approved by	Colin Walden
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** None.

- Presence of water flow patterns:** None, except following extremely high intensity storms where short flow patterns may appear.

- Number and height of erosional pedestals or terracettes:** None

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0-10% bare ground randomly distributed throughout in small and non-connected areas.

- Number of gullies and erosion associated with gullies:** None

- Extent of wind scoured, blowouts and/or depositional areas:** None

- Amount of litter movement (describe size and distance expected to travel):** Under normal rainfall, little litter

movement should be minimal and short (<6").

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Reference condition soil surface is resistant to erosion. Stability class range is expected to be 5-6.
-

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is very dark grayish brown silty clay 9 inches thick that contains soft bodies of calcium carbonate. Moderately alkaline. SOM is 1-3%.
-

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The savannah of tallgrasses, midgrasses, forbs and trees having adequate litter and little bare ground provide for maximum infiltration and little runoff under normal rainfall events.
-

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No evidence of compaction.
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Warm-season tallgrasses

Sub-dominant: Warm-season midgrasses Trees

Other: Forbs Cool-season grasses = Forbs = Warm-season shortgrasses Shrubs

Additional: Forbs make up 4%, Shrubs make up 2% and Trees make up 4%.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Perennial grasses will naturally exhibit a minor amount (less than 5%) of senescence and some mortality every year.
-

14. **Average percent litter cover (%) and depth (in):** Dominant litter is herbaceous.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 970 to 2,820 pounds per year
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site: Ashe Juniper is dominant, honey mesquite, prickly pear, bermudagrass, Johnsongrass, King Ranch bluestem.

17. **Perennial plant reproductive capability:** All perennial species should be capable of reproducing every year unless disrupted by extended drought, overgrazing, wildfire, insect damage, or other events occurring immediately prior to, or during the reproductive phase.
-