

Ecological site R083BY019TX Gray Sandy Loam

Last updated: 9/19/2023
Accessed: 05/13/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 083B–Western Rio Grande Plain

Major Land Resource Area (MLRA) 83B It makes up about 9,285 square miles (24,060 square kilometers). The border towns of Del Rio, Eagle Pass, Laredo, and Zapata are in this MLRA. Interstate 35 crosses the area just north of Laredo. The Amistad National Recreation Area is just outside this MLRA, northwest of Del Rio, and the Falcon State Recreation Area is southeast of Laredo. Laughlin Air Force Base is just east of Del Rio. This area is comprised of inland, dissected coastal plains.

Classification relationships

USDA-Natural Resources Conservation Service, 2006.
-Major Land Resource Area (MLRA) 83B

Ecological site concept

The Gray Sandy Loam refers to the gray-colored, sandy loam surfaces found on the ecological site. High amounts of calcium carbonates in the upper soil profile are responsible for the gray colors and alkalinity.

Associated sites

R083BY018TX	Clay Flat
R083BY004TX	Shallow Sandy Loam
R083BY005TX	Shallow
R083BY012TX	Ramadero
R083BY025TX	Clay Loam
R083BY003TX	Gravelly Ridge
R083BY011TX	Claypan Prairie

Similar sites

R083AY019TX	Gray Sandy Loam
R083CY019TX	Gray Sandy Loam
R083DY019TX	Gray Sandy Loam

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Bernardia myricifolia</i> (2) <i>Leucophyllum frutescens</i>
Herbaceous	(1) <i>Heteropogon contortus</i> (2) <i>Trichloris pluriflora</i>

Physiographic features

Sites were formed on nearly level to gently sloping interfluves and ridges on the inland, dissected Coastal Plains. Surfaces are linear to convex. Slope ranges from 0 to 5 percent but are mostly less than 3 percent. Elevation is 150 to 880 feet.

Table 2. Representative physiographic features

Landforms	(1) Coastal plain > Interfluve (2) Coastal plain > Ridge
Runoff class	Negligible to medium
Flooding frequency	None
Ponding frequency	None
Elevation	46–268 m
Slope	0–5%
Aspect	Aspect is not a significant factor

Climatic features

MLRA 83B mainly has a subtropical steppe climate along the Rio Grande River and subtropical subhumid climates in La Salle and McMullen counties. Winters are dry and mild and the summers are hot. Tropical maritime air masses predominate throughout spring, summer and fall. Modified polar air masses exert considerable influence during winter, creating a continental climate characterized by large variations in temperature. Peak rainfall occurs late in spring and a secondary peak occurs early in fall. Most heavy thunderstorm activities occur during the summer months. July is hot and dry with little weather variations. Rainfall increases again in late August and September as tropical disturbances increase and become more frequent as the storms dissipate. Tropical air masses from the Gulf of Mexico dominate during the spring, summer and fall. Prevailing winds are southerly to southeasterly throughout the year except in December when winds are predominately northerly.

Table 3. Representative climatic features

Frost-free period (characteristic range)	231-321 days
Freeze-free period (characteristic range)	313-365 days
Precipitation total (characteristic range)	508 mm
Frost-free period (actual range)	214-365 days
Freeze-free period (actual range)	260-365 days
Precipitation total (actual range)	483-533 mm
Frost-free period (average)	270 days
Freeze-free period (average)	340 days
Precipitation total (average)	508 mm

Climate stations used

- (1) FALCON DAM [USC00413060], Roma, TX
- (2) LAREDO 2 [USC00415060], Laredo, TX
- (3) CATARINA [USC00411528], Asherton, TX
- (4) CRYSTAL CITY [USC00412160], Crystal City, TX
- (5) DEL RIO 2 NW [USC00412361], Del Rio, TX
- (6) EAGLE PASS 3N [USC00412679], Eagle Pass, TX
- (7) ZAPATA 1 S [USC00419976], Zapata, TX
- (8) DEL RIO INTL AP [USW00022010], Del Rio, TX

Influencing water features

Water features do not influence this site.

Wetland description

N/A.

Soil features

The soil are moderately deep to very deep, moderate to moderately slowly permeable over weakly to strongly cemented sandstone. The site gets its name from the gray colors in the soil resulting from calcium carbonates, making the soils alkaline. Soil series correlated to this site include: Aguillares, Choke, Copita, Lenocita, and Tonio.

Table 4. Representative soil features

Parent material	(1) Residuum—sedimentary rock
Surface texture	(1) Fine sandy loam (2) Sandy clay loam (3) Silty clay loam
Family particle size	(1) Fine-loamy (2) Fine-silty
Drainage class	Well drained
Permeability class	Moderate to moderately slow
Soil depth	51–203 cm
Surface fragment cover <=3"	0–1%
Surface fragment cover >3"	0%

Available water capacity (0-101.6cm)	10.16–17.78 cm
Calcium carbonate equivalent (0-101.6cm)	1–35%
Electrical conductivity (0-101.6cm)	0–4 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–10
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–10%
Subsurface fragment volume >3" (Depth not specified)	0–1%

Ecological dynamics

The accounts of early explorers and settlers suggest that the Rio Grande Plains was likely a vast mosaic of open grassland, savannah, and shrubland. While moving in 1691 out of Maverick County and into Zavala County, Don Domingo de Teran found after crossing the Nueces River “the country was level and covered with mesquites and cats’ claw.” In 1849, Michler described south Texas as “concerning the land both on the Frio and the Leona, from these rivers back, that it may be divided into four parallel strips-the first, next to the river, consisting of heavy timber, and a heavy black soil, the second, a mesquite flat, of small width, and the soil of a lighter nature, and very fertile; the third, a range of low hills, covered with loose stones, and thick chaparral; the fourth, a wide-open prairie.” Lehman indicates, “thus while it is quite true that the Rio Grande Plains once had fewer woody plants and more grass than now, it is also true that an ample seed stock of shrubs and trees has been widely distributed for as long as man has known.” The vegetation structure likely varied from place-to-place depending on topography, soil properties, and time since the last major disturbance.

Large numbers of domestic livestock grazed South Texas as early as the mid-1700’s. Formal deeds to properties from the Spanish and Mexican governments came in the late 1760’s with much larger blocks granted in the decades to follow. Lehman indicated, “in 1757, the official Spanish census showed residents of Camargo and Reynosa in the lower Rio Grande owning over 90,000 sheep and goats. By way of contrast, combined numbers of cattle, oxen, horses, mules and burros were less than 16,000.” By the mid-1800’s, according to Lehman’s figures from the U. S. Census of 1889, “there were a minimum of 1,644,268 sheep-fully 45 percent of Texas total population, grazing south of the Nueces River.” According to Inglis, “the Rio Grande Plains had the four-leading sheep producing counties in the state and ten of the top fifteen sheep producing counties were in South Texas. The peak decade was 1880 to 1890, at times exceeding two million head.” These domestic animals were in addition to bison, antelope, deer, and large herds of wild horses. It is obvious from early accounts, that much of the Rio Grande Plains was periodically grazed hard by both domestic animals and wild populations as early as the early to mid-1700’s. It may be that overgrazing by sheep and goats could have suppressed the many shrubs, reduced shrub canopy, and arrested shrub seedlings.


With the arrival of European man, the South Texas area was fenced and, in many instances, stocked beyond its capability to sustain forage. This overstocking led to a reduced fire frequency and intensity, creating an opportunity for woody shrubs to increase across the landscape. As the natural graze-rest cycles were altered and stocking rates continued to exceed the natural carrying capacity of the land, midgrasses were replaced by shortgrasses and the ground cover was opened so additional annual and perennial forbs also increased. Drought certainly enhanced this effect. As prolonged overgrazing continued, shrub cover increased. Shortgrasses became dominant and forage production decreased. This change in plant cover and structure further decreased fire frequency and intensity, favoring shrub establishment and dominance.

The plant communities of this site are dynamic varying in relation to fire, periodic drought, and wet cycles. Periodic fires were set by either Native Americans or started naturally by lightning. Fire did not play as important a role on this site as in deeper more productive sites due to lower production of grasses to burn. Because of large amounts of gravel in the soil, available water holding capacity is greatly reduced. This causes highly variable forage

production and minimal grass production during dry years. The historic community of this site was influenced to some extent by periodic grazing by herds of buffalo and wild horses. Herds of buffalo and wild horses would come into an area, graze it down, and then not come back for many months or even years depending upon the availability of water. This long deferment period allowed recovery of the grasses and forbs which served as fuel load. More than likely, fire occurred following years of good rainfall followed by a dry season. The fire frequency for this area is interpreted to be four to six years (Frost, 1998).

Presently, the Gray Sandy Loam is a community of woody shrubs exceeding 50 percent canopy, with the interspaces dominated by shortgrasses such as common curly mesquite (*Hilaria berlanderi*), fall witchgrass (*Digitaria cognata*), Hall's panicum (*Panicum hallii*), perennial threeawn (*Aristida purpurea*), and tumblegrass (*Schedonnardus paniculatus*). If drought and/or grazing denude the site, soils will cap over and infiltration of rainfall will be reduced significantly making the site much more droughty. When in this condition, this site recovers very slowly, and mechanical manipulation will be required to reduce shrub canopy and break the soil crust.

State and transition model



- 1.1A – Heavy Continuous Grazing, No Fire, No Brush Management
- 1.2A – Prescribed Grazing, Prescribed Burning, Brush Management
- 2.1A – Heavy Continuous Grazing, No Fire, Brush Invasion
- 2.2A – Prescribed Grazing, Prescribed Burning, Brush Management
- T1A – Heavy Continuous Grazing, No Fire, Brush Invasion
- R2A – Brush Management, Prescribed Burning, Prescribed Grazing
- T2A – Brush Management, Range Planting, Pasture Planting
- T3A – Heavy Continuous Grazing, No Fire, Brush Invasion
- T1B – Brush Management, Pasture Planting, Range Planting, Prescribed Grazing

Figure 8. STM

State 1
Grassland

Dominant plant species

- multiflower false Rhodes grass (*Trichloris pluriflora*), grass
- tanglehead (*Heteropogon contortus*), grass

Community 1.1
Midgrass Dominant

The reference community for the Gray Sandy Loam is an open grassland dominated by midgrasses, perennial forbs, and interspersed with occasional woody shrubs. Dominant midgrasses include multi-flowered false rhodesgrass, tanglehead, plains bristlegrass, Arizona cottontop, and silver bluestem. There are shortgrasses present but in limited amounts. Perennial forbs include bush sunflower, orange zexmenia, daleas, snoutbeans, and bundleflowers. Scattered woody plants occur making up less than 10 percent of the total composition. These included guajillo, granjeno, cenizo, wolfberry, blackbrush acacia and many others.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	897	2578	3699
Forb	168	224	336
Shrub/Vine	112	168	224
Tree	—	—	—
Total	1177	2970	4259

Figure 10. Plant community growth curve (percent production by month). TX4800, Midgrass Dominant Community. Warm-season midgrasses with forbs and shrubs..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	2	10	20	20	5	8	15	10	6	2

Community 1.2
Midgrass/Shortgrass Dominant



Figure 11. 1.2 Midgrass/Shortgrass Dominant Community

This community is resilient, still influenced of fire, and can easily be transitioned back to community 1.1. This community is very stable and is often maintained for wildlife purposes. Continued heavy grazing coupled with drought cycles has caused the dominant midgrasses to decrease in composition. The opening of the midgrass canopy causes shortgrasses, forbs and woody species to increase. Reduction in stocking rates, periodic rest, and

increased fire frequency will maintain this community, or even transition back to the reference community. But, if over grazing continues, midgrasses will continue to decline, fire frequency and intensity will decrease, and the plant community will continue towards the Shrubland State. Such species as tanglehead, multi-flowered false Rhodes grass, Arizona cottontop, and plains bristlegrass diminish and are replaced by species like pink pappusgrass, hooded windmillgrass, and perennial three-awns. Short grasses such as curly mesquite, Hall's panicum, tumble windmillgrass, and sand dropseed also increase.

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	673	2242	3138
Forb	196	280	392
Shrub/Vine	196	252	336
Tree	–	–	–
Total	1065	2774	3866

Figure 13. Plant community growth curve (percent production by month). TX4800, Midgrass Dominant Community. Warm-season midgrasses with forbs and shrubs..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	2	10	20	20	5	8	15	10	6	2

Pathway 1.1A Community 1.1 to 1.2

A shift to the 1.2 Community occurs if the Midgrass Community is weakened by excessive leaf removal. Drought hastens the process. A reduction in midgrass also corresponds in a reduction of fuel loading needed for fire to effectively suppress woody species.

Pathway 1.2A Community 1.2 to 1.1

Managerial activities that restore the hydrologic cycle, the energy capture by midgrasses, and ground cover will move the 1.2 Community toward the Midgrass Community (1.1). Utilizing historic ecological disturbances such as herbivory and fire in constructive amounts are needed. Selective brush management may also be needed.

State 2 Shrubland

Dominant plant species

- guajillo (*Acacia berlandieri*), shrub
- spiny hackberry (*Celtis ehrenbergiana*), shrub

Community 2.1 Mid/Shortgrass Shrubland Complex

The Mid/Shortgrass Shrubland Complex represents a community where thresholds have been crossed. Continued heavy grazing, no rest, greatly reduced fire frequency, and increasing shrub canopy cover have altered this plant community drastically. Midgrasses, though still present, are relegated to a position within the thorny shrubs. Interspaces between shrubs are dominantly species of hooded windmillgrass, sand dropseed, curly mesquite, and Hall's panicum. Water, energy, and mineral cycles are drastically altered. Although rainfall still infiltrates within the shrub community, woody plants harvest the water, limiting the amount available for herbaceous production. In this state, fire is rare to non-existent due to decreased fine fuel loads and absence of litter accumulation. In above normal rainfall years, some fine fuel loads may build in the shrub interspaces allowing mid-summer lightning fires to

burn. These wild fires will be of much less intensity, will essentially burn only in the interspaces and will do little to damage shrub canopy. In most instances, prescribed burning is not feasible due to decreased fuel loading and discontinuous fuel beds.

Table 7. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	336	1457	2018
Shrub/Vine	336	448	560
Forb	336	448	560
Tree	–	–	–
Total	1008	2353	3138

Figure 15. Plant community growth curve (percent production by month). TX4801, Mid/Shortgrasses Shrubland Community. Mid and shortgrasses with forbs and 20-50% woody canopy..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	2	10	20	20	5	8	15	10	6	2

Community 2.2 Wooded Grassland



Figure 16. 2.2 Wooded Grassland Community

The Wooded Grassland Community (2.2) has woody canopies exceeding 50, and often 80, percent. Midgrasses are found only within thorny shrubs and the few interspaces are dominated by shortgrasses. Fire on the site is almost totally non-existent. This community can be brought back to Community 2.1 or even the Grassland State (1) but not without significant restoration efforts. Because of the diverse woody plant community, this site in this community is most often manipulated by roller-chopping to enhance it for white-tailed deer, northern bobwhite, and/or scaled quail. This is a very good wildlife site in this state if the woody plant community is manipulated in some pattern and grazing use is very carefully managed. Wildlife enterprises are commonly utilize this community.

Table 8. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	112	560	1009
Shrub/Vine	560	897	1009
Forb	224	392	560
Tree	–	–	–
Total	896	1849	2578

Figure 18. Plant community growth curve (percent production by month). TX4802, Wooded Grassland Community. Wooded Grassland Community with 50 to 80% woody canopy cover..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	5	14	16	16	10	13	15	6	2	1

Pathway 2.1A

Community 2.1 to 2.2

A shift to the to Community 2.2 occurs if brush management is not accomplished. Drought hastens the process. A lack of brush management allows existing brush to gain in stature. Seedlings are introduced through droppings from livestock and wildlife. A reduction in midgrass also corresponds in a reduction of fuel loading needed for fire to effectively suppress woody species, although fire is a questionable at this point.

Pathway 2.2A

Community 2.2 to 2.1

Managerial activities that restore the hydrologic cycle, such as the energy captured by midgrasses, and restored ground cover will tend to move the Community 2.2 toward the Mid/Shortgrass Shrubland Complex (2.1). Selective brush management is needed to accomplish the desired canopy level and spatial arrangement of woody species. Integrated brush management and utilizing historic ecological disturbances such as herbivory and fire in are needed to maintain the desired brush densities. The time to shift back to the 20 to 50 percent canopy is dependent upon favorable growing conditions and could take three to five years.

State 3

Seeded

Dominant plant species

- buffelgrass (*Pennisetum ciliare*), grass

Community 3.1

Introduced and Native Seeding

The Seeded State is attained by mechanical manipulation of the woody plant community either by rhome disking or by root-plowing. Following mechanical control, the site has traditionally been seeded to buffelgrass (Cenchrus ciliaris). Historically, native-grass mixtures have been used very little in South Texas due to the lack of availability. Following mechanical management, woody plants will re-establish. The woody plant community that returns will be limited in species diversity and little will be desirable for. The Seeded State can be maintained for long periods of time with the use of fire, periodic brush management, and the use of prescribed grazing.

Table 9. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1009	2690	3811
Forb	168	224	336
Shrub/Vine	45	78	112
Tree	–	–	–
Total	1222	2992	4259

Figure 20. Plant community growth curve (percent production by month). TX4762, Introduced Grass Community. Planted into introduced grasses for pasture planting..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	10	20	20	5	10	15	10	5	0

Transition T1A State 1 to 2

The Grassland State will cross a threshold to Shrubland (State 2) with abusive grazing and without brush management or fire. Severe drought is also a significant factor to accelerate this crossing of a threshold. In State 2 more rainfall is being utilized by woody plants than the herbaceous plants. Because of the increased canopy, sunlight is being captured by the woody plants and converted to energy instead of the herbaceous plants.

Transition T1B State 1 to 3

The transition to the Converted Land State is triggered by major ground disturbing mechanical treatment and planting to native or introduced forages. Planting is usually done following brush management.

Restoration pathway R2A State 2 to 1

Brush management is the key driver in restoring State 2 back to the Grassland State (1). Reduction in woody canopy below 20 percent will take large energy inputs depending on the canopy cover. A prescribed grazing plan and prescribed burning plan will keep the state functioning.

Transition T2A State 2 to 3

The transition to the Seeded State is triggered by major ground disturbing mechanical treatment and planting to native or introduced forages. Planting is usually done following brush management.

Transition T3A State 3 to 2

The transition from the Seeded State to the Shrubland State is triggered by neglect or no management over long periods of time. Shrubs re-establish from the seed bank and introduction from wildlife and livestock. A complete return to a previous state is not possible if adapted non-native plants have been established.

Additional community tables

Table 10. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
0	Midgrass			179–740	
1	Midgrasses			359–1480	
	multiflower false Rhodes grass	TRPL3	<i>Trichloris pluriflora</i>	224–785	–
	silver beardgrass	BOLAT	<i>Bothriochloa laguroides</i> ssp. <i>torreyana</i>	22–336	–
	large-spike bristlegrass	SEMA5	<i>Setaria macrostachya</i>	34–336	–
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	56–224	–
2	Mid/Shortgrasses			179–740	
	pink pappusgrass	PABI2	<i>Pappophorum bicolor</i>	56–336	–

	nooded windmill grass	CHCU2	<i>Chloris cucullata</i>	34–224	–
	plains lovegrass	ERIN	<i>Eragrostis intermedia</i>	22–112	–
	lovegrass tridens	TRER	<i>Tridens eragrostoides</i>	22–112	–
3	Mid/Shortgrasses			90–370	
	purple threeawn	ARPU9	<i>Aristida purpurea</i>	0–168	–
	marsh bristlegrass	SEPA10	<i>Setaria parviflora</i>	0–112	–
	Texas bristlegrass	SETE6	<i>Setaria texana</i>	11–112	–
	slim tridens	TRMUM	<i>Tridens muticus</i> var. <i>muticus</i>	6–56	–
4	Shortgrasses			90–370	
	curly-mesquite	HIBE	<i>Hilaria belangeri</i>	34–168	–
	Hall's panicgrass	PAHA	<i>Panicum hallii</i>	6–112	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	6–56	–
	fall witchgrass	DICO6	<i>Digitaria cognata</i>	0–56	–
	Madagascar dropseed	SPPY2	<i>Sporobolus pyramidatus</i>	6–11	–
	slender grama	BORE2	<i>Bouteloua repens</i>	6–11	–
	Texas grama	BORI	<i>Bouteloua rigidiseta</i>	1–6	–
	red grama	BOTR2	<i>Bouteloua trifida</i>	1–6	–
	tumble windmill grass	CHVE2	<i>Chloris verticillata</i>	0–6	–
	sand crabgrass	DIAR7	<i>Digitaria arenicola</i>	0–6	–
	southern crabgrass	DICI	<i>Digitaria ciliaris</i>	0–6	–
	knot grass	SEREF	<i>Setaria reverchonii</i> ssp. <i>firmula</i>	0–6	–
	gummy lovegrass	ERCU	<i>Eragrostis curtipedicellata</i>	0–6	–
Forb					
5	Forbs			84–168	
	awnless bushsunflower	SICA7	<i>Simsia calva</i>	11–56	–
	whitemouth dayflower	COER	<i>Commelina erecta</i>	6–34	–
	Riddell's dozedaisy	APRI	<i>Aphanostephus riddellii</i>	6–22	–
	violet wild petunia	RUNU	<i>Ruellia nudiflora</i>	6–11	–
	Gregg's tube tongue	JUPI5	<i>Justicia pilosella</i>	6–11	–
	trailing krameria	KRLA	<i>Krameria lanceolata</i>	0–6	–
6	Forbs			50–101	
	Texas sleepydaisy	XATE	<i>Xanthisma texanum</i>	6–22	–
	yellow puff	NELU2	<i>Neptunia lutea</i>	0–17	–
	wild tantan	DEVI3	<i>Desmanthus virgatus</i>	1–17	–
	hoary milkpea	GACA	<i>Galactia canescens</i>	0–11	–
	littleleaf sensitive-briar	MIMI22	<i>Mimosa microphylla</i>	0–11	–
	prairie clover	DALEA	<i>Dalea</i>	0–11	–
	American snoutbean	RHAM	<i>Rhynchosia americana</i>	0–11	–
	globemallow	SPHAE	<i>Sphaeralcea</i>	1–11	–
7	Forbs			34–67	
	Forb, perennial	2FP	<i>Forb, perennial</i>	11–22	–
	fanpetals	SIDA	<i>Sida</i>	6–17	–
	woody crinklemat	TICA3	<i>Tiquilia canescens</i>	6–11	–
	Forb, annual	2FA	<i>Forb, annual</i>	6–11	–

	Texas crownbeard	VEMI	<i>Verbesina microptera</i>	0–6	–
	bristleleaf pricklyleaf	THTE7	<i>Thymophylla tenuiloba</i>	1–6	–
	Texas Indian mallow	ABFR3	<i>Abutilon fruticosum</i>	1–6	–
	prairie false foxglove	AGHE4	<i>Agalinis heterophylla</i>	0–6	–
	wealeaf bur ragweed	AMCO3	<i>Ambrosia confertiflora</i>	0–6	–
	prairie broomweed	AMDR	<i>Amphiachyris dracunculoides</i>	0–6	–
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	1–6	–
	desert goosefoot	CHPR5	<i>Chenopodium pratericola</i>	0–6	–
	Texas bindweed	COEQ	<i>Convolvulus equitans</i>	0–6	–
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	1–6	–
	slimleaf heliotrope	HETO	<i>Heliotropium torreyi</i>	0–6	–
	cheeseweed mallow	MAPA5	<i>Malva parviflora</i>	0–6	–
	povertyweed	MONOL	<i>Monolepis</i>	0–6	–
	Santa Maria feverfew	PAHY	<i>Parthenium hysterophorus</i>	0–6	–
	cutleaf groundcherry	PHAN5	<i>Physalis angulata</i>	1–6	–
	smartweed leaf-flower	PHPO3	<i>Phyllanthus polygonoides</i>	0–6	–

Shrub/Vine

8	Shrubs/Vines			112–224	
	mouse's eye	BEMY	<i>Bernardia myricifolia</i>	34–56	–
	Texas barometer bush	LEFR3	<i>Leucophyllum frutescens</i>	22–56	–
	spiny hackberry	CEEH	<i>Celtis ehrenbergiana</i>	22–45	–
	guajillo	ACBE	<i>Acacia berlandieri</i>	22–45	–
	catclaw acacia	ACGRW	<i>Acacia greggii</i> var. <i>wrightii</i>	11–22	–
	blackbrush acacia	ACRI	<i>Acacia rigidula</i>	11–22	–
	Texan goatbush	CAERT	<i>Castela erecta</i> ssp. <i>texana</i>	11–22	–
	Texas persimmon	DITE3	<i>Diospyros texana</i>	11–22	–
	Berlandier's wolfberry	LYBE	<i>Lycium berlandieri</i>	11–22	–
	Texas kidneywood	EYTE	<i>Eysenhardtia texana</i>	11–22	–
	pricklypear	OPUNT	<i>Opuntia</i>	11–22	–
	lotebush	ZIOB	<i>Ziziphus obtusifolia</i>	6–22	–
	Texas paloverde	PATE10	<i>Parkinsonia texana</i>	6–17	–
	shrubby blue sage	SABA5	<i>Salvia ballotiflora</i>	6–17	–
	Texas babybonnets	COAX	<i>Coursetia axillaris</i>	6–17	–
	Brazilian bluewood	COHO	<i>Condalia hookeri</i>	6–17	–
	algerita	MATR3	<i>Mahonia trifoliolata</i>	6–11	–
	leatherstem	JADI	<i>Jatropha dioica</i>	6–11	–
	coyotillo	KAHU	<i>Karwinskia humboldtiana</i>	6–11	–
	crown of thorns	KOSP	<i>Koeberlinia spinosa</i>	6–11	–
	West Indian shrubverbena	LAUR2	<i>Lantana urticoides</i>	6–11	–
	catclaw acacia	ACGRG3	<i>Acacia greggii</i> var. <i>greggii</i>	6–11	–
	Christmas cactus	CYLE8	<i>Cylindropuntia leptocaulis</i>	6–11	–
	lime pricklyash	ZAFA	<i>Zanthoxylum fagara</i>	6–11	–
	desert yaupon	SCCU4	<i>Schaefferia cuneifolia</i>	1–6	–
	yucca	YUCCA	<i>Yucca</i>	1–6	–

	littleleaf sumac	RHMI3	<i>Rhus microphylla</i>	1–6	–
	Rio Grande beebrush	ALMA9	<i>Aloysia macrostachya</i>	0–6	–
	Texas swampprivet	FOAN	<i>Forestiera angustifolia</i>	1–6	–
	Texan hogplum	COTE6	<i>Colubrina texensis</i>	0–6	–
	clapweed	EPAN	<i>Ephedra antisyphilitica</i>	1–6	–

Animal community

As a historic tall/midgrass prairie, this site was occupied by bison, antelope, deer, quail, turkey, and dove. This site was also used by many species of grassland songbirds, migratory waterfowl, and coyotes. This site now provides forage for livestock and is still used by quail, dove, migratory waterfowl, grassland birds, coyotes, and deer.

Feral hogs (*Sus scrofa*) can be found on most ecological sites in Texas. Damage caused by feral hogs each year includes, crop damage by rutting up crops, destroyed fences, livestock watering areas, and predation on native wildlife. Feral hogs have few natural predators, thus allowing their population to grow to high numbers.

Wildlife habitat is a complex of many different plant communities and ecological sites across the landscape. Most animals use the landscape differently to find food, shelter, protection, and mates. Working on a conservation plan for the whole property, with a local professional, will help managers make the decisions that allow them to realize their goals for wildlife and livestock.

Grassland State (1): This state provides the maximum amount of forage for livestock such as cattle. It is also utilized by deer, quail and other birds as a source of food. When a site is in the reference plant community phase (1.1) it will also be used by some birds for nesting, if other habitat requirements like thermal and escape cover are near.

Tree/Shrubland (2): This state can be maintained to meet the habitat requirements of cattle and wildlife. Land managers can find a balance that meets their goals and allows them flexibility to manage for livestock and wildlife. Forbs for deer and birds like quail will be more plentiful in this state. There will also be more trees and shrubs to provide thermal and escape cover for birds as well as cover for deer.

Seeded State (3): The quality of wildlife habitat this site will produce is extremely variable and is influenced greatly by the timing of rain events. This state is often manipulated to meet landowner goals. If livestock production is the main goal, it can be converted to pastureland. It can also be planted to a mix of grasses and forbs that will benefit both livestock and wildlife. A mix of forbs in the pasture could attract pollinators, birds and other types of wildlife. Food plots can also be planted to provide extra nutrition for deer.

This rating system provides general guidance as to animal preference for plant species. It also indicates possible competition between kinds of herbivores for various plants. Grazing preference changes from time to time, especially between seasons, and between animal kinds and classes. Grazing preference does not necessarily reflect the ecological status of the plant within the plant community. For wildlife, plant preferences for food and plant suitability for cover are rated. Refer to habitat guides for a more complete description of a species habitat needs.

Hydrological functions

The grassland and the shrubland communities on this site use all the water from rainfall events. Research has shown that the evapotranspiration rate on the grassland and the shrubland is nearly the same. Very little water could be harvested from this site if the woody plant community is replaced by a grass-dominated community.

Recreational uses

White-tailed deer, quail, javelina, and feral hogs are hunted on the site. Bird watching may also be done.

Inventory data references

Information presented was derived from the revised Range Site, literature, limited NRCS clipping data (417s), field

observations, and personal contacts with range-trained personnel.

Other references

Archer, S. 1995. Herbivore mediation of grass-woody plant interactions. *Tropical Grasslands*, 29:218-235.

Archer, S. 1995. Tree-grass dynamics in a *Prosopis*-thornscrub savanna parkland: reconstructing the past and predicting the future. *Ecoscience*, 2:83-99.

De Leon, A. 2003. Itineraries of the De León Expeditions of 1689 and 1690. In *Spanish Exploration in the Southwest, 1542-1706*. Edited by H. E. Bolton. Charles Scribner's Sons, New York, NY.

Dillehay T. 1974. Late quaternary bison population changes on the Southern Plains. *Plains Anthropologist*, 19:180-96.

Duaine, C. L. 1971. *Caverns of Oblivion*. Packrat Press, Oak Harbor, WA.

Everitt, J. H., D. L. Drawe, and R. I. Leonard. 2002. *Trees, Shrubs, and Cacti of South Texas*. Texas Tech University Press, Lubbock, TX.

Everitt, J. H., D. L. Drawe, and R. I. Leonard. 1999. *Field Guide to the Broad-Leaved Herbaceous Plants of South Texas*. Texas Tech University Press. Lubbock, TX.

Frost, C. C. 1998. Presettlement fire frequency regimes of the United States: a first approximation. In *Fire in ecosystem management: shifting the paradigm from suppression to prescription*. Tall Timbers Fire Ecology Conference Proceedings. 20:70-81.

Gilbert, L. E. 1982. An ecosystem perspective on the role of woody vegetation, especially mesquite, in the Tamaulipan biotic region of South Texas. *Proceeding Symposium of the Tamaulipan Biotic Province*, Corpus Christi, TX.

Hanselka, W., R. Lyons, and M. Moseley. 2009. *Grazing Land Stewardship: A Manual for Texas Landowners*. Texas AgriLife Extension Service, College Station, TX.

Hart, C. R., T. Garland, A. C. Barr, B. B. Carpenter, and J.C. Reagor. 2003. *Toxic Plants of Texas: Integrated Management Strategies to Prevent Livestock Losses*. Texas Cooperative Extension Bulletin B-6103 11-03.

Heitschmidt R. K., Stuth J. W., eds. 1991. *Grazing management: an ecological perspective*. Timberline Press, Portland, OR.

Inglis, J. M. 1964. A history of vegetation of the Rio Grande Plains. *Texas Parks and Wildlife Department Bulletin No. 45*, Austin, TX.

Lehman, V. W. 1969. *Forgotten legions: sheep in the Rio Grande Plains of Texas*. Texas Western Press, University of Texas at El Paso, El Paso, TX.

McGinty A., D. N. Ueckert. 2001. The Brush Busters success story. *Rangelands*, 23:3-8.

McLendon T. 1991. Preliminary description of the vegetation of South Texas exclusive of coastal saline zones. *Texas Journal of Science*, 43: 13-32

Norwine, J. 1978. Twentieth-century semiarid climates and climatic fluctuations in Texas and northeastern Mexico. *Journal of Arid Environments*, 1:313-325.

Norwine, J. and R. Bingham. 1986. Frequency and severity of droughts in South Texas: 1900-1983, 1-17. In *Livestock and wildlife management during drought*. Edited by R. D. Brown. Caesar Kleberg Wildlife Research Institute, Kingsville, TX.

Parvin, R. W. 2003. Rio Bravo Resource Conservation and Development. Llanos Mestenos South Texas Heritage Trail. Zapata, TX.

Scifres, C. J. and W. T. Hamilton. 1993. Prescribed burning for brushland management: the South Texas example. Texas A&M Press, College Station, TX.

Scifres C. J., W. T. Hamilton, J. R. Conner, J. M. Inglis, and G. A. Rasmussen. 1985. Integrated Brush Management Systems for South Texas: Development and Implementation. Texas Agricultural Experiment Station, College Station, TX.

Texas Parks and Wildlife Department. 2007. List of White-tailed Deer Browse and Ratings. District 8.

Thurrow, T. L. and J. W. Hester. 1997. How an increase or reduction in juniper cover alters rangeland hydrology. Juniper Symposium Proceedings. Texas A&M University, San Angelo, TX.

Weltz, M. A. and W. H. Blackburn. 1995. Water budget for south Texas rangelands. Journal of Range Management, 48:45-52.

Wright, B. D., R. K. Lyons, J. C. Cathey, and S. Cooper. 2002. White-tailed deer browse preferences for South Texas and the Edwards Plateau. Texas Cooperative Extension Bulletin B-6130.

Contributors

Gary Harris, MSSSL, NRCS, Robstown, Texas

Approval

Bryan Christensen, 9/19/2023

Acknowledgments

Reviewers:

Shanna Dunn, RSS, NRCS, Corpus Christi, Texas

Vivian Garcia, RMS, NRCS, Corpus Christi, Texas

Jason Hohlt, RMS, NRCS, Kingsville, Texas

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Vivian Garcia, Zone RMS, NRCS, Corpus Christi, Texas
Contact for lead author	361-241-0609
Date	09/17/2007
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

-
2. **Presence of water flow patterns:** None except following extremely high intensity storms when short flow patterns may appear.
-
3. **Number and height of erosional pedestals or terracettes:** Few.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 0 to 5 percent bare ground. Small and non-connected areas.
-
5. **Number of gullies and erosion associated with gullies:** Few.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Minimal and short.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability class anticipated to be 5 to 6 at the surface.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A-horizon is 0 to 2 inches thick with light, brownish-gray fine sandy loam. Structure is weak, fine, subangular blocky. The surface is hard, friable, slightly sticky, and slightly plastic with few snail shell fragments. The soil is strongly effervescent, moderately alkaline, and has an abrupt smooth boundary. SOM is 0 to 3 percent.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** High canopy, basal cover and density with small interspaces should make rainfall impact negligible. This site has well drained soils, deep with 0 to 3 percent slopes which allow for negligible runoff and erosion.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm-season midgrasses >>
- Sub-dominant: Warm-season shortgrasses >

Other: Forbs > Trees/Shrubs

Additional: Forbs make up five percent species composition while shrubs and trees compose up to five percent species composition.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Grasses due to their growth habit will exhibit some mortality and decadence, though very slight.
-

14. **Average percent litter cover (%) and depth (in):** Litter is dominantly herbaceous.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 1,050 to 3,800 air-dry pounds per acre.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Mesquite, blackbrush, guajillo, lime pricklyash, and cenizo are the primary invaders.
-

17. **Perennial plant reproductive capability:** All species should be capable of plant reproduction, except during periods of prolonged drought, heavy natural herbivory, and/or wild fires.
-