

Ecological site R084BY172TX Sandy 29-33" PZ

Last updated: 9/21/2023
Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 084B–West Cross Timbers

MLRA 84B is characterized by nearly level to strongly sloping, dissected plains with narrow valleys that deepen eastward. Soils are generally deep and formed in sediments of Cretaceous age. Average annual precipitation is 25 to 35 inches, and elevation ranges from 1000 to 1300 feet.

Classification relationships

This ecological site is correlated to soil components at the Major Land Resource Area (MLRA) level which is further described in USDA Ag Handbook 296.

Ecological site concept

These sites occur on deep sandy soils on uplands. The reference vegetation includes native tallgrasses with scattered oaks. Many forbs can also be found along with scattered native shrubs. Without fire or other brush management, woody species may increase and dominate the site.

Associated sites

R084BY169TX	Deep Sand 29-33" PZ Very deep sandy soils on uplands.
R084BY171TX	Loamy Sand 29-33" PZ Loamy sands on uplands
R084BY168TX	Claypan 29-33" PZ Dense clay subsoils.
R084BY173TX	Sandy Bottomland 29-33" PZ Sandy soils on floodplains

Similar sites

R084BY169TX	Deep Sand 29-33" PZ Very deep sandy soils on uplands.
R084BY171TX	Loamy Sand 29-33" PZ Loamy sands on uplands

Table 1. Dominant plant species

Tree	(1) <i>Quercus stellata</i> (2) <i>Quercus marilandica</i>
Shrub	Not specified
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Andropogon gerardii</i>

Physiographic features

This site occurs on interfluves and side slopes of hillslopes and on treads of stream terraces in the West Cross Timbers. Slopes are typically less than 8 percent.

Table 2. Representative physiographic features

Landforms	(1) Hills > Hillslope (2) Hills > Ridge (3) Alluvial plain > Stream terrace
Runoff class	Negligible to medium
Elevation	183–640 m
Slope	0–8%
Water table depth	51 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate is subtropical. Precipitation varies from an average of 33 inches in the eastern part of the Cross Timbers to 29 inches in the western part. Winters are dry and summers are hot and humid. Tropical maritime air masses control the weather during the spring , summer and fall. Large variations in temperature sometimes accompany polar air masses in winter.

Table 3. Representative climatic features

Frost-free period (characteristic range)	193-200 days
Freeze-free period (characteristic range)	220-225 days
Precipitation total (characteristic range)	787-838 mm
Frost-free period (actual range)	192-204 days

Freeze-free period (actual range)	216-226 days
Precipitation total (actual range)	711-864 mm
Frost-free period (average)	197 days
Freeze-free period (average)	222 days
Precipitation total (average)	813 mm

Climate stations used

- (1) MINERAL WELLS AP [USW00093985], Millsap, TX
- (2) BRIDGEPORT [USC00411063], Bridgeport, TX
- (3) PUTNAM [USC00417327], Baird, TX
- (4) RISING STAR 1S [USC00417633], Rising Star, TX
- (5) PROCTOR RSVR [USC00417300], Comanche, TX

Influencing water features

These sites may receive some runoff from adjacent areas and also shed water to areas lower on the landscape. The presence of deep rooted tallgrasses can help facilitate infiltration of water into the soil profile. These sites are not associated with wetlands.

Wetland description

NA

Figure 7-1 The hydrologic cycle with factors that affect hydrologic processes




Figure 8.

Soil features

Representative soil components for this ecological site include: Demona, Heaton, and Nimrod

The site is characterized by deep soils with a high infiltration rate.

Table 4. Representative soil features

Parent material	(1) Alluvium–sandstone and siltstone (2) Residuum–sandstone and siltstone
Surface texture	(1) Loamy fine sand (2) Fine sand
Drainage class	Moderately well drained to well drained
Permeability class	Slow to moderate

Soil depth	102–183 cm
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0–5%
Available water capacity (0-101.6cm)	12.7–25.4 cm
Calcium carbonate equivalent (0-101.6cm)	0–5%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	5.1–7.8
Subsurface fragment volume <=3" (Depth not specified)	0–5%
Subsurface fragment volume >3" (Depth not specified)	0–5%

Ecological dynamics

The reference plant community for the Sandy ecological site is a post oak and blackjack oak savanna with tall and midgrass understory. The grasses are primarily little bluestem (*Schizachyrium scoparium*), Indiangrass (*Sorghastrum nutans*), big bluestem (*Andropogon gerardii*) and sand lovegrass (*Eragrostis trichodes*). Little bluestem and Indiangrass are the most commonly occurring grass species. Smaller amounts of big bluestem and sand lovegrass occur as well. Shrubs and trees are present and consist of post oak (*Quercus stellata*), blackjack oak (*Quercus marilandica*), plums (*Prunus* spp), hawthorns (*Crataegus* spp), greenbriar (*Smilax* spp), grapes (*Vitis* spp) coralberry (*Symphoricarpos orbiculatus*), hackberry (*Celtis reticulata*) and bumelia (*Sideroxylon lanuginose*). Woody plants have increased on virtually all of the sandy sites over the past 100 to 150 years. The production potential of the site is moderate. Pre-settlement grazers included bison and deer. The grasses are palatable and the site provides year round grazing. The most limiting soil factor is erodibility followed closely by fertility. In very dry periods, the soils can appear rather droughty. When good rainfall is received, the site produces fairly well.

Fire played a role in the ecology of the site as is true for most of the grasslands. The main effect of fire on this site was to hold woody shrubs and cactus in check. The grass species such as little bluestem and Indiangrass are considered fire neutral as far as their response to fire. Climate and soils are the most important and limiting factors affecting grass vegetation on the site. Fire stimulated forbs growth if the timing was right and the fires of pre-settlement days were probably more severe due to more fuel being available which could have been more damaging to woody plants. Fire usually creates more diversity in this site for a year or two post-burn. Prescribed fire is sometimes used as a tool to promote diversity, mainly for wildlife. Fire will usually not produce much mortality in older woody plants. After brush has been controlled with herbicides or mechanically, fire can sometimes be used effectively to suppress regrowth. Small juniper can be killed by fire. Fuel loads are often the most limiting factor for the effective use of prescribed fire on this site. In general, the uses of fire on mature (larger) or dense stands of woody plants do not result in the same positive effects that burning has in tall/midgrass communities.

With abusive grazing practices, the vigorous Indiangrass and big bluestem will become lower in vigor while little bluestem will increase. Secondary successional species such as sand dropseed (*Sporobolus cryptandrus*), and silver bluestem (*Bothriochloa laguroides*) then will begin to increase along with an increase of woody plants. The little bluestem is a tough, resistant species tolerant of some fairly heavy grazing for long periods, but at some point, a threshold is crossed and the ground cover is opened up resulting in bare places where weedy species can establish. Western ragweed (*Ambrosia psilostachya*), crotons (*Croton setigerus*), and cool-season annuals will quickly invade if the principal species are in a weakened condition. Grazing management probably has minimal effect on the proliferation of woody plants, but a good cover of perennial grasses likely minimizes the seed to soil contact the woody plants need to establish. Prescribed fire where it can be safely carried out provides a much better method to control the spread of woody plants. Selective individual removal of woody plants is easy and economical

when a few plants begin to show up on the site, but the increase of number of plants can be fairly rapid and the number of woody plants per acre will soon become too numerous for individual control to be feasible. Prescribed grazing with a reasonable stocking rate can sustain the grass species composition and production at a near reference level until the brush canopy is so dense that the shade starts to interfere with photosynthesis. The sandy site can be abused to the point that the perennial warm-season grasses thin out and lower successional grasses along with annual forbs begin to dominate. This process of degradation usually takes many years and is further exacerbated by summer drought and above average winter moisture.

Long-term droughts that occur only three to four times in a century can effect some change in plant communities. Short-term droughts are common and usually do not have a lasting effect in changing stable plant communities, although production will be affected. When a brush canopy becomes established which shades the ground sufficiently it tends to favor cool season annual species. Once a state of brush and cool-season annuals is reached, recovery to a good perennial grass cover is unlikely without major input with brush management and reseeding. In summary, the change in states of vegetation depend on the type of grazing management applied over many years, and the rate of invasion and establishment of woody species. The effects of seasonal moisture and short term dry spells become more pronounced after the site crosses thresholds to a lower ecological condition. Plant communities that consist of warm-season perennial grasses such as little bluestem are able to persist and withstand climatic extremes with only minor shifts in the overall plant community.

This site was inhabited by grassland wildlife species such as bison, grassland birds and small mammals. Over the years, as the site has changed to a more mixed grass and shrub community, more wildlife species have come to utilize it for habitat. Woody plants provide cover for white-tailed deer and bob-white quail. These wildlife species have both increased along with the brushy plants due to the cover that these plants provide. More forbs are needed to meet these species food requirements and woody plants for browse are important for deer. It is often the objective of many land owners to strike a balance in plant community so that these wildlife species can exist along with domestic livestock. This can be accomplished by a carefully thought out grazing and brush management program. It must be realized that managing at a lower successional level may meet some wildlife species requirements very well, but may not be nearly as productive for grazing purposes, and may not be as capable of satisfying functions such as nutrient cycling, hydrologic protection, plant community stability or soil protection. A proper balance can be achieved with careful planning that considers all resources.

Hydrologically, the site contributes runoff to the various draws, creeks, and streams that are common in the MLRA. If the perennial grass cover is maintained in good vigor, then maximum infiltration occurs and runoff is reduced. More water getting into the ground means a healthier, more productive plant community. If infiltration is minimal, then the effect is an artificially shallow soil with plant roots retreating to near the soil surface. More perennial grass cover means less runoff may result; but the runoff that does occur is less laden with sediment. Overall watershed protection is enhanced by a healthy grassland community, as is nutrient cycling.


State and Transitional Pathways: Narrative

The following diagram suggests some pathways that the vegetation on this site might take in response to various treatment or natural stimuli over time. There may be other states that are not shown on this diagram. This information is to show that changes in plant community do occur due to management and natural factors; and can be changed by implementing certain practices. The plant communities described are commonly observed on this site. Before making plans for plant community manipulation for specific purposes, consult local professionals.

As a site changes in plant community makeup, the changes may be due many factors. Change may occur slowly or in some cases, fairly rapidly. As vegetative changes occur, certain thresholds are crossed. This means that once a certain point is reached during the transition of one community to another, a return to the first state may not be possible without the input of some form of energy. This often means intervention with practices that are not part of natural processes. An example might be the application of herbicide to control some woody species in order to reduce its population and encourage more grass and forbs growth. Merely adjusting grazing practices would probably not accomplish any significant change in plant community once certain thresholds are crossed. The amount of energy required to effect change in community would depend on the present vegetative state and the desired change.

State and transition model

Sandy 29-33" PZ
R084BY172TX



Legend

- 1.1A Heavy Continuous Grazing, No Fire, Idle
- 1.2A Prescribed Grazing, Brush Management, Prescribed Burning
- T1A Heavy Continuous Grazing, No Fire, Idle
- R2A Prescribed Grazing, Brush Management, Range Planting, Prescribed Burning
- T1B Pasture Planting, Crop Cultivation, Pest Mgmt, Nutrient Mgmt, Prescribed Grazing
- R3A Prescribed Grazing, Range Planting, Prescribed Burning, Brush Management, Abandonment
- 2.1A Heavy Continuous Grazing, No Fire, Idle
- T2A Pasture Planting, Crop Cultivation, Pest Management, Nutrient Management
- T3B Heavy Continuous Grazing, No Brush Management, No Fire, Abandonment
- 3.1A Heavy Continuous Grazing, No Brush Management, Idle
- T3A Idle, No Fire
- T4A Heavy Continuous Grazing, No Brush Management, No Fire

State 1
Tallgrass/Oak Savannah State - Reference

The interpretive plant community for this site is a Tallgrass/Oak Savanna. The community is dominated by warm-season perennial tallgrasses with blackjack oak and post oak. The major perennial grass species are well dispersed through the community. Perennial forbs and shrubs are well represented throughout the community. Annual production ranges from 3345 to 4755 pounds per acre. Grasses make up 75% of species composition while trees/shrubs make up 17.5% species composition. The Tallgrass/Shrubland Transition Community occurs when post oak, blackjack oak and elm starts regeneration and the tallgrasses will start to disappear from the plant community. Invader brush appears and becomes established. Bumelia and hackberry starts to increase. Texas wintergrass and purpletop tridens increases as brush canopy increases. The plant community consists of about a 15 percent canopy of mature trees with an understory canopy of shrubs and young oaks, elm and pecan encompassing 5 to 35%. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 65% species composition while trees/shrubs make up 25% species composition.

Dominant plant species

- post oak (*Quercus stellata*), tree
- blackjack oak (*Quercus marilandica*), tree
- little bluestem (*Schizachyrium scoparium*), grass

Community 1.1
Tallgrass/Oak Savannah Community



Figure 9. 1.1 Tallgrass/Oak Savannah Community

The interpretive plant community for this site is a Tallgrass/Oak Savanna. The community is dominated by warm-season perennial tallgrasses with blackjack oak and post oak. The major perennial grass species are well dispersed through the community. Perennial forbs and shrubs are well represented throughout the community. This plant community evolved with a short duration of heavy use by large herbivores followed by long rest periods due to herd migration along with occasional fire. Annual production ranges from 3345 to 4755 pounds per acre. Grasses make up 75% of species composition while trees/shrubs make up 17.5% species composition.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	2942	3407	3867
Tree	392	633	874
Forb	258	325	392
Shrub/Vine	157	174	196
Total	3749	4539	5329

Figure 11. Plant community growth curve (percent production by month). TX5514, Post oak/Blackjack Oak Savannah. Post oak and blackjack oak savannah with tall grasses..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2	2	8	10	20	23	5	3	10	10	5	2

Community 1.2

Tallgrass/Shrubland Transition Community



Figure 12. 1.2 Tallgrass/Shrubland Transition Community

This transition state occurs without fire or brush management. Post oak, blackjack oak and elm starts regeneration and the tallgrasses will start to disappear from the plant community. Invader brush appears and becomes established. Bumelia and hackberry starts to increase. Texas wintergrass and purpletop tridens increases as brush canopy increases. The plant community consists of about a 15 percent canopy of mature trees with an understory canopy of shrubs and young oaks, elm and pecan encompassing 5 to 35%. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 65% species composition while trees/shrubs make up 25% species composition. This transition state can revert back to the tallgrass savannah with prescribed burning and/or prescribed grazing. Without prescribed burning and/or prescribed grazing this transition state will continue to shift toward the Oak/Shrubland Transition State (2.1).

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	2589	2998	3403
Tree	588	684	773
Shrub/Vine	392	454	516
Forb	353	409	464
Total	3922	4545	5156

Figure 14. Plant community growth curve (percent production by month). TX5511, Shrubland Transition. Increasing percentage of shrubs invading site (20-50 % canopy).

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	1	3	10	30	25	8	5	10	5	2	1

Pathway 1.1A

Community 1.1 to 1.2



Tallgrass/Oak Savannah Community



Tallgrass/Shrubland Transition Community

With heavy continuous grazing, no fire, and land in idle, the Tallgrass/Oak Savannah Community will shift to the Tallgrass/Shrubland Transition Community.

Pathway 1.2A Community 1.2 to 1.1



Tallgrass/Shrubland Transition Community



Tallgrass/Oak Savannah Community

The Tallgrass/Shrubland Transition Community can shift back to the Tallgrass/Oak Savannah Community with the use of Prescribed Grazing, Brush Management, and Prescribed Burning conservation practices.

Conservation practices

Brush Management
Prescribed Burning
Prescribed Grazing

State 2 Oak Shrubland State

The Oak/Shrubland Transition Community (2.1) consists of 50 to 80% canopy of woody plants. As the shrubland community ages, oak matures and the invader species increase. Warm-season perennial tallgrasses such as Indiangrass and switchgrass have all but disappeared. In the early stages of this transition stage, little bluestem tends to dominate the grasses. However, as brush canopy continues to increase, Texas wintergrass, three-awns and annuals continue to increase while the little bluestem decreases. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 5% species composition while trees/shrubs make up 90% species composition. The Oak/Woodland Community is a closed overstory (greater than 80% canopy) woodland dominated by post oak and blackjack oak. Understory shrubs, sub-shrubs, and woody vines are also present. The herbaceous understory is almost nonexistent except for shade-tolerant species occurring in small amounts. Small isolated clearings will contain little bluestem, perennial three-awns, Texas wintergrass and small amounts of other grasses. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 5% species composition while trees/shrubs make up 90% species composition.

Dominant plant species

- post oak (*Quercus stellata*), tree
- elm (*Ulmus*), tree
- Ashe's juniper (*Juniperus ashei*), tree
- Texas wintergrass (*Nassella leucotricha*), grass

Community 2.1 Oak/Shrubland Transition Community



Figure 15. 2.1 Oak/Shrubland Transition Community

The Oak/Shrubland Transition Community (2.1) consists of 50 to 80% canopy of woody plants. As the shrubland community ages, oak matures and the invader species increase. Warm-season perennial tallgrasses such as Indiangrass and switchgrass have all but disappeared. In the early stages of this transition stage little bluestem tends to dominate the grasses: however as brush canopy continues to increase, Texas wintergrass, three-awns (*Aristida* spp) and annuals continue to increase while the little bluestem decreases. Where severe erosion has occurred or cropland has been abandoned mesquite and/or juniper may dominate the site. Continuous grazing by domestic livestock has accelerated the shift. The shift to this state has occurred due to the absence of fire or other means of brush suppression. Where this state has been reached from cropland or pasture, mesquite and/or juniper dominate the woody vegetation, but post oak and blackjack oak are beginning to occur. The grass species that dominate the site are splitbeard bluestem (*Andropogon ternarius*), silver bluestem (*Bothriochloa laguroides*), Texas wintergrass and three-awns along with the seeded introduced grass species. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 5% species composition while trees/shrubs make up 90% species composition. This state can be reverted back to near reference condition by some means of brush suppression and good grazing management. Without this treatment the site will continue to shift toward the Oak Woodland State.

Table 7. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Tree	1132	1314	1495
Forb	1098	1271	1444
Grass/Grasslike	1098	1271	1444
Shrub/Vine	594	684	773
Total	3922	4540	5156

Figure 17. Plant community growth curve (percent production by month). TX5512, Oak Shrubland Transition. Continued increase of invader species and post oaks maturity. Approximately 50-80 percent canopy cover..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	3	10	30	30	5	5	8	5	1	1

Community 2.2

Oak/Woodland Community



Figure 18. 2.2 Oak/Woodland Community

This plant community is a closed overstory (greater than 80% canopy) woodland dominated by post oak and blackjack oak. Other species present in small amounts are cedar elm, hackberry, bois d'arc (*Maclura pomifera*). Understory shrubs and sub-shrubs include yaupon, hawthorns, and American beautyberry (*Callicarpa americana*). Woody vines are also present and include greenbriars, poison oak (*Toxicodendron* spp), Virginia creeper (*Parthenocissus quinquefolia*) and grapes. The herbaceous understory is almost nonexistent. Shade tolerant species such as purpletop tridens (*Tridens flavus*) and Canada wildrye (*Elymus canadensis*) occur in small amounts. Small isolated clearings will contain little bluestem, perennial three-awns, Texas wintergrass and small amounts of other grasses. Continuous grazing by domestic livestock has accelerated the shift. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 5% species composition while trees/shrubs make up 90% species composition. This Oak/Woodland Community (2.2) has developed due to the absence of fire (or some other method of brush suppression). Livestock grazing yearlong accelerates the shift. The tallgrass savannah (1.1) can be restored by prescribed burning but will require many years of burning due to light fuel load of fine fuel and the absence of a seed source for the tall grasses. Chemical control alone is usually not a good option for treatment on a large scale due to the resistance of some of the woody plant species to herbicides. Mechanical treatment of this site along with seeding is generally the best method for conversion back to a tallgrass savannah. The cost of doing this type of treatment is usually so expensive as to be not economically feasible so in most instances it is planted to introduced grass species and converted to pastureland. The soils of this site are very sensitive to erosion. During the transition states from the Historic State due to the reduction of grass litter, sheet and rill erosion and in some instances wind erosion has accelerated and by the time it has reached the Woodland State the topsoil depth is greatly reduced. This in turn has reduced the soils natural fertility. While in the transition stages the organic matter in the soil is also reduced and may never be what was present in the tallgrass prairie. In the Woodland Community (2.2) the amount of litter cover is similar to Plant Community (1.1) but this cover is now mostly leaves. The leaves of the trees and underbrush intercept rainfall from lighter intensity rainfall which evaporates before reaching the ground resulting in less water reaching the soil surface. When runoff does occur there are more tendencies for the litter to drift until it catches on the stems of dense underbrush or what little grass is present. When the woodland is grazed the amount of litter decreases along with a decrease of surface vegetation which increases the drifting of the litter with runoff. Due to the presence of shade the amount of grass cover is greatly reduced which in turn reduces forage production from the reference state. When this state is reached following abandonment of cropland or pastureland of introduced plants quite often severe erosion has occurred. The absence of topsoil has greatly decreased water infiltration which in turn increases runoff.

Table 8. Annual production by plant type


Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Tree	1950	2242	2578
Shrub/Vine	1569	1816	2062
Forb	247	280	308
Grass/Grasslike	157	179	207
Total	3923	4517	5155

Figure 20. Plant community growth curve (percent production by month).
TX5513, Postoak Woodland. Postoak dominated closed overstory with over 80 percent canopy cover..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	1	2	10	50	14	5	5	5	5	1	1

Pathway 2.1A

Community 2.1 to 2.2



Oak/Shrubland Transition Community



Oak/Woodland Community

The Oak/Shrubland Transition Community shifts to the Oak/Woodland Community when heavy continuous grazing, no fires, and land in idle occur.

State 3

Converted Land State

Conversion of the Tallgrass/Oak Savannah (1.1) to Converted Land Community (3.1) (mainly for cotton production) occurred from first settlement by European settlers during the middle 1800's and continued until early 1900's. Some remains in cropland today. This site is often planted to introduced grasses following crop production or brush control. Typical species planted include bermudagrass varieties and yellow bluestems. Many of these species may become invasive and once established they are difficult to remove and hinders the establishment of native species. In the Threeawn/Invasive Brush Transition Community, the plant community will move toward a transition of invasive brush species such as mesquite and/or cacti without the annual application of the cultural practices, especially fertilization. The introduced grass species will start to disappear and be replaced by three-awns and annuals. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 40% species composition while trees/shrubs make up 25% species composition.

Dominant plant species

- Bermudagrass (*Cynodon dactylon*), grass

Community 3.1

Converted Land Community

Conversion of the Tallgrass/Oak Savannah to cropland occurred from first settlement by European settlers during the middle 1800's and continued until early 1900's. Some remains in cropland today. The early cropping with little regard for erosion control leads to severe erosion by water and wind. Erosion changes fertility, soil structure and moisture holding capacity of the soil. While restoration of this site to some semblance of the tallgrass savannah is possible with seeding, prescribed grazing and prescribed burning; a complete restoration of the reference plant community in a reasonable time is very unlikely. In recent years if cropping is abandoned the land is usually planted to introduced grass and manages as pastureland. Refer to cropland capability classes for production potentials and limitations. This site is often planted to introduced grasses following crop production or brush control. These grasses are planted mostly for livestock grazing and some hay production. Typical species planted include Bermudagrass varieties and yellow bluestems. Many of these species may become invasive and once established they are difficult to remove and hinders the establishment of native species. The establishment and maintenance of these species requires fertilization, weed control and prescribed grazing management. Without the annual application of these cultural practices the plant community will move toward a transition of invasive brush species. Refer to pastureland suitability groups for species suitability, production potentials and limitations.

Figure 21. Plant community growth curve (percent production by month).
TX5520, Pastureland. Coastal Bermudagrass or Introduced Species..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	21	22	10	5	14	10	8	0

Community 3.2

Threeawn/Invasive Brush Transition Community



Figure 22. 3.2 Three-awn/Invasive Brush Transition Community

Without the annual application of the cultural practices, especially fertilization, the plant community will move toward a transition of invasive brush species such as mesquite and/or cacti. The introduced grass species will start to disappear and be replaced by threeawns and annuals. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 40% species composition while trees/shrubs make up 25% species composition.

Table 9. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1485	1726	1961
Forb	1457	1681	1905
Tree	588	678	773
Shrub/Vine	392	454	516
Total	3922	4539	5155

Figure 24. Plant community growth curve (percent production by month). TX5517, Threeawn/Invasive Shrubs Community. Threeawns and Invasive Shrub dominant community. Converted from old cropland into threeawn shrub community..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2	2	8	10	20	23	5	3	10	10	5	2

Pathway 3.1A

Community 3.1 to 3.2

The Converted Land Community will shift to the Three-awn/Invasive Brush Transition Community with the use of heavy continuous grazing, no brush management, and land being abandoned.

State 4

Invasive Brush State

The Invasive Brush/Midgrass Community occurs due to continued absence of added fertility the plant community will continue change toward a brushy condition. The shrubs will continue to thicken especially mesquite. At this state the introduced grasses have almost disappeared. Annual production ranges from 3500 to 4600 pounds per

acre. Grasses make up 30% species composition while trees and shrubs make up 43% species composition.

Dominant plant species

- Ashe's juniper (*Juniperus ashei*), tree
- Bermudagrass (*Cynodon dactylon*), grass

Community 4.1
Invasive Brush/Midgrass Transition Community



Figure 25. 4.1 Invasive Brush/Midgrass Community

With continued absence of added fertility the plant community will continue change toward a brushy condition. The shrubs will continue to thicken especially mesquite. The grasses in this community will change toward more perennials, where the fertility is especially low splitbeard bluestem appears and may dominate the grasses. At this state the introduced grasses have almost disappeared. Annual production ranges from 3500 to 4600 pounds per acre. Grasses make up 30% species composition while trees and shrubs make up 43% species composition.

Table 10. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1233	1412	1592
Forb	1009	1171	1334
Tree	897	1042	1188
Shrub/Vine	785	913	1042
Total	3924	4538	5156

Figure 27. Plant community growth curve (percent production by month).
TX5523, Midgrass & Invasive Brush Community. old cropland field
dominated by plant community consisting of midgrasses and invasive
brush..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2	2	8	10	20	23	5	3	10	10	5	2

Transition T1A
State 1 to 2

The Tallgrass/Oak Savannah State will transition into the Oak Shrubland State due to continuous heavy grazing, no fires, and land not in use.

Transition T1B
State 1 to 3

The Tallgrass/Oak Savannah State will transition into the Converted Land State with the application of various conservation practices including Pasture/Hay Planting, Crop Cultivation, Pest Management, Nutrient Management, and Prescribed Grazing.

Restoration pathway R2A State 2 to 1

With the use of Prescribed Grazing, Brush Management, Range Planting, and Prescribed Burning conservation practices, the Oak Shrubland State can be restored into the Tallgrass/Oak Savannah State.

Conservation practices

Brush Management
Prescribed Burning
Prescribed Grazing
Range Planting

Transition T2A State 2 to 3

The Oak Shrubland State will transition into the Converted Land State with the use of various conservation practices including Pasture/Hay Planting, Crop Cultivation, Pest Management, and Nutrient Management.

Restoration pathway R3A State 3 to 1

With the application of various conservation practices including Prescribed Grazing, Range Planting, Prescribed Burning, and Brush Management, the Converted Land State can be restored to the Tallgrass/Oak Savannah State.

Transition T3B State 3 to 2

The Converted Land State may transition into the Oak Shrubland State through the continued use of heavy continuous grazing, no brush management, no fires, land abandonment, no pest and nutrient management.

Transition T3A State 3 to 4

The Converted Land State will transition into the Invasive Brush State due to land being abandoned from land conversion such as cropland and pastureland as well as no fires.

Transition T4A State 4 to 2

The Invasive Brush State is transitioned to the Oak Shrubland State through heavy continuous grazing, no brush management, and no fires.

Additional community tables

Table 11. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Tallgrass			981–1289	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	981–1289	–

2	Tallgrass			392–516	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	392–516	–
3	Tallgrass			392–516	
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	392–516	–
4	Midgrass			196–258	
	silver beardgrass	BOLAT	<i>Bothriochloa laguroides</i> ssp. <i>torreyana</i>	196–258	–
5	Cool Season			196–258	
	Scribner's rosette grass	DIOLS	<i>Dichanthelium oligosanthes</i> var. <i>scribnerianum</i>	196–258	–
6	Midgrass			196–258	
	purpletop tridens	TRFL2	<i>Tridens flavus</i>	196–258	–
7	Midgrass			196–258	
	sand lovegrass	ERTR3	<i>Eragrostis trichodes</i>	196–258	–
8	Midgrass			196–258	
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	196–258	–
9	Midgrass			196–258	
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	196–258	–
10	Mid/Shortgrasses			0–1	
	fall witchgrass	DICO6	<i>Digitaria cognata</i>	0–1	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	0–1	–
	Virginia wildrye	ELVI3	<i>Elymus virginicus</i>	0–1	–
	crowngrass	PASPA2	<i>Paspalum</i>	0–1	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–1	–
Forb					
11	Forbs			196–258	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	0–67	–
	whitemouth dayflower	COER	<i>Commelina erecta</i>	0–67	–
	prairie clover	DALEA	<i>Dalea</i>	0–67	–
	purple prairie clover	DAPU5	<i>Dalea purpurea</i>	0–67	–
	lespedeza	LESPE	<i>Lespedeza</i>	0–67	–
	yellow puff	NELU2	<i>Neptunia lutea</i>	0–67	–
	amberique-bean	STHE9	<i>Strophostyles helvola</i>	0–67	–
Shrub/Vine					
12	Shrubs/Vines			118–157	
	fragrant sumac	RHAR4	<i>Rhus aromatica</i>	0–39	–
	blackberry	RUBUS	<i>Rubus</i>	0–39	–
	roundleaf greenbrier	SMRO	<i>Smilax rotundifolia</i>	0–39	–
	coralberry	SYOR	<i>Symphoricarpos orbiculatus</i>	0–39	–
	grape	VITIS	<i>Vitis</i>	0–39	–
Tree					
13	Tree			588–773	
	blackjack oak	QUMA3	<i>Quercus marilandica</i>	0–773	–
	post oak	QUST	<i>Quercus stellata</i>	0–773	–
14	Tree			78–104	

14	Tree			10-101	
	hackberry	CELT1	<i>Celtis</i>	0-101	—
	hawthorn	CRATA	<i>Crataegus</i>	0-101	—
	plum	PRUNU	<i>Prunus</i>	0-101	—
	bully	SIDER2	<i>Sideroxylon</i>	0-101	—
	Hercules' club	ZACL	<i>Zanthoxylum clava-herculis</i>	0-101	—

Animal community

The historic oak-tallgrass savannah was habitat to migratory bison herds, deer, turkey, migratory birds and large predators such as wolves, coyotes, mountain lions and black bear. White-tail deer, turkey, bobcats and coyotes along with resident and migratory birds and small mammals find suitable habitat today. Domestic livestock and white-tail deer are the dominant grazers and browsers of the site. As the savannah changes through the various vegetative states towards the Oak Woodland, the quality of the habitat may improve for some species and decline for others. Management must be applied to maintain a vegetative state in optimum habitat quality for the desired animal species.

Hydrological functions

Peak rainfall periods occur in April, May, June, September and October. Rainfall amounts may be high (3 to 10 inches per event) and events may be intense. The soil of this site are very susceptible to erosion and severe erosion occurs where adequate herbaceous cover is not maintained and on heavy use areas such as roads and livestock trails. Periods of 60 plus days of little or no rainfall during the growing season are common. The hydrology of this site may be manipulated with management to yield higher runoff volumes or greater infiltration to groundwater. Management for less herbaceous cover will favor higher surface runoff while dense herbaceous cover favors ground water recharge. Potential movement of soil (erosion), pesticides and both organic and inorganic nutrients(fertilizer) should always be considered when managing for higher volumes of surface runoff.

Recreational uses

Hunting, hiking, camping, equestrian, bird watching and off road vehicle use.

Wood products

Oaks and mesquite are used for firewood. Mesquite is also used for barbecue wood.

Other products

None.

Other information

None.

Inventory data references

Data from the ESI Database were obtain from the following counties: Erath, Hamilton, Hood, Montague, Parker, Somervell, and Wise.

References

. 2021 (Date accessed). USDA PLANTS Database. <http://plants.usda.gov>.

Other references

White-tailed Deer, Their Foods and Management in the Cross Timbers
By Kenneth L. Gee, Michael D Porter, Steve Demarais, Fred C. Bryant, and Gary Van Vreede. A Samuel Roberts Noble Foundation Publication, 1991

Technical Review:

Homer Sanchez, State Rangeland Management Specialist, NRCS, Temple, Texas
Mark Moseley, State Rangeland Management Specialist, NRCS, Stillwater, Oklahoma
Kent Ferguson, Zone Rangeland Management Specialist, NRCS, Weatherford, Texas
Dr. Jack Eckroat, Grazing Lands Specialist, NRCS, Stillwater, Oklahoma
Justin Clary, Rangeland Management Specialist, NRCS, Temple, Texas

Contributors

Earl V. Hogan
PES edits by Colin Walden, Stillwater Soil Survey Office

Approval

Bryan Christensen, 9/21/2023

Acknowledgments

Site Development and Testing Plan:

Future work, as described in a Project Plan, to validate the information in this Provisional Ecological Site Description is needed. This will include field activities to collect low, medium and high intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document. Annual reviews of the Project Plan are to be conducted by the Ecological Site Technical Team.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Colin Walden, Range Management Specialist, Soil Survey Region 9.
Contact for lead author	colin.walden@ok.usda.gov
Date	01/25/2018
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Minimal evidence of current or past rill formation.

-
2. **Presence of water flow patterns:** Few water flow patterns on steep areas. Short and stable, not incising.

-
3. **Number and height of erosional pedestals or terracettes:** No pedestals terracettes present.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground less than 10 percent. Bare areas small and not connected.
-
5. **Number of gullies and erosion associated with gullies:** No gullies present.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** No wind scoured areas.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Litter movement less than 3 feet. Vegetative cover should restrict litter movement over long distances. Only herbaceous litter less than .25 inches expected to move.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability scores of 5 or greater expected.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** A--0 to 4 inches; grayish brown (10YR 5/2) fine sand, dark grayish brown (10YR 4/2) moist; single grained;
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Presence of perennial tall and midgrasses help to facilitate percolation into the soil. Some runoff expected on steeper slopes during moderate precipitation events.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction under reference conditions. Beware texture change of Bt horizon not product of compaction.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Tallgrasses (group 1,2,3)

Sub-dominant: Trees (13)

Forbs (11)

Other: All other groups

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Possible mortality only during prolonged drought. Less than 5%.

14. **Average percent litter cover (%) and depth (in):** Litter expected to be at 75% cover at average .25 inch depth.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual production 4000 lb/acre. Ranging from 3300 to 4800 lbs.

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Juniper(ashe juniper/eastern redcedar) most common invader. Also greenbriar, poison ivy, and other woodies will increase without fire.

17. **Perennial plant reproductive capability:** Plants should be capable of reproducing every year with exception of prolonged growing season drought.
