

## Ecological site R086BY001TX Chalky Ridge

Last updated: 9/21/2023  
Accessed: 05/18/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 086B–Texas Blackland Prairie, Southern Part

MLRA 86B, the Southern Part of the Texas Blackland Prairie is located in east-central Texas. It makes up about 2,925 square miles (7,585 square kilometers). The towns of Brenham, Caldwell, La Grange, Schulenberg, Hallettsville, and Navasota are in the eastern part while Lincoln, Benchley, and Normangee are in the western part. The area supports tall and mid-grass prairies, but improved pasture, croplands, and urban development account for the majority of the acreage.

### Classification relationships

USDA-Natural Resources Conservation Service, 2006.  
-Major Land Resource Area (MLRA) 86B

### Ecological site concept

The Chalky Ridge sites are characterized by their shallow soils on ridges with high concentrations of chalk, or calcium carbonates.

## Associated sites

R086BY003TX	<b>Clay Loam</b> The Clay Loam site is often downslope from the Chalky Ridge site. It differs from the site by having deeper soils, higher soil fertility, low to moderate runoff, and lower erosion rates.
R086BY005TX	<b>Blackland</b> The Blackland site is often downslope from the Chalky Ridge site. It differs from the site by having deeper soils, higher soil fertility, and higher soil clay content.

## Similar sites

R086AY001TX	<b>Northern Chalky Ridge</b> Similar but different MLRA
R086AY002TX	<b>Southern Chalky Ridge</b> Similar but different MLRA

Table 1. Dominant plant species

Tree	(1) <i>Quercus virginiana</i> (2) <i>Celtis laevigata</i> var. <i>laevigata</i>
Shrub	(1) <i>Symphoricarpos orbiculatus</i> (2) <i>Smilax rotundifolia</i>
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Andropogon gerardii</i>

## Physiographic features

This site occurs as gently sloping to rolling upland ridges. Slope gradients range from 1 to 20 percent but are usually less than 8 percent.

Table 2. Representative physiographic features

Landforms	(1) Plains > Ridge
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	61–183 m
Slope	1–8%
Water table depth	183 cm
Aspect	Aspect is not a significant factor

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	Not specified
Slope	1–20%
Water table depth	Not specified

## Climatic features

The climate for MLRA 86B is humid subtropical and is characterized by hot summers, especially in July and August,

and relatively mild winters. Tropical maritime air controls the climate during spring, summer and fall. In winter and early spring, frequent surges of Polar Canadian air cause sudden drops in temperatures and add considerable variety to the daily weather. When these cold air masses stagnate and are overrun by moist air from the south, several days of cold, cloudy, and rainy weather follow. Generally, these occasional cold spells are of short duration with rapid clearing following cold frontal passages. The summer months have little variation in day-to-day weather except for occasional thunderstorms that dissipate the afternoon heat. The moderate temperatures in spring and fall are characterized by long periods of sunny skies, mild days, and cool nights. Rainfall during the spring and summer months generally falls during thunderstorms, and fairly large amounts of rain may fall in a short time. High intensity rains of short duration are likely to produce rapid runoff almost anytime during the year. The amount of rain that falls varies considerably from month-to-month and from year-to-year.

**Table 4. Representative climatic features**

Frost-free period (average)	254 days
Freeze-free period (average)	280 days
Precipitation total (average)	1,067 mm

## Climate stations used

- (1) HALLETTSVILLE 2 N [USC00413873], Hallettsville, TX
- (2) LEXINGTON [USC00415193], Lexington, TX
- (3) BRENHAM [USC00411048], Brenham, TX
- (4) FLATONIA 4SE [USC00413183], Moulton, TX
- (5) WASHINGTON SP [USC00419491], Navasota, TX

## Influencing water features

A stream does not influence the plant community of this site.

## Wetland description

Wetlands are not associated with this site.

## Soil features

The soils are dark brown to brownish-gray calcareous clay loams. The surface soil, usually low in organic matter, is underlain by strongly cemented, calcareous sandstone at depths of 11 to 15 inches. The shallow soils severely limit moisture and fertility holding capacity. Roots readily penetrate the parent material, but moisture and minerals are generally not sufficient to support dense stands of high producing grasses. Surface compaction is common. Runoff is high and erosion is a problem when the soil is bare. Soils correlated to this site include: Renish and Shiner.

**Table 5. Representative soil features**

Parent material	(1) Residuum—calcareous sandstone
Surface texture	(1) Clay loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Very slow
Soil depth	28–38 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0–10%
Available water capacity (0-101.6cm)	4.57–6.1 cm


Calcium carbonate equivalent (0-101.6cm)	10–35%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	6–7%
Subsurface fragment volume >3" (Depth not specified)	3–20%

## Ecological dynamics

This is a prairie site that was protected from extensive cultivation because of its shallow topsoil and the hard sandstone that lies beneath. Large live oak (*Quercus virginiana*) trees are scattered sparingly on this site in the western portion of the MLRA while hackberry (*Celtis laevigata*) and bumelia (*Sideroxylon lanuginosum*) are the most common woody species on the eastern fringe. Woody canopy is less than 10 percent in the reference plant community. Little bluestem (*Schizachyrium scoparium*), Indiangrass (*Sorghastrum nutans*), big bluestem (*Andropogon gerardii*), sideoats grama (*Bouteloua curtipendula*), and meadow dropseed (*Sporobolus compositus*) are the dominant grass species on this site. Gayfeather (*Liatris* sp.), black sampson (*Echinacea angustifolia*), cobea penstemon (*Penstemon cobaea*), Engelmann's daisy (*Engelmannia peristenia*), Illinois bundleflower (*Desmanthus illinoensis*), and western indigo (*Indigofera miniata*) are important forbs in the herbaceous community.

## State and transition model


### Ecosystem states



**T1A** - No fire, no brush management, improper grazing management, drought

**R2A** - Fire, brush management, proper grazing, range planting


### State 1 submodel, plant communities



**1.1A** - No fire, no brush management, improper grazing management, drought

**1.2A** - Fire, brush management, proper grazing

State 2 submodel, plant communities



2.1A - No fire, no brush management, improper grazing management, drought

State 1  
Prairie/Savannah

Two communities exist in the Prairie/Savannah State: the 1.1 Tallgrass Prairie Community and the 1.2 Live Oak/Red Cedar Savannah Community. Community 1.1 is characterized by tallgrasses dominating the understory, with woody cover less than 10 percent. Community 1.2 is characterized by a decrease in tallgrasses and an increase in midgrasses and forbs. Woody cover increases to 10 to 25 percent.

Community 1.1  
Tallgrass Prairie



The reference plant community of this site is a prairie or very open savannah sparsely covered with single live oak, hackberry, or bumelia trees, or scattered woody mottes that provide 10 percent or less canopy. The scattered trees, tall native grasses, colorful wildflowers, and rolling topography make this an attractive site. The herbaceous plant community is dominated by little bluestem which generally produces 50 to 75 percent of the total annual production. Indiangrass, big bluestem, sideoats grama, and meadow dropseed are important components of the warm season grass population. Virginia wildrye (*Elymus virginicus*), Canada wildrye (*Elymus canadensis*), and Texas wintergrass (*Nassella leucotricha*) are components of the cool season grass population. Important forbs include gayfeather, Maximillian sunflower (*Helianthus maximilliani*), Engelmann’s daisy, cobeia penstemon, black samson, blue sage (*Salvia azurea*), compass plant (*Silphium* sp.), bundleflower, and western indigo. Grazing prescriptions that permit acceptable grazing periods and allow adequate rest periods along with prescribed fire every three to five years are important in the maintenance of the reference plant community and the prairie landscape structure. Continuous over grazing, or over rest, and the absence of fire tend to favor a transition towards a community dominated by hackberry, prickly pear (*Opuntia* sp.), eastern red cedar (*Juniperus virginiana*), cedar elm (*Ulmus crassifolia*), threeawn (*Aristida* sp.), Texas wintergrass, and annual broomweed (*Amphiachyris amoena*). Without corrective grazing management and burning, the transition will continue to a woodland state dominated by cedar, bumelia, elm, and hackberry.

Table 6. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	2102	3110	4203
Forb	280	420	560
Tree	112	168	224
Shrub/Vine	56	84	112
<b>Total</b>	<b>2550</b>	<b>3782</b>	<b>5099</b>

**Table 7. Ground cover**

Tree foliar cover	0-10%
Shrub/vine/liana foliar cover	0-10%
Grass/grasslike foliar cover	30-60%
Forb foliar cover	5-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	10-20%
Surface fragments >0.25" and <=3"	0-20%
Surface fragments >3"	0-10%
Bedrock	0%
Water	0%
Bare ground	10-20%

**Table 8. Soil surface cover**

Tree basal cover	0-10%
Shrub/vine/liana basal cover	0-5%
Grass/grasslike basal cover	10-30%
Forb basal cover	1-3%
Non-vascular plants	0%
Biological crusts	0%
Litter	10-20%
Surface fragments >0.25" and <=3"	0-20%
Surface fragments >3"	0-10%
Bedrock	0%
Water	0%
Bare ground	10-20%

**Table 9. Canopy structure (% cover)**

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	—	0-5%
>0.15 <= 0.3	—	—	5-10%	0-5%
>0.3 <= 0.6	—	—	10-20%	0-5%
>0.6 <= 1.4	—	0-10%	20-40%	5-10%
>1.4 <= 4	—	—	0-5%	0-5%
>4 <= 12	0-10%	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Community 1.2

### Live Oak/Red Cedar Savannah

This plant community is a transitional community between the Prairie and the Shrubland State. It develops in the absence of fire and proper grazing management, or the absence of chemical or biological brush management. It is usually the result of abandonment following yearly continuous grazing. In addition to the naturally occurring trees, eastern red cedar and bois d'arc (*Maclura pomifera*) increase in density and canopy coverage (10 to 25 percent canopy). In some cases, eastern red cedar may dominate the component of the community. Species whose seed is wind blown (elm) or animal dispersed (cedar, bumelia, bois d'arc) are the first to invade and dominate the site. Remnants of little bluestem, Indiangrass, and sideoats grama may still occur, but the herbaceous component of the community becomes dominated by lesser producing grasses and forbs. Silver bluestem (*Bothriochloa laguroides*), fall witchgrass (*Digitaria cognata*), threeawn, western ragweed (*Ambrosia psilostachya*), annual broomweed (*Amphiachyris dracunculoides*), and snow on the prairie (*Euphorbia bicolor*) commonly occur. Since the woody shrub canopy has not exceeded 40 percent, prescribed burning on a five to seven year interval in conjunction with prescribed grazing is a viable option for returning this community to a tallgrass prairie that may resemble the reference plant community.

Table 10. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	1345	2130	2914
Forb	560	1009	1345
Shrub/Vine	224	336	448
Tree	112	168	224
<b>Total</b>	<b>2241</b>	<b>3643</b>	<b>4931</b>

## Pathway 1.1A

### Community 1.1 to 1.2

The Tallgrass Prairie Plant Community will shift to the Live Oak/Red Cedar Savannah Plant Community when there is continued growing-season stress on reference grass species. These stresses include improper grazing management that creates insufficient critical growing-season deferment, excess intensity of defoliation, repeated, long-term growing-season defoliation, long-term drought, and/or other repeated critical growing-season stress. Increaser species (midgrasses and woody species) are generally endemic species released by disturbance. Woody species canopy exceeding 10 percent and a species shift from tallgrass to midgrass composition indicate a transition. The driver for the shift is heavy grazing, no fire, no brush management, and/or abandonment.

## Pathway 1.2A

### Community 1.2 to 1.1

The 1.2 Plant Community will return to the Tallgrass Prairie Plant Community under grazing management that provides sufficient critical growing season deferment in combination with proper grazing intensity as long as the seedbank or seed source is still present. Favorable moisture conditions will facilitate or accelerate this transition. The understory component may return to dominance by tallgrasses in the absence of fire or brush control. However, reduction of the woody component to reference conditions of 10 percent or less canopy cover will require inputs of fire or brush control.

## State 2 Shrubland

Two communities exist in the Shrubland State: the 2.1 Live Oak/Red Cedar Shrubland Community and the 2.2 Live Oak/Red Cedar Woodland Community. Community 2.1 is characterized by a shift in grass composition to silver bluestem and threeawns. The overstory canopy increases to 25 to 40 percent. Community 2.2 is characterized by a shortgrasses and low forbs with greater than 40 percent overstory canopy cover.

### Community 2.1 Live Oak/Red Cedar Shrubland



The Live Oak/Red Cedar Shrubland develops in the absence of fire and proper grazing management, or the absence of chemical or biological brush management. It is usually the result of abandonment following yearly continuous grazing. In addition to the naturally occurring trees, eastern red cedar and bois d'arc (*Maclura pomifera*) increase in density and canopy coverage (25 to 40 percent). In some cases, eastern red cedar may dominate the component of the community. Species whose seed is wind blown (elm) or animal dispersed (cedar, bumelia, bois d'arc) are the first to invade and dominate the site. Remnants of little bluestem, Indiangrass, and sideoats grama may still occur, but the herbaceous component of the community becomes dominated by lesser producing grasses and forbs. Silver bluestem, fall witchgrass, threeawn, Western ragweed, annual broomweed, and snow on the prairie commonly occur. Since the woody shrub canopy still has not exceeded 40 percent, prescribed burning on a five to seven year interval in conjunction with prescribed grazing is a viable option for returning this community to a tallgrass prairie that may resemble the reference plant community. The presence of sandstone close to the soil surface precludes most methods of mechanical control other than hand cutting.

Table 11. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Forb	1065	1597	2130
Grass/Grasslike	785	1177	1569
Shrub/Vine	336	504	673
Tree	112	168	224
<b>Total</b>	<b>2298</b>	<b>3446</b>	<b>4596</b>

Table 12. Ground cover



Tree foliar cover	10-40%
Shrub/vine/liana foliar cover	10-20%
Grass/grasslike foliar cover	30-40%
Forb foliar cover	20-30%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-20%
Surface fragments >0.25" and <=3"	0-20%
Surface fragments >3"	0-10%
Bedrock	0%
Water	0%
Bare ground	20-40%

**Table 13. Soil surface cover**

Tree basal cover	15-20%
Shrub/vine/liana basal cover	5-15%
Grass/grasslike basal cover	10-20%
Forb basal cover	5-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-20%
Surface fragments >0.25" and <=3"	0-20%
Surface fragments >3"	0-10%
Bedrock	0%
Water	0%
Bare ground	20-40%

**Table 14. Canopy structure (% cover)**

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	5-10%	—
>0.15 <= 0.3	—	—	5-10%	—
>0.3 <= 0.6	—	—	20-40%	5-15%
>0.6 <= 1.4	—	—	5-15%	5-15%
>1.4 <= 4	—	10-20%	—	—
>4 <= 12	20-40%	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Community 2.2

### Live Oak/Red Cedar Woodland



This plant community is dominated by woody species including eastern red cedar, bumelia, hackberry, cedar elm, and live oak. Canopy cover exceeds 40 percent. Understory shrubs and vines include coralberry (*Symphoricarpos orbiculatus*), greenbriar (*Smilax* sp.), grape (*Vitis* sp.), prickly pear, and tasajillo (*Opuntia leptocaulis*). Herbaceous composition and production varies with woody canopy cover but Texas wintergrass, cedar sedge (*Carex planostachys*), silver bluestem, threeawn, annual broomweed, and snow on the prairie are common occupants of the site. Treatment options to move this community back towards the reference community are limited. If the woody canopy has exceeded 40 percent, chemical brush control must be applied. The presence of sandstone close to the soil surface precludes most methods of mechanical treatments other than handcutting or tree dozing. Commercial aerial herbicide application is usually impractical due to the limited extent of the site, while the presence of large trees hinders the effectiveness of ground applied herbicides. Prescribed burning is usually not a viable alternative due to the lack of fine fuel.

**Table 15. Annual production by plant type**

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Tree	841	1261	1681
Forb	729	1093	1457
Grass/Grasslike	448	673	897
Shrub/Vine	336	504	673
<b>Total</b>	<b>2354</b>	<b>3531</b>	<b>4708</b>

**Table 16. Ground cover**

Tree foliar cover	40-60%
Shrub/vine/liana foliar cover	20-40%
Grass/grasslike foliar cover	10-20%
Forb foliar cover	15-30%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-10%
Surface fragments >0.25" and <=3"	0-20%
Surface fragments >3"	0-10%
Bedrock	0%
Water	0%
Bare ground	20-40%

**Table 17. Soil surface cover**

Tree basal cover	20-40%
Shrub/vine/liana basal cover	15-25%
Grass/grasslike basal cover	5-10%
Forb basal cover	5-10%
Non-vascular plants	0%
Biological crusts	0%
Litter	5-10%
Surface fragments >0.25" and <=3"	0-20%
Surface fragments >3"	0-10%
Bedrock	0%
Water	0%
Bare ground	20-40%

**Table 18. Canopy structure (% cover)**

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	5-10%	5-10%
>0.15 <= 0.3	—	—	—	—
>0.3 <= 0.6	—	—	5-10%	10-20%
>0.6 <= 1.4	—	—	5-10%	—
>1.4 <= 4	—	15-25%	—	—
>4 <= 12	40-60%	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Pathway 2.1A Community 2.1 to 2.2



Live Oak/Red Cedar Shrubland



Live Oak/Red Cedar Woodland

The Shrubland Community will transition to the Woodland Community with continued heavy grazing, no fire, no brush management and/or abandonment. Indicators of the transition include an increased species composition of shortgrasses and trees becoming more mature.

## Transition T1A State 1 to 2

The Prairie/Savannah State will transition to the Shrubland State when continued heavy grazing pressure, no fire, no brush management, and/or field abandonment continues. The transition is evident when woody species canopy cover exceeds 25 percent and grasses shift composition to silver bluestem, threeawns, and shortgrasses.

## Restoration pathway R2A State 2 to 1

Restoration of the Shrubland State to the Prairie/Savannah State requires substantial energy input. Mechanical or herbicidal brush control treatments can be used to remove woody species. A long-term prescribed fire program may sufficiently reduce brush density particularly if the woody component is dominated by species that are not re-sprouters. However, fire may not be sufficient to remove mature trees. Brush control in combination with prescribed fire, proper grazing management, and favorable growing conditions may be the most economical means of creating and maintaining the desired plant community. If remnant populations of tallgrasses, midgrasses, and desirable forbs are not present at sufficient levels, range planting will be necessary to restore a plant community similar to the reference community.

## Additional community tables

Table 19. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Tallgrass</b>			1345–2690	
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	1345–2690	–
2	<b>Tallgrasses</b>			364–729	
	big bluestem	ANGE	<i>Andropogon gerardii</i>	364–729	–
	Indiangrass	SONU2	<i>Sorghastrum nutans</i>	364–729	–
3	<b>Midgrasses</b>			252–504	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	252–504	–
	silver beardgrass	BOLAT	<i>Bothriochloa laguroides</i> ssp. <i>torreyana</i>	252–504	–
	composite dropseed	SPCOC2	<i>Sporobolus compositus</i> var. <i>compositus</i>	252–504	–
4	<b>Mid/Shortgrasses</b>			112–224	
	purple threeawn	ARPU9	<i>Aristida purpurea</i>	112–224	–
	Wright's threeawn	ARPUW	<i>Aristida purpurea</i> var. <i>wrightii</i>	112–224	–
	Canada wildrye	ELCA4	<i>Elymus canadensis</i>	112–224	–
	Virginia wildrye	ELVI3	<i>Elymus virginicus</i>	112–224	–
	Texas wintergrass	NALE3	<i>Nassella leucotricha</i>	112–224	–
	Florida paspalum	PAFL4	<i>Paspalum floridanum</i>	112–224	–
	slim tridens	TRMU	<i>Tridens muticus</i>	112–224	–
<b>Forb</b>					
5	<b>Forbs</b>			280–504	
	purple dalea	DALA4	<i>Dalea lasiathera</i>	280–504	–
	pointedleaf ticktrefoil	DEGL5	<i>Desmodium glutinosum</i>	280–504	–
	Illinois bundleflower	DEIL	<i>Desmanthus illinoensis</i>	280–504	–
	blacksamson echinacea	ECANA	<i>Echinacea angustifolia</i> var. <i>angustifolia</i>	280–504	–
	Engelmann's daisy	ENPE4	<i>Engelmannia peristenia</i>	280–504	–
	Maximilian sunflower	HEMA2	<i>Helianthus maximiliani</i>	280–504	–
	coastal indigo	INMI	<i>Indigofera miniata</i>	280–504	–
	dotted blazing star	LIPUP	<i>Liatris punctata</i> var. <i>punctata</i>	280–504	–
	Nuttall's sensitive-briar	MINU6	<i>Mimosa nuttallii</i>	280–504	–
	yellow puff	NELU2	<i>Neptunia lutea</i>	280–504	–
	whitest evening	OEAL	<i>Oenothera albicaulis</i>	280–504	–

	primrose				
	cobaea beardtongue	PECO4	<i>Penstemon cobaea</i>	280–504	–
	Texas snoutbean	RHSE4	<i>Rhynchosia senna</i>	280–504	–
	fringeleaf wild petunia	RUHU	<i>Ruellia humilis</i>	280–504	–
	azure blue sage	SAAZ	<i>Salvia azurea</i>	280–504	–
	compassplant	SILA3	<i>Silphium laciniatum</i>	280–504	–
	amberique-bean	STHE9	<i>Strophostyles helvola</i>	280–504	–
6	<b>Forbs</b>			28–56	
	Cuman ragweed	AMPS	<i>Ambrosia psilostachya</i>	28–56	–
	downy Indian paintbrush	CAPU11	<i>Castilleja purpurea</i>	28–56	–
	California larkspur	DECA	<i>Delphinium californicum</i>	28–56	–
	snow on the prairie	EUBI2	<i>Euphorbia bicolor</i>	28–56	–
	plains beeblossom	GABR2	<i>Gaura brachycarpa</i>	28–56	–
	Texas lupine	LUTE	<i>Lupinus texensis</i>	28–56	–
	Drummond's skullcap	SCDR2	<i>Scutellaria drummondii</i>	28–56	–
<b>Shrub/Vine</b>					
7	<b>Vines</b>			56–112	
	roundleaf greenbrier	SMRO	<i>Smilax rotundifolia</i>	56–112	–
	coralberry	SYOR	<i>Symphoricarpos orbiculatus</i>	56–112	–
	muscadine	VIRO3	<i>Vitis rotundifolia</i>	56–112	–
<b>Tree</b>					
8	<b>Trees</b>			168–336	
	sugarberry	CELA	<i>Celtis laevigata</i>	168–336	–
	live oak	QUVI	<i>Quercus virginiana</i>	168–336	–
	gum bully	SILAA4	<i>Sideroxylon lanuginosum</i> ssp. <i>albicans</i>	168–336	–
	cedar elm	ULCR	<i>Ulmus crassifolia</i>	168–336	–

## Animal community

Historical accounts of early Texas explorers indicate that the tallgrass prairie sites were used by bison, deer, antelope, prairie chicken, wolf, mountain lion, and black bear. Wild horses and wild cattle occupied these prairie sites into the middle 1800's. While most prairie sites were converted to cropland, the shallow rocky nature of this site precluded that. Habitat values and potential plant communities for this site have been influenced by past grazing management or lack of. Deer, bobcat, coyote, quail, turkey, dove, and a host of other mammals, birds, and reptiles use this site when woody cover is sufficient. Being of limited extent, the habitat values of this site are influenced by the quality of the habitat of the surrounding ecological sites. Manipulation of the plant community and habitat types by methods other than fire and grazing management are somewhat limited by the shallow rocky soil.

## Hydrological functions

Because of the fractured sandstone horizon and the potential herbaceous plant community, this site may provide ground water recharge or contribute to surface runoff. Peak rainfall periods occur in May and June from frontal passage thunderstorms and in September and October from tropical systems, as well as frontal passages. Rainfall amounts may be high (three to five inches per event) and events may be intense. The site is subject to erosion where adequate herbaceous cover is not maintained and on heavy use areas such as roads and livestock trails. Extended periods (60 days) of little to no rainfall during the growing season are common and can limit herbaceous production and ground cover. The hydrology of this site may be manipulated through management to yield higher runoff volumes or greater infiltration to ground water. Management for less herbaceous cover will favor higher

surface runoff while dense herbaceous cover will favor ground water recharge. Potential pollution from sediment and pesticides should always be considered when managing for higher volumes of surface runoff.

## Recreational uses

Recreational uses of this site may include hunting, bird watching, photography, horseback riding, cycling, hiking, and camping.

## Inventory data references

These site descriptions were developed as part a Provisional Ecological Site project using historic soil survey manuscripts, available site descriptions, and low intensity field traverse sampling. Future work to validate the information is needed. This will include field activities to collect low, medium, and high-intensity sampling, soil correlations, and analysis of that data. A final field review, peer review, quality control, and quality assurance review of the will be needed to produce the final document.

## Other references

Diggs, G.M., B.L. Lipscomb, and R.J. O'Kennon. 1999. Illustrated Flora of North Central Texas Botanical Research Institute of Texas and Austin College, Fort Worth, TX.

Hatch, S.L, K.N. Gandhi, and L.E. Brown. 1990. Checklist of the vascular plants of Texas. Texas Agricultural Experiment Station, College Station, TX.

## Contributors

Mark Moseley  
Tyson Hart

## Approval

Bryan Christensen, 9/21/2023

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Mike Stellbauer
Contact for lead author	Mike Stellbauer, Zone RMS, NRCS, Bryan, Texas
Date	05/23/2005
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** A few short rills may be present due to landscape position, slope, and production potential of the site.
-

2. **Presence of water flow patterns:** Some water flow patterns are normal for this site due to landscape position and slope.
- 
3. **Number and height of erosional pedestals or terracettes:** A few slightly elevated pedestals or terracettes may occur due to slope, landscape position, and natural lack of cover on this site.
- 
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** No more than 30 percent bare ground randomly distributed.
- 
5. **Number of gullies and erosion associated with gullies:** None
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Small to medium-size litter movement for short distances should be expected for this site during intense rainfall events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface is resistant to erosion. Stability class range is 4 to 6.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is 7 to 15 inches thick with colors of very dark brown to very dark grayish brown with moderate fine to very fine subangular blocky structure. SOM is 1 to 3 percent.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The predominance of tallgrasses, midgrasses, perennial forbs, and adequate litter provide for maximum infiltration and reduced runoff under normal rainfall events.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm-season tallgrasses >>
- Sub-dominant: Warm-season midgrasses > Forbs >
- Other: Trees > Shrubs

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** There should be little mortality or decadence in any functional groups.
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 2,275 pounds per acre for below average moisture years to 4,550 pounds per acre for above average moisture years.
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Potential invasive species include hackberry, elm, yellow bluestem, common Bermudagrass, eastern red cedar, pricklypear and osage orange.
- 
17. **Perennial plant reproductive capability:** Perennial plants are capable of adequate reproduction, except during periods of prolonged drought conditions, heavy natural herbivory, and intense wildfires.
-