

## Ecological site R086BY001TX Chalky Ridge

Last updated: 9/21/2023  
Accessed: 05/04/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Mike Stellbauer
Contact for lead author	Mike Stellbauer, Zone RMS, NRCS, Bryan, Texas
Date	05/23/2005
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- Number and extent of rills:** A few short rills may be present due to landscape position, slope, and production potential of the site.

---
- Presence of water flow patterns:** Some water flow patterns are normal for this site due to landscape position and slope.

---
- Number and height of erosional pedestals or terracettes:** A few slightly elevated pedestals or terracettes may occur due to slope, landscape position, and natural lack of cover on this site.

---
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** No more than 30 percent bare ground randomly distributed.

---
- Number of gullies and erosion associated with gullies:** None

---
- Extent of wind scoured, blowouts and/or depositional areas:** None

---

7. **Amount of litter movement (describe size and distance expected to travel):** Small to medium-size litter movement for short distances should be expected for this site during intense rainfall events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface is resistant to erosion. Stability class range is 4 to 6.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is 7 to 15 inches thick with colors of very dark brown to very dark grayish brown with moderate fine to very fine subangular blocky structure. SOM is 1 to 3 percent.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The predominance of tallgrasses, midgrasses, perennial forbs, and adequate litter provide for maximum infiltration and reduced runoff under normal rainfall events.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Warm-season tallgrasses >>
- Sub-dominant: Warm-season midgrasses > Forbs >
- Other: Trees > Shrubs
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** There should be little mortality or decadence in any functional groups.
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 2,275 pounds per acre for below average moisture years to 4,550 pounds per acre for above average moisture years.
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not**

**invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Potential invasive species include hackberry, elm, yellow bluestem, common Bermudagrass, eastern red cedar, pricklypear and osage orange.

---

17. **Perennial plant reproductive capability:** Perennial plants are capable of adequate reproduction, except during periods of prolonged drought conditions, heavy natural herbivory, and intense wildfires.
-