

## Ecological site F096XA014MI Snowy Mucky Depression

Last updated: 10/03/2019  
Accessed: 04/27/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 096X–Northwestern Michigan Fruit Belt

This area is dominated by outwash plains and moraines. Lake plains, till plains, drumlins, and sand dunes are found locally across the area. The terrain is steep on stream carved moraines, ice contact ridges, and sand dunes, and flat on outwash plains and lake plains. Elevation ranges from 177 to 369 m (580 to 1210 ft). Local topographic relief averages 11 m (35 ft) in the south to 20 m (65 ft) in the north and ranges up to a maximum of 158 m (520 ft) at Empire Bluff (Sleeping Bear Dunes). Much of the area rises sharply from the lakeshore to the adjoining hilltops. The Manistee River is the longest river in this area. Its trout fishery is maintained by constant inflow of cool ground water from the porous sand dominated landscape. The Pine and Pere Marquette Rivers also occur in this MLRA. Surficial topography are formed of glacial deposits except for local areas with dune building near Lake Michigan. Most of the bedrock surface is at or below the elevation of Lake Michigan, and is exposed in only in limited extents near Charlevoix. The bedrock, all Paleozoic in age, is the Traverse Group and the Dundee Limestone. These Silurian-Devonian rocks are mostly limestone and dolomite with some interbedded shale, chert, and anhydrite stringers. The drumlin belts in the northern portion of the area is the most affected by the limestone nearer to the surface in terms of carbonates in the till.

About two-thirds of this area is in small, privately owned holdings, and one-third consists mostly of State forests. The forests are used mainly for timber production and recreation. The growth of orchard crops and other crops and dairy and beef operations are important enterprises in the area. Forage and feed grains for dairy and other livestock

are the principal crops. Asparagus, wheat, oats, corn, and hay are commonly grown in the area. Orchard products include sweet and tart cherries, apples, plums, and peaches. The Manistee National Forest and Sleeping Bear Dunes National Lakeshore are among the more notable conservation lands in the area. Nordhouse Dunes Wilderness Area is within the Manistee National Forest. Sections of the Pere Marquette, Pine, and Manistee Rivers, and Bear Creek have been designated as National Wild and Scenic Rivers.

Summary of existing land use:

- Upland Forest (47%)
- Hardwood (38%)
- Conifer (7%)
- Developed (16%)
- Swamps and Marshes (12%)
- Agricultural (10%)
- Open Water (8%)
- Grassland (6%)

### Classification relationships

According to the USFS (Bailey) system of ecoregions, the site is located mostly within 212Hd (Grand Traverse Ground Moraine) and 212Hf (Grand Traverse Drumlin Fields) subsections. According to the EPA (Omernik) system of ecoregions, the site is located in 51m (Manistee-Leelanau Shore) and 51n (Platte River Outwash) level IV ecoregions. This site is outside the environmental range of the Kotar system. This site corresponds to the Organic Wetland, ecological land type phase, 82, in the USFS Ecological Land Type system.

### Ecological site concept

The central concept of Snowy Mucky Depression is lowlands on hydric organic or marl soils with a pH greater than 4.5 (euic histosols). Site is in the heavy annual snowfall belt, mostly north of Manistee River, where fire was rare. Area with a more northern flora than sites further south. Vegetation ranging from rich conifer swamp to fen.

Table 1. Dominant plant species

Tree	(1) <i>Thuja occidentalis</i> (2) <i>Fraxinus nigra</i>
Shrub	Not specified
Herbaceous	(1) <i>Cystopteris bulbifera</i> (2) <i>Caltha palustris</i>

### Physiographic features

Site found in depressions and lower slope positions of various glacial landforms, especially if coarse textured, allowing for groundwater movement.

Table 2. Representative physiographic features

Landforms	(1) Depression
-----------	----------------

### Climatic features

Mean annual temperatures are 7.1 to 8.0 °C (45 to 46 °F). The warmest six months average 15.3 to 16.2 °C (60 to 61 °F). Mean July temperatures range from 19.8 to 20.7 °C (68 to 69 °F). Mean January temperatures range from -6.7 to -4.2 °C (20 to 24 °F). The maximum monthly average daily highs are 24.2 to 27.4 °C (76 to 81 °F). The minimum monthly average daily lows are -10.8 to -7.2 °C (13 to 19 °F). Mean annual precipitation ranges from 780 to 880 mm (31 to 35 in). Prevailing winds pick up moisture from the Great Lakes in the form of lake effect rain and snow showers during fall and winter seasons, and in the form of fog during spring and summer. Thunderstorm intensity is reduced by temperature inversions over the lake during the spring and early summer when lake water is cools the air flowing over it. Average 0 °C (32 °F) frost-free season ranges from 108 to 161 days. Average -2 °C (28

°F) freeze-free season is 141 to 194 days increasing in length from north to south and decreasing in length from the lakeshore inward. Mean annual snowfall ranges from 2.1 to 3.4 m (60 to 130 in). Mean annual extreme minimum temperatures range from -29 to -18.9 °C (-20 to -2 °F), or hardiness zones 4b to 6b.

**Table 3. Representative climatic features**

Frost-free period (average)	122 days
Freeze-free period (average)	155 days
Precipitation total (average)	34 in

### **Climate stations used**

- (1) NORTHPORT 2W [USC00206007], Northport, MI
- (2) TRAVERSE CITY CHERRY CPTL AP [USW00014850], Traverse City, MI
- (3) CHARLEVOIX [USC00201468], Charlevoix, MI
- (4) MAPLE CITY 1E [USC00205097], Cedar, MI
- (5) PETOSKEY [USC00206507], Petoskey, MI
- (6) FRANKFORT 2NE [USC00202984], Frankfort, MI
- (7) NW MICHIGAN RSCH FM [USC00206012], Suttons Bay, MI

### **Influencing water features**

Site has a stable source of minerotrophic groundwater flowing through it at less than 25 cm in depth.

### **Soil features**

Soils are very poorly drained circumneutral mucks to calcareous marl. They are commonly classified Terric Borosaprists, Typic Haplosaprists, and Terric Haplosaprists, and commonly mapped as Adrian, Houghton, and Tawas series or components. The top 50 cm has a typical pH of 6.4 and is 5% sand and 54.1% organic matter. At depth, pH ranges up to 7.1, and texture averages 30% sand and 10% clay. Depth to impeded hydraulic conductivity or root restrictive layers averages >200 cm. Depth to carbonates averages 200 cm.

### **Ecological dynamics**

Snowy Mucky Depression tends to share the same ecological dynamics as Natureserve/Landfire system, Laurentian-Acadian Alkaline Conifer-Hardwood Swamp or Laurentian-Acadian Alkaline Fen. Stand replacing fires occurred every 500-2000 years, while light surface fires were very rare. Overstory was dominated by species like whitecedar (*Thuja occidentalis*) and black ash (*Fraxinus nigra*), which thrive in soils saturated with minerotrophic groundwater. The minerotrophic groundwater seeping into the mucks supports an understory of marsh marigold (*Caltha palustris*) and skunk cabbage (*Symplocarpus foetidus*). Sloping mucks may support carpets of bladder fern (*Cystopteris bulbifera*).

### **State and transition model**

# F096XA014MI Snowy Mucky Depression




Figure 6. stm

## Legend

1.1A	Extreme winds every 222 years or replacement fire every 1000 years
1.2A	Succession
1.3A	Succession
1.3B	Extreme winds every 333 years or replacement fire every 1000 years
1.4A	Succession
1.4B	Extreme winds every 222 years or replacement fire every 1000 years
R2	Restoration
R4	Restoration
R5	Restoration?
R6	Restoration?
T1A	Artificial drainage
T1B	Invasive species introduction
T2	Clearing + cultivation of crops
T3	Abandonment + invasive species
T4	Cultivation

Figure 7. Legend

### State 1

#### Reference State

#### Community 1.1

Rich Conifer Swamp: *Thuja occidentalis* - *Fraxinus nigra* Swamp Forest

#### Community 1.2

Wet Meadow: *Carex utriculata* - *Carex lacustris* - (*Carex vesicaria*, *Carex stricta*) Wet Meadow

#### Community 1.3

Southern Shrub-carr: *Cornus sericea* - *Salix* spp. - (*Rosa palustris*) Shrub Swamp

#### Community 1.4

Rich Tamarack Swamp: *Larix laricina* / *Alnus incana* Swamp Forest

#### Community 1.5

Fen: *Betula pumila* - *Dasiphora fruticosa* ssp. *floribunda* / *Carex lasiocarpa* - *Trichophorum alpinum* Fen

#### Community 1.6

Emergent Marsh: *Schoenoplectus tabernaemontani* - *Typha* spp. - (*Sparganium* spp., *Juncus*

## spp.) Marsh

### Community 1.7

Inundated Shrub Swamp: *Cornus sericea* - *Salix* spp. - (*Rosa palustris*) Shrub Swamp

#### Pathway 1.1A

##### Community 1.1 to 1.2

Temporary prolonged inundation/Fire.

##### Conservation practices

Prescribed Burning
--------------------

#### Pathway 1.1B

##### Community 1.1 to 1.3

Clearcut/Blowdown/Fire.

##### Conservation practices

Prescribed Burning
--------------------

Early Successional Habitat Development/Management
---

Forest Stand Improvement
--------------------------

#### Pathway 1.2A

##### Community 1.2 to 1.1

Succession.

##### Conservation practices

Tree/Shrub Site Preparation
-----------------------------

Tree/Shrub Establishment
--------------------------

#### Pathway 1.2B

##### Community 1.2 to 1.3

Succession.

##### Conservation practices

Tree/Shrub Site Preparation
-----------------------------

Tree/Shrub Establishment
--------------------------

#### Pathway 1.2C

##### Community 1.2 to 1.5

Increase peat or marl thickness, decrease nitrogen or phosphorus availability.

#### Pathway 1.2D

##### Community 1.2 to 1.6

Permanent inundation.

**Pathway 1.2E**  
**Community 1.2 to 1.7**

Permanent inundation.

**Pathway 1.3A**  
**Community 1.3 to 1.1**

Succession.

**Conservation practices**

Tree/Shrub Site Preparation
Tree/Shrub Establishment

**Pathway 1.3B**  
**Community 1.3 to 1.2**

Temporary prolonged inundation/Fire.

**Conservation practices**

Prescribed Burning
--------------------

**Pathway 1.3C**  
**Community 1.3 to 1.6**

Permanent inundation.

**Pathway 1.3D**  
**Community 1.3 to 1.7**

Permanent inundation.

**Pathway 1.4A**  
**Community 1.4 to 1.1**

Succession.

**Pathway 1.4B**  
**Community 1.4 to 1.5**

Clearcut/Blowdown/Fire.

**Conservation practices**

Prescribed Burning
Early Successional Habitat Development/Management
Forest Stand Improvement

**Pathway 1.5A**  
**Community 1.5 to 1.2**

Decrease peat or marl thickness, increase nitrogen or phosphorus availability.

**Pathway 1.5B**  
**Community 1.5 to 1.4**

Succession.

**Conservation practices**

Tree/Shrub Site Preparation
-----------------------------

Tree/Shrub Establishment
--------------------------

**Pathway 1.5C**  
**Community 1.5 to 1.6**

Permanent inundation.

**Pathway 1.6A**  
**Community 1.6 to 1.2**

Drop water table.

**Pathway 1.6B**  
**Community 1.6 to 1.5**

Drop water table.

**Pathway 1.6C**  
**Community 1.6 to 1.7**

Temporary drop water table; shrub establishment.

**Conservation practices**

Tree/Shrub Site Preparation
-----------------------------

Tree/Shrub Establishment
--------------------------

**Pathway 1.7A**  
**Community 1.7 to 1.2**

Drop water table, fire.

**Conservation practices**

Prescribed Burning
--------------------

**Pathway 1.7B**  
**Community 1.7 to 1.3**

Drop water table.

**Pathway 1.7C**  
**Community 1.7 to 1.6**

Temporary drought and fire; shrub mortality.

**Conservation practices**



Brush Management
Prescribed Burning

**State 2  
Cultural State**

**Community 2.1  
Sustainable Crop, Pasture, or Plantation**

**Community 2.2  
Unsustainable Cultural Phase**

**Community 2.3  
Conservation Feature**

Can be a grassed waterway, conservation reserve, a small patch pollinator garden, or other land taken out of its primary cultural production to mitigate or reduce impacts of adjacent land use, and is not by itself a permanent restoration of a complete native biological community and associated ecosystem services.

**Pathway 2.1A  
Community 2.1 to 2.2**

Revert to unsustainable cultural practices.

**Pathway 2.1B  
Community 2.1 to 2.3**

Establish conservation feature.

**Conservation practices**

Conservation Cover
Grassed Waterway

**Pathway 2.2A  
Community 2.2 to 2.1**

Implement sustainable cultural practices.

**Conservation practices**

Conservation Crop Rotation
Cover Crop
Nutrient Management
Integrated Pest Management (IPM)

**Pathway 2.2B  
Community 2.2 to 2.3**

Establish conservation feature.

**Conservation practices**

Conservation Cover
--------------------

Grassed Waterway

**Pathway 2.3A**  
**Community 2.3 to 2.1**

Implement sustainable cultural practices.

**Conservation practices**

Conservation Crop Rotation
Cover Crop
Nutrient Management
Integrated Pest Management (IPM)

**Pathway 2.3B**  
**Community 2.3 to 2.2**

Revert to unsustainable cultural practices.

**State 3**  
**Seminatural Drained State**

**Community 3.1**  
**Ruderal Drained Meadow & Shrub: *Dactylis glomerata* - *Phleum pratense* - *Festuca* spp. - *Solidago* spp. Ruderal Meadow**

**Community 3.2**  
**Exotic Ruderal Drained Forest: *Acer platanoides* Ruderal Forest**

**Pathway 3.1A**  
**Community 3.1 to 3.2**

Succession

**Pathway 3.2A**  
**Community 3.2 to 3.1**

Blowdown/clearcut.

**Conservation practices**

Early Successional Habitat Development/Management
Forest Stand Improvement

**State 4**  
**Seminatural State**

**Community 4.1**  
**Ruderal Wet Meadow & Shrub Swamp: *Phalaris arundinacea* Eastern Ruderal Marsh**

**Community 4.2**  
**Exotic Ruderal Swamp Forest: *Acer negundo* Ruderal Floodplain Forest**

**Pathway 4.1A**  
**Community 4.1 to 4.2**

Succession.

**Pathway 4.2A**  
**Community 4.2 to 4.1**

Blowdown/clearcut.

**Conservation practices**

Early Successional Habitat Development/Management
Forest Stand Improvement

**Transition T1A**  
**State 1 to 2**

Drain; clear vegetation; cultivate domesticated species.

**Transition T1B**  
**State 1 to 3**

Drain; clear vegetation, invasive species introduced.

**Transition T1C**  
**State 1 to 4**

Clear vegetation, invasive species introduced.

**Restoration pathway R2**  
**State 2 to 1**

Restore hydrology; remove domesticated species; restore native species.

**Conservation practices**

Brush Management
Restoration and Management of Rare and Declining Habitats
Wetland Wildlife Habitat Management
Wetland Restoration
Herbaceous Weed Control

**Transition T2A**  
**State 2 to 3**

Abandon, succession.

**Transition T2B**  
**State 2 to 4**

Restore hydrology; abandon; succession.

**Conservation practices**

Wetland Restoration
---------------------

**Restoration pathway R3  
State 3 to 1**

Restore hydrology; control invasive species; restore native species

**Conservation practices**

Brush Management
Restoration and Management of Rare and Declining Habitats
Wetland Wildlife Habitat Management
Wetland Restoration
Herbaceous Weed Control

**Transition T3A  
State 3 to 2**

Clear vegetation; cultivate domesticated species.

**Transition T3B  
State 3 to 4**

Restore hydrology.

**Conservation practices**

Wetland Restoration
---------------------

**Restoration pathway R4  
State 4 to 1**

Control invasive species; restore native species.

**Conservation practices**

Brush Management
Restoration and Management of Rare and Declining Habitats
Wetland Wildlife Habitat Management
Herbaceous Weed Control

**Transition T4A  
State 4 to 2**

Drain; clear vegetation; cultivate domesticated species.

**Transition T4B  
State 4 to 3**

Drain.

**Additional community tables**

## Other references

A PROVISIONAL ECOLOGICAL SITE is a conceptual grouping of soil map unit components within a major land resource area (MLRA) based on the similarities in response to management. A provisional ecological site is a first approximation based on a cursory literature review, personal experience, and limited field reconnaissance. As more adequate literature review, expert opinion, and intensive plot data are collected, the site concept is subject to shifting, broadening, narrowing, subdivision, or re-aggregation in definition. Likewise, the community dynamics will be more elaborate in content, and may also change in structure, upon reaching approved status.

Future work, as described in a project plan, to validate the information in this provisional ecological site description is needed. This will include field activities to collect low and medium intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final document. Annual reviews of the project plan are to be conducted by the Ecological Site Technical Team.

Albert, D. A. et al., 1995. Vegetation circa 1800 of Michigan. Michigan's native landscape as interpreted from the General Land Office Surveys 1816-1856 (digital map), Lansing: Michigan Natural Features Inventory.

Baker, M.E. and Barnes, B.V., 1998. Landscape ecosystem diversity of river floodplains in northwestern Lower Michigan, USA. *Canadian Journal of Forest Research*, 28(9), pp.1405-1418.

Barnes, B. V. and Wagner, W. H., 2004. Michigan trees: a guide to the trees of the Great Lakes region. Ann Arbor (Michigan): University of Michigan Press.

Burger, T. L. and Kotar, J., 2003. A Guide to Forest Communities and Habitat Types of Michigan. Madison, Wisconsin: Department of Forest Ecology and Management, University of Wisconsin.

Cleland, D. T. et al., 1994. Field guide: Ecological classification and inventory system of the Huron-Manistee National Forests, s.l.: USDA Forest Service, North Central Forest Experiment Station.

Eichenlaub, V.L., 1979. Weather and climate of the Great Lakes region. University of Notre Dame Press, Indiana. 335 pages.

GHCN, 2016. Global Historical Climatology Network Monthly Versions 2 and 3 (temperature and precipitation data). NOAA. <https://www.ncdc.noaa.gov/ghcnm/>

Kost, M. A. et al., 2010. Natural Communities of Michigan: Classification and Description, Lansing, MI: Michigan Natural Features Inventory.

Landfire, 2017. Landfire Biophysical Settings Review Site. Accessed May, 2017  
<http://www.landfirereview.org/descriptions.html>.

National Ocean Service, 2017. Tides and Currents (historic water level data for US coastal waters).  
<https://tidesandcurrents.noaa.gov/stations.html?type=Water+Levels>

NDBC, 2017. National Data Buoy Center (wave height and period data for US coastal waters). NOAA.  
<http://www.ndbc.noaa.gov/>

PRISM Climate Group. 2013. Gridded 30 Year Normals, 1981-2010. Oregon State University,  
<http://prism.oregonstate.edu>

U.S. Department of the Interior, Geological Survey, 2011. LANDFIRE: LANDFIRE 1.1.0 Existing Vegetation Type layer. <http://landfire.cr.usgs.gov/viewer/>

USFS, Witness Tree data for northern Lower Michigan.

## Contributors

## Approval

Nels Barrett, 10/03/2019

## Acknowledgments

The following individuals made substantive comments regarding the development of the Provisional Ecological Sites: Randy Swaty, The Nature Conservancy; Trevor Hobbs, USFS; Richard A. Corner, USFS; Andy Henriksen, NRCS; Dan Zay, NRCS.

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 
17. **Perennial plant reproductive capability:**

---