

Ecological site R104XY014IA Ponded Floodplain Marsh

Last updated: 5/18/2020
Accessed: 05/05/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 104X–Eastern Iowa and Minnesota Till Prairies

The Eastern Iowa and Minnesota Till Prairies (MLRA 104) includes the Iowan Surface, Oak Savanna, and Western Coulee and Ridges landforms (Prior 1991; MDNR 2005; WDNR 2015). It spans three states (Iowa, 74 percent; Minnesota, 22 percent; Wisconsin, 4 percent), encompassing approximately 9,660 square miles (Figure 1). The elevation ranges from approximately 1,310 feet above sea level (ASL) on the highest ridges to about 985 feet ASL in the lowest valleys. Local relief is mainly 10 to 20 feet. Glacial till and outwash deposits cover the uplands of the MLRA with recent alluvium located in the major river valleys. Paleozoic bedrock sediments, comprised primarily of shale and limestone, lies beneath the glacial material. The depth to limestone is shallow, resulting in karst topography across much of the area (USDA-NRCS 2006).

The vegetation in the MLRA has undergone drastic changes over time. Spruce forests dominated the landscape 30,000 to 21,500 years ago. As the last glacial maximum peaked 21,500 to 16,000 years ago, they were replaced with open tundras and parklands. The end of the Pleistocene Epoch saw a warming climate that initially prompted the return of spruce forests, but as the warming continued, spruce trees were replaced by deciduous trees (Baker et al. 1990). Not until approximately 9,000 years ago did the vegetation transition to prairies as climatic conditions continued to warm and subsequently dry. Between 4,000 and 3,000 years ago, oak savannas began intermingling within the prairie landscape, while the more wooded and forested areas maintained a foothold in sheltered areas. This prairie-forest transition ecosystem formed the dominant landscapes until the arrival of European settlers (Baker

et al. 1992).

Classification relationships

USFS Subregions: North Central U.S. Driftless and Escarpment (222L), Minnesota and Northeast Iowa Morainal-Oak Savannah (222M), Central Dissected Till Plains (251C) Sections; Menominee Eroded Pre-Wisconsin Till (222La), Oak Savannah Till and Loess Plains (222Me), Southeast Iowa Rolling Loess Hills (251Ch) Subsections (Cleland et al. 2007)

U.S. EPA Level IV Ecoregion: Eastern Iowa and Minnesota Drift Plains (47c), Rolling Loess Prairies (47f), Lower St. Croix and Vermillion Valleys (47g), Rochester/Paleozoic Plateau Upland (52c) (USEPA 2013)

National Vegetation Classification – Ecological Systems: North-Central Interior Floodplain (CES202.694) (NatureServe 2015)

National Vegetation Classification - Plant Associations: Schoenoplectus fluviatilis – Schoenoplectus spp. Marsh (CEGL002221) (Nature Serve 2015)

Biophysical Settings: Central Interior and Appalachian Floodplain Systems (BpS 4214710) (LANDFIRE 2009)

Natural Resources Conservation Service – Iowa Plant Community Species List: Marsh, River Bulrush; Marsh, Bulrush- Cattail – Bur-reed Shallow (USDA-NRCS 2007)

Iowa Department of Natural Resources: Floodplain Marsh (INAI 1984)

Minnesota Department of Natural Resources: MRp93 Prairie Bulrush – Arrowhead Marsh (MDNR 2005)

U.S. Army Corps of Engineers: Marshes (Eggers and Reed 2015)

Ecological site concept

Ponded Floodplain Marshes are located within the green areas on the map (Figure 1). They occur on floodplains in river valleys. The soils are Entisols and Mollisols that are very poorly to poorly-drained and deep, formed in alluvium. The site experiences seasonal flooding and subsequent ponding for a significant portion of the growing season.

The historic pre-European settlement vegetation on this ecological site was dominated by emergent herbaceous vegetation adapted to flooded and saturated conditions. River bulrush (*Bolboschoenus fluviatilis* (Torr.) Soják) and broadleaf arrowhead (*Sagittaria latifolia* Willd.) are the dominant and diagnostic species for the site, respectively. Woolgrass (*Scirpus cyperinus* (L.) Kunth), broadfruit bur-reed (*Sparganium eurycarpum* Engelm.), and common bladderwort (*Utricularia macrorhiza* Leconte) are other common emergent associates. Herbaceous species typical of an undisturbed plant community associated with this ecological site can include sweetflag (*Acorus americanus* (Raf.) Raf), threepetal bedstraw (*Galium trifidum* L.), northern bog bedstraw (*Galium labradoricum* (Wiegand) Wiegand), and hairy sedge (*Carex lacustris* L.) (Drobney et al. 2001). Depth and duration of flooding are the primary disturbance factors that maintain this ecological site, while native mammal herbivory is a secondary factor (LANDFIRE 2009; Eggers and Reed 2015).

Associated sites

R104XY016IA	Wet Terrace Sedge Meadow Alluvial parent material on low stream terraces that experiences rare flooding including Bremer, Harcot, Marshan, Selmass, Talcot, and Udolpho
R104XY018IA	Wet Floodplain Sedge Meadow Alluvial parent material on floodplains that experiences occasional flooding including Calco, Coland, Colo, Sawmill, Udifluvents, and Zook

F104XY019IA	Floodplain Swamp Forest Alluvial parent material on floodplains that experiences occasional flooding including Calco, Coland, Colo, Kalmarville, Sawmill, Udifluvents, and Zook
-------------	---

Similar sites

R104XY018IA	Wet Floodplain Sedge Meadow Wet Floodplain Sedge Meadows experience flooding only and are dominated by a sedge meadow plant community
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Bolboschoenus fluviatilis</i> (2) <i>Sagittaria latifolia</i>

Physiographic features

Ponded Floodplain Marshes occur on floodplains in river valleys (Figure 2). They are situated on elevations ranging from approximately 499 to 1401 feet ASL. The site experiences occasional to frequent flooding and frequent ponding that can last from 2 to 30 days at a time. Ponding depths on average are about 12 inches (Table 1).




Figure 2. Figure 1. Location of Pondered Floodplain Marsh ecological site within MLRA 104.




Figure 3. Figure 2. Representative block diagram of Pondered Floodplain Marsh and associated ecological sites.

Table 2. Representative physiographic features

Slope shape across	(1) Concave (2) Linear
Slope shape up-down	(1) Concave (2) Linear
Landforms	(1) River valley > Flood plain
Runoff class	Negligible to low
Flooding duration	Long (7 to 30 days)
Flooding frequency	Rare to frequent
Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	Occasional to frequent
Elevation	152–457 m
Slope	0–2%
Ponding depth	0–30 cm
Water table depth	0–15 cm
Aspect	Aspect is not a significant factor

Climatic features

The Eastern Iowa and Minnesota Till Prairies falls into the hot-summer humid continental climate (Dfa) and warm-summer humid continental climate (Dfb) Köppen-Geiger climate classifications (Peel et al. 2007). In winter, dry, cold air masses periodically shift south from Canada. As these air masses collide with humid air, snowfall and rainfall result. In summer, moist, warm air masses from the Gulf of Mexico migrate north, producing significant frontal or convective rains. Occasionally, hot, dry winds originating from the Desert Southwest will stagnate over the region, creating extended droughty periods in the summer from unusually high temperatures. Air masses from the Pacific Ocean can also spread into the region and dominate producing mild, dry weather in the autumn known as Indian Summers (NCDC 2006).

The soil temperature regime of MLRA 104 is classified as mesic, where the mean annual soil temperature is between 46 and 59°F (USDA-NRCS 2006). Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south one travels. The average freeze-free period of this ecological site is about 154 days, while the frost-free period is about 127 days (Table 2). The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season. Average annual precipitation is approximately 36 inches, which includes rainfall plus the water equivalent from snowfall (Table 3). The average annual low and high temperatures are 36 and 57°F, respectively.

Climate data and analyses are derived from 30-year averages gathered from six National Oceanic and Atmospheric Administration (NOAA) weather stations contained within the range of this ecological site (Table 4).

Table 3. Representative climatic features

Frost-free period (characteristic range)	119-135 days
Freeze-free period (characteristic range)	141-167 days
Precipitation total (characteristic range)	889-965 mm
Frost-free period (actual range)	115-137 days
Freeze-free period (actual range)	139-169 days
Precipitation total (actual range)	864-965 mm
Frost-free period (average)	127 days
Freeze-free period (average)	154 days
Precipitation total (average)	914 mm

Climate stations used

- (1) BYRON 4NORTH [USC00211174], Byron, MN
- (2) AUSTIN WWT FAC [USC00210355], Austin, MN
- (3) OSAGE [USC00136305], Osage, IA
- (4) TRIPOLI [USC00138339], Tripoli, IA
- (5) ANAMOSA 1 WNW [USC00130213], Anamosa, IA
- (6) CEDAR RAPIDS NO 1 [USC00131319], Marion, IA

Influencing water features

Ponded Floodplain Marshes are classified as a RIVERINE: flooded, ponded, herbaceous wetland under the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008) and as a Palustrine, Persistent, Emergent, Seasonally Flooded-Saturated wetland under the National Wetlands Inventory (FGDC 2013). Overbank flow and subsurface hydraulic connections are the main sources of water for this ecological site, but other sources may be from surface runoff from adjacent uplands and precipitation (Smith et al. 1995). Infiltration is very slow (Hydrologic Group D) for undrained soils, and surface runoff is negligible to low (Figure 5).

Primary wetland hydrology indicators for an intact Ponded Floodplain Marsh may include: A1 Surface water, A2 High water table, A3 Saturation, and B14 True aquatic plants. Secondary wetland hydrology indicators may include: B10: Drainage patterns, C2 Dry-season water table, D2 Geomorphic position, and D5 FAC-neutral test (USACE 2010).




Figure 10. Figure 5. Hydrologic cycling in Ponded Floodplain Marsh ecological site.

Soil features

Soils of Ponded Floodplain Marshes are in the Entisols and Mollisols orders, further classified as Aeric Fluvaquents, Cumulic Endoaquolls, Typic Endoaquolls, and Cumulic Epiaquolls with very slow infiltration and negligible to low runoff potential. The soil series associated with this site includes Aquent, Aquolls, Epsom, Fluvaquents, Granby, and Shandep. The parent material is alluvium, and the soils are very poorly to poorly-drained and deep with seasonal high-water tables. Soil pH classes are moderately acid to moderately alkaline. No rooting restrictions are noted for the soils of this ecological site (Table 5).

Some soil map units in this ecological site, if not drained, may meet the definition of hydric soils and are listed as meeting criteria 2, 3, and 4 of the hydric soils list (77 FR 12234).




Figure 11. Figure 6. Profile sketch of soil series associated with Poned Floodplain Marsh.

Table 4. Representative soil features

Parent material	(1) Alluvium
Family particle size	(1) Fine-silty (2) Fine-loamy
Drainage class	Very poorly drained to poorly drained
Permeability class	Very slow to moderately slow
Depth to restrictive layer	203 cm
Soil depth	203 cm

Ecological dynamics

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

The MLRA lies within the transition zone between the eastern deciduous forests and the tallgrass prairies. The heterogeneous topography of the area results in variable microclimates and fuel matrices that in turn support prairies, savannas, woodlands, and forests. Poned Floodplain Marshes form an aspect of this vegetative continuum. This ecological site occurs on very poorly to poorly-drained soils. Species characteristic of this ecological site consist of hydrophytic and aquatic herbaceous vegetation.

Flooding is the dominant disturbance factor in Poned Floodplain Marshes (LANDFIRE 2009). Seasonal flooding likely occurred annually from spring snow melt and heavy rains. The depth and duration of ponded water affects species diversity, composition, and productivity. Little to no ponded water allows more of a sedge meadow community to dominate, while deep water depths create a deep marsh community populated with emergent and aquatic vegetation.

Animal herbivory also played a role in shaping this ecological site. Foraging muskrats can alter the extent of emergent vegetation, creating larger patches of open water. Left unchecked, muskrats can remove all the emergent vegetation, which won't re-establish until the next drought or drawdown event (Eggers and Reed 2015).

Today, Poned Floodplain Marshes have been greatly reduced as the land has mostly been converted for agricultural production. Remnants that do exist show evidence of indirect anthropogenic influences from hydrological alterations as non-native species have replaced the natural vegetation. A return to the historic plant community may not be possible due to significant hydrologic and water quality changes in the watershed, but long-term conservation agriculture or habitat reconstruction efforts can help to restore some natural diversity and ecological function. The state-and-transition model that follows provides a detailed description of each state,

community phase, pathway, and transition. This model is based on available experimental research, field observations, literature reviews, professional consensus, and interpretations.

STATE 1 – REFERENCE STATE

The reference plant community is categorized as a marsh community, dominated by hydrophytic and aquatic vegetation. The three community phases within the reference state are dependent on seasonal flooding and subsequent ponding. The depth and duration of ponding alters species composition, cover, and extent. Animal herbivory has more localized impacts in the reference phases, but does contribute to overall species composition, diversity, cover, and productivity.

Community Phase 1.1 River Bulrush – Broadleaf Arrowhead – Sites in this reference community phase are dominated by hydrophytic herbaceous vegetation. River bulrush and broadleaf arrowhead are the dominant species. Some sites may be dominated by other bulrushes, such as woolrush, stalked bulrush (*Scirpus pedicellatus* Fernald), and slender bulrush (*Schoenoplectus heterochaetus* (Chase) Soják). Characteristic forbs can include broadfruit bur-reed, common bladderwort, and water knotweed (*Polygonum amphibium* L.). Water depths between 1 and 2 feet will maintain this phase, but a reduced water level (below 1 foot) will shift the community to phase 1.2 while an increase in water level (above 2 feet) will shift the community to phase 1.3.

Pathway 1.1A – Ponded water depths decrease to <12 inches.

Pathway 1.1B – Ponded water depths increase to >24 inches.

Community Phase 1.2 Hairy Sedge – Broadleaf Arrowhead – This reference community phase can occur when the frequency and depth of ponding are reduced to less than 1 foot. Bulrushes are still present, but hairy sedge becomes the dominant monocot on the site. Broadleaf arrowhead is still the dominant forb, but forb diversity increases and can include smallspike false nettle (*Boehmeria cylindrica* (L.) Sw.), marsh skullcap (*Scutellaria galericulata* L.), and tufted loosestrife (*Lysimachia thyrsiflora* L.). Shallow ponded water depths (less than 1 foot) will maintain this phase, but an increase in water depths can shift the community back to phase 1.1.

Pathway 1.2A – Ponded water depths increase to 12 – 24 inches.

Community Phase 1.3 River Bulrush – Common Duckweed – This reference community phase can occur when the frequency and depth of ponding are greater than 2 feet. Bulrushes and cattails (*Typha* L.) are the dominant monocots, and common duckweed (*Lemna minor* L.) is the dominant floating-leaved vegetation. Aquatic vegetation becomes important characteristic species during this phase and can include species such as coon's tail (*Ceratophyllum demersum* L.), common duckmeat (*Spirodela polyrrhiza* (L.) Schleid), and American white waterlily (*Nymphaea odorata* Aiton ssp. *tuberosa* (Paine) Wiersma & Hellquist). Deep ponded water depths (greater than 2 feet) will maintain this phase, but a decrease in water depths can shift the community back to phase 1.1.

Pathway 1.3A – Ponded water depths decrease to 12 – 24 inches.

Transition 1A – Direct and indirect alterations to the landscape hydrology from human-induced land development transition the site to the hydrologically-altered state (2).

Transition 1B – Cultural treatments to enhance forage quality and yield transitions this site to the forage state (3).

Transition 1C – Installation of drain tiles, tillage, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

STATE 2 – HYDROLOGICALLY-ALTERED STATE

Hydrology is the most important determinant of wetlands and wetland processes. Hydrology modifies and determines the physiochemical environment (i.e., sediments, soil chemistry, water chemistry) which in turn directly affects the vegetation, animals, and microbes (Mitsch and Gosselink 2007). Human activities on landscape hydrology have greatly altered Pondered Floodplain Marshes. Alterations such as agricultural tile draining and conversion to cropland on adjacent lands in addition to stream channelization and damming have changed the natural hydroperiod and rate of sedimentation as well as increased nutrient pollution (Mitsch and Gosselink 2007; Eggers and Reed 2015).

Community Phase 2.1 Reed Canarygrass – River Bulrush – This community phase represents the early changes to the natural wetland hydroperiod, increasing sedimentation, and unabated nutrient runoff. Native monocots, such as river bulrush, softstem bulrush, and cattails, continue to form a component of the herbaceous layer, but the highly invasive reed canarygrass (*Phalaris arundinacea* L.) co-dominates (Waggy 2010). As reed canarygrass invades, it can not only alter species composition, but vegetation structure as well (Annen et al. 2008).

Pathway 2.1A – Continuing alterations to the natural hydrology and increasing sedimentation

Community Phase 2.2 Reed Canarygrass – Sites falling into this community phase have experienced significant sedimentation and nutrient enrichment and are dominated by a monoculture of reed canarygrass (Eggers and Reed 2015). Reed canarygrass stands can significantly alter the physiochemical environment as well as the biotic communities, making the site only suitable to reed canarygrass. These monotypic stands create a positive feedback loop that perpetuates increasing sedimentation, altered hydrology, and dominance by this non-native species, especially in sites affected by nutrient enrichment from agricultural runoff (Vitousek 1995; Bernard and Lauve 1995; Kercher et al. 2007; Waggy 2010; Eggers and Reed 2015).

Transition 2A – Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

Transition 2B – Installation of drain tiles, tillage, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

Restoration 2A – Hydroperiod restoration, site preparation, non-native species control, and seeding native species transition the site to the reconstructed marsh state (5).

STATE 3 – FORAGE STATE

The forage state arises when the site is converted to a farming system that emphasizes domestic livestock production, known as grassland agriculture. Periodic cultural treatments (e.g., clipping, drainage, soil amendment applications, planting new species and/or cultivars, mechanical harvesting), hydrologic alterations and/or grazing by domesticated livestock transition and maintain this state (USDA-NRCS 2003). Early settlers seeded non-native species, such as smooth brome (*Bromus inermis* Leyss.) and Kentucky bluegrass (*Poa pratensis* L.), to help extend the grazing season (Smith 1998). Over time, as lands were continuously harvested or grazed by herds of cattle, these species were able to spread and expand across the landscape, reducing the native species diversity and ecological function.

Community Phase 3.1 Hayfield – Sites in this community phase consist of forage plants that are planted and mechanically harvested. Mechanical harvesting removes much of the aboveground biomass and nutrients that feed the soil microorganisms (Franzluebbers et al. 2000; USDA-NRCS 2003). As a result, soil biology is reduced leading to decreases in nutrient uptake by plants, soil organic matter, and soil aggregation. Frequent biomass removal can also reduce the site's carbon sequestration capacity (Skinner 2008).

Pathway 3.1A – Mechanical harvesting is replaced with domestic livestock utilizing continuous grazing.

Pathway 3.1B – Mechanical harvesting is replaced with domestic livestock utilizing rotational grazing.

Community Phase 3.2 Continuous Pastured Grazing System – This community phase is characterized by continuous grazing where domestic livestock graze a pasture for the entire season. Depending on stocking density, this can result in lower forage quality and productivity, weed invasions, and uneven pasture use. Continuous grazing can also increase the amount of bare ground and erosion and reduce soil organic matter, cation exchange capacity, water-holding capacity, and nutrient availability and retention (Bharati et al. 2002; Leake et al. 2004; Teague et al. 2011). Smooth brome, Kentucky bluegrass, and white clover (*Trifolium repens* L.) are common pasture species used in this phase. Their tolerance to continuous grazing has allowed these species to dominate, sometimes completely excluding the native vegetation.

Pathway 3.2A – Domestic livestock are removed, and mechanical harvesting is implemented.

Pathway 3.2B – Rotational grazing replaces continuous grazing.

Community Phase 3.3 Rest-Rotation Pastured Grazing System – This community phase is characterized by

rotational grazing where the pasture has been subdivided into several smaller paddocks. Through the development of a grazing plan, livestock utilize one or a few paddocks, while the remaining area is rested allowing plants to restore vigor and energy reserves, deepen root systems, develop seeds, as well as allow seedling establishment (Undersander et al. 2002; USDA-NRCS 2003). Rest-rotation pastured grazing systems include deferred rotation, rest rotation, high intensity – low frequency, and short duration methods. Vegetation is generally more diverse and can include orchardgrass (*Dactylis glomerata* L.), timothy (*Phleum pratense* L.), red clover (*Trifolium pratense* L.), and alfalfa (*Medicago sativa* L.). The addition of native prairie species can further bolster plant diversity and, in turn, soil function. This community phase promotes numerous ecosystem benefits including increasing biodiversity, preventing soil erosion, maintaining and enhancing soil quality, sequestering atmospheric carbon, and improving water yield and quality (USDA-NRCS 2003).

Pathway 3.3A – Continuous grazing replaces rotational grazing.

Pathway 3.3B – Domestic livestock are removed, and mechanical harvesting is implemented.

Transition 3A – Land is abandoned and left fallow; natural succession by opportunistic species transition this site the hydrologically-altered state (2).

Transition 3B – Installation of drain tiles, tillage, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

Restoration 3A – Hydroperiod restoration, site preparation, non-native species control, and seeding native species transition the site to the reconstructed marsh state (5).

STATE 4 – CROPLAND STATE

The low topographic relief across the MLRA has resulted in nearly the entire area being converted to agriculture (Eilers and Roosa 1994). Agricultural tile drains used to lower the water table and the continuous use of tillage, row-crop planting, and chemicals (i.e., herbicides, fertilizers, etc.) have effectively eliminated the reference community and many of its natural ecological functions in favor of crop production. Corn and soybeans are the dominant crops for the site, and oats (*Avena* L.) and alfalfa (*Medicago sativa* L.) may be rotated periodically. These areas are likely to remain in crop production for the foreseeable future.

Community Phase 4.1 Conventional Tillage Field – Sites in this community phase typically consist of monoculture row-cropping maintained by conventional tillage practices. They are cropped in either continuous corn or corn-soybean rotations. The frequent use of deep tillage, low crop diversity, and bare soil conditions during the non-growing season negatively impacts soil health. Under these practices, soil aggregation is reduced or destroyed, soil organic matter is reduced, erosion and runoff are increased, and infiltration is decreased, which can ultimately lead to undesirable changes in the hydrology of the watershed (Tomer et al. 2005).

Pathway 4.1A – Tillage operations are greatly reduced, crop rotation occurs on a regular interval, and crop residue remains on the soil surface.

Pathway 4.1B – Tillage operations are greatly reduced or eliminated, crop rotation occurs on a regular interval, crop residue remains on the soil surface, and cover crops are planted following crop harvest.

Community Phase 4.2 Conservation Tillage Field – This community phase is characterized by rotational crop production that utilizes various conservation tillage methods to promote soil health and reduce erosion. Conservation tillage methods include strip-till, ridge-till, vertical-till, or no-till planting systems. Strip-till keeps seedbed preparation to narrow bands less than one-third the width of the row where crop residue and soil consolidation are left undisturbed in-between seedbed areas. Strip-till planting may be completed in the fall and nutrient application either occurs simultaneously or at the time of planting. Ridge-till uses specialized equipment to create ridges in the seedbed and vegetative residue is left on the surface in between the ridges. Weeds are controlled with herbicides and/or cultivation, seedbed ridges are rebuilt during cultivation, and soils are left undisturbed from harvest to planting. Vertical-till systems employ machinery that lightly tills the soil and cuts up crop residue, mixing some of the residue into the top few inches of the soil while leaving a large portion on the surface. No-till management is the most conservative, disturbing soils only at the time of planting and fertilizer application. Compared to conventional tillage systems, conservation tillage methods can improve soil ecosystem function by reducing soil erosion, increasing organic matter and water availability, improving water quality, and reducing soil

compaction.

Pathway 4.2A – Intensive tillage is utilized, and monoculture row-cropping is established.

Pathway 4.2B – Cover crops are implemented to minimize soil erosion.

Community Phase 4.3 Conservation Tillage with Cover Crop Field – This community phase applies conservation tillage methods as described above as well as adds cover crop practices. Cover crops typically include nitrogen-fixing species (e.g., legumes), small grains (e.g., rye, wheat, oats), or forage covers (e.g., turnips, radishes, rapeseed). The addition of cover crops not only adds plant diversity but also promotes soil health by reducing soil erosion, limiting nitrogen leaching, suppressing weeds, increasing soil organic matter, and improving the overall soil ecosystem. In the case of small grain cover crops, surface cover and water infiltration are increased, while forage covers can be used to graze livestock or support local wildlife. Of the three community phases for this state, this phase promotes the greatest soil sustainability and improves ecological functioning within a cropland system.

Pathway 4.3A – Cover crop practices are abandoned.

Pathway 4.3B – Intensive tillage is utilized, cover crops practices are abandoned, monoculture row-cropping is established, and crop rotation is reduced or eliminated.

Transition 4A – Agricultural production abandoned and left fallow; natural succession by opportunistic species transition this site to the hydrologically-altered state (2).

Transition 4B – Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

Restoration 4A – Hydroperiod restoration, site preparation, non-native species control and seeding native species transition this site to the reconstructed marsh state (5).

STATE 5 – RECONSTRUCTED MARSH STATE

Marsh habitats provide multiple ecosystem services including flood abatement, water quality improvement, and biodiversity support (Mitsch and Gosselink 2007). However, many marsh communities have been eliminated as a result of type conversions to agricultural production, changes to the natural hydrologic regime, and invasion of non-native species, thereby significantly reducing these services (Annen et al. 2008). The extensive alterations of lands adjacent to Pondered Floodplain Marshes may not allow for restoration back to the historic reference condition. But ecological reconstruction can aim to aid the recovery of degraded, damaged, or destroyed functions. A successful reconstruction will have the ability to structurally and functionally sustain itself, demonstrate resilience to the natural ranges of stress and disturbance, and create and maintain positive biotic and abiotic interactions (SER 2002; Mitsch and Jørgensen 2004).

Community Phase 5.1 Early Successional Reconstructed Marsh – This community phase represents the early community assembly from marsh habitat reconstruction and is highly dependent on invasive species control, hydroperiod repair, and planting (Adams and Galatowitsch 2006). In addition, adaptive restoration tactics that incorporate multiple restoration methods should be implemented to more clearly identify cause-effect relationships of vegetative development (Zedler 2005).

Pathway 5.1A – Maintenance of proper hydrology and nutrient balances in line with a developed wetland management plan.

Community Phase 5.2 Late Successional Reconstructed Marsh – Appropriately timed disturbance regimes (e.g. hydroperiod, invasive species control) and nutrient management applied to the early successional community phase can help increase the species richness and improve ecosystem function, pushing the site into a late successional community phase over time (Mitsch and Gosselink 2007).

Pathway 5.2A – Reconstruction experiences a setback from extreme weather event or improper timing of management actions.


Transition 5A – Land is abandoned and left fallow; natural succession by opportunistic species transition this site the hydrologically-altered state (2).

Transition 5B – Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

Transition 5C – Installation of drain tiles, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

State and transition model

R104XY014IA PONDED FLOODPLAIN MARSH



Code	Process
1.1A	Ponded water depths <12 inches
1.1B	Ponded water depths >24 inches
1.2A, 1.3A	Ponded water depths 12-24 inches
T1A, T3A, T4A, T5A	Changes to natural hydroperiod and/or land abandonment
2.1A	Increasing changes to hydrology and increasing sedimentation
T1B, T2A, T4B, T5B	Cultural treatments are implemented to increase forage quality and yield
3.1A	Mechanical harvesting is replaced with domestic livestock and continuous grazing
3.1B	Mechanical harvesting is replaced with domestic livestock and rest-rotational grazing
3.2A, 3.3B	Tillage, forage crop planting, and mechanical harvesting replace grazing
3.2B	Implementation of rest-rotational grazing
3.3A	Implementation of continuous grazing
T1C, T2B, T3B, T5C	Agricultural conversion via drainage, tillage, seeding, and non-selective herbicide
4.1A	Less tillage, residue management
4.1B	Less tillage, residue management, and implementation of cover cropping
4.2B	Implementation of cover cropping
4.2A, 4.3B	Intensive tillage, remove residue, and reinitiate monoculture row cropping
4.3A	Remove cover cropping
R2A, R3A, R4A	Site preparation, non-native species control, hydroperiod repair, and native seeding
5.1A	Invasive species control and implementation of disturbance regimes
5.2A	Drought or improper timing/use of management actions

State 1

Reference State

The reference plant community is categorized as a marsh community, dominated by hydrophytic and aquatic vegetation. The three community phases within the reference state are dependent on seasonal flooding and subsequent ponding. The depth and duration of ponding alters species composition, cover, and extent. Animal herbivory has more localized impacts in the reference phases, but does contribute to overall species composition, diversity, cover, and productivity.

Community 1.1

River Bulrush – Broadleaf Arrowhead

Sites in this reference community phase are dominated by hydrophytic herbaceous vegetation. River bulrush and broadleaf arrowhead are the dominant species. Some sites may be dominated by other bulrushes, such as woolrush, stalked bulrush (*Scirpus pedicellatus* Fernald), and slender bulrush (*Schoenoplectus heterochaetus* (Chase) Soják). Characteristic forbs can include broadfruit bur-reed, common bladderwort, and water knotweed (*Polygonum amphibium* L.). Water depths between 1 and 2 feet will maintain this phase, but a reduced water level (below 1 foot) will shift the community to phase 1.2 while an increase in water level (above 2 feet) will shift the community to phase 1.3.

Community 1.2

Hairy Sedge – Broadleaf Arrowhead

This reference community phase can occur when the frequency and depth of ponding are reduced to less than 1 foot. Bulrushes are still present, but hairy sedge becomes the dominant monocot on the site. Broadleaf arrowhead is still the dominant forb, but forb diversity increases and can include smallspike false nettle (*Boehmeria cylindrica* (L.) Sw.), marsh skullcap (*Scutellaria galericulata* L.), and tufted loosestrife (*Lysimachia thyrsiflora* L.). Shallow ponded water depths (less than 1 foot) will maintain this phase, but an increase in water depths can shift the community back to phase 1.1.

Community 1.3

River Bulrush – Common Duckweed

This reference community phase can occur when the frequency and depth of ponding are greater than 2 feet. Bulrushes and cattails (*Typha* L.) are the dominant monocots, and common duckweed (*Lemna minor* L.) is the dominant floating-leaved vegetation. Aquatic vegetation becomes important characteristic species during this phase and can include species such as coon's tail (*Ceratophyllum demersum* L.), common duckmeat (*Spirodela polyrrhiza* (L.) Schleid), and American white waterlily (*Nymphaea odorata* Aiton ssp. *tuberosa* (Paine) Wiersma & Hellquist). Deep ponded water depths (greater than 2 feet) will maintain this phase, but a decrease in water depths can shift the community back to phase 1.1.

Pathway 1.1A

Community 1.1 to 1.2

Ponded water depths decrease to <12 inches.

Pathway 1.1B

Community 1.1 to 1.3

Ponded water depths increase to >24 inches.

Pathway 1.2A

Community 1.2 to 1.1

Ponded water depths increase to 12 – 24 inches.

Pathway 1.3A

Community 1.3 to 1.1

Ponded water depths decrease to 12 – 24 inches.

State 2

Hydrologically-altered State

Hydrology is the most important determinant of wetlands and wetland processes. Hydrology modifies and determines the physiochemical environment (i.e., sediments, soil chemistry, water chemistry) which in turn directly affects the vegetation, animals, and microbes (Mitsch and Gosselink 2007). Human activities on landscape hydrology have greatly altered Ponded Floodplain Marshes. Alterations such as agricultural tile draining and conversion to cropland on adjacent lands in addition to stream channelization and damming have changed the natural hydroperiod and rate of sedimentation as well as increased nutrient pollution (Mitsch and Gosselink 2007; Eggers and Reed 2015).

Community 2.1

Reed Canarygrass – River Bulrush

This community phase represents the early changes to the natural wetland hydroperiod, increasing sedimentation, and unabated nutrient runoff. Native monocots, such as river bulrush, softstem bulrush, and cattails, continue to form a component of the herbaceous layer, but the highly invasive reed canarygrass (*Phalaris arundinacea* L.) co-dominates (Waggy 2010). As reed canarygrass invades, it can not only alter species composition, but vegetation structure as well (Annen et al. 2008).

Community 2.2

Reed Canarygrass

Sites falling into this community phase have experienced significant sedimentation and nutrient enrichment and are dominated by a monoculture of reed canarygrass (Eggers and Reed 2015). Reed canarygrass stands can significantly alter the physiochemical environment as well as the biotic communities, making the site only suitable to reed canarygrass. These monotypic stands create a positive feedback loop that perpetuates increasing sedimentation, altered hydrology, and dominance by this non-native species, especially in sites affected by nutrient enrichment from agricultural runoff (Vitousek 1995; Bernard and Lauve 1995; Kercher et al. 2007; Waggy 2010; Eggers and Reed 2015).

Pathway 2.1A

Community 2.1 to 2.2

Continuing alterations to the natural hydrology and increasing sedimentation

State 3

Forage State

The forage state arises when the site is converted to a farming operation that emphasizes domestic livestock production, known as grassland agriculture. Periodic cultural treatments (e.g., clipping, drainage, soil amendment applications, planting new species and/or cultivars, mechanical harvesting), hydrologic alterations and/or grazing by domesticated livestock transition and maintain this state (USDA-NRCS 2003). Early settlers seeded non-native species, such as smooth brome (*Bromus inermis* Leyss.) and Kentucky bluegrass (*Poa pratensis* L.), to help extend the grazing season (Smith 1998). Over time, as lands were continuously harvested or grazed by herds of cattle, these species were able to spread and expand across the landscape, reducing the native species diversity and ecological function.

Community 3.1

Hayfield

Sites in this community phase consist of forage plants that are planted and mechanically harvested. Mechanical

harvesting removes much of the aboveground biomass and nutrients that feed the soil microorganisms (Franzluebbers et al. 2000; USDA-NRCS 2003). As a result, soil biology is reduced leading to decreases in nutrient uptake by plants, soil organic matter, and soil aggregation. Frequent biomass removal can also reduce the site's carbon sequestration capacity (Skinner 2008).

Community 3.2

Continuous Pastured Grazing

This community phase is characterized by continuous grazing where domestic livestock graze a pasture for the entire season. Depending on stocking density, this can result in lower forage quality and productivity, weed invasions, and uneven pasture use. Continuous grazing can also increase the amount of bare ground and erosion and reduce soil organic matter, cation exchange capacity, water-holding capacity, and nutrient availability and retention (Bharati et al. 2002; Leake et al. 2004; Teague et al. 2011). Smooth brome, Kentucky bluegrass, and white clover (*Trifolium repens* L.) are common pasture species used in this phase. Their tolerance to continuous grazing has allowed these species to dominate, sometimes completely excluding the native vegetation.

Community 3.3

Periodic-rest Pastured Grazing

This community phase is characterized by periodic-rest grazing where the pasture has been subdivided into several smaller paddocks. Subdividing the pasture in this way allows livestock to utilize one or a few paddocks, while the remaining area is rested allowing plants to restore vigor and energy reserves, deepen root systems, develop seeds, as well as allow seedling establishment (Undersander et al. 2002; USDA-NRCS 2003). Periodic-rest pastured grazing includes deferred periods, rest periods, and periods of high intensity – low frequency, and short duration methods. Vegetation is generally more diverse and can include orchardgrass (*Dactylis glomerata* L.), timothy (*Phleum pratense* L.), red clover (*Trifolium pratense* L.), and alfalfa (*Medicago sativa* L.). The addition of native prairie species can further bolster plant diversity and, in turn, soil function. This community phase promotes numerous ecosystem benefits including increasing biodiversity, preventing soil erosion, maintaining and enhancing soil quality, sequestering atmospheric carbon, and improving water yield and quality (USDA-NRCS 2003).

Pathway 3.1A

Community 3.1 to 3.2

Mechanical harvesting is replaced with domestic livestock utilizing continuous grazing.

Pathway 3.1B

Community 3.1 to 3.3

Mechanical harvesting is replaced with domestic livestock utilizing periodic-rest grazing.

Pathway 3.2A

Community 3.2 to 3.1

Domestic livestock are removed, and mechanical harvesting is implemented.

Pathway 3.2B

Community 3.2 to 3.3

Periodic-rest grazing replaces continuous grazing.

Pathway 3.3B

Community 3.3 to 3.1

Domestic livestock are removed, and mechanical harvesting is implemented.

Pathway 3.3A

Community 3.3 to 3.2

Continuous grazing replaces periodic-rest grazing.

State 4

Cropland State

The low topographic relief across the MLRA has resulted in nearly the entire area being converted to agriculture (Eilers and Roosa 1994). Agricultural tile drains used to lower the water table and the continuous use of tillage, row-crop planting, and chemicals (i.e., herbicides, fertilizers, etc.) have effectively eliminated the reference community and many of its natural ecological functions in favor of crop production. Corn and soybeans are the dominant crops for the site, and oats (*Avena L.*) and alfalfa (*Medicago sativa L.*) may be rotated periodically. These areas are likely to remain in crop production for the foreseeable future.

Community 4.1

Conventional Tillage Field

Sites in this community phase typically consist of monoculture row-cropping maintained by conventional tillage practices. They are cropped in either continuous corn or alternating periods of corn and soybean crops. The frequent use of deep tillage, low crop diversity, and bare soil conditions during the non-growing season negatively impacts soil health. Under these practices, soil aggregation is reduced or destroyed, soil organic matter is reduced, erosion and runoff are increased, and infiltration is decreased, which can ultimately lead to undesirable changes in the hydrology of the watershed (Tomer et al. 2005).

Community 4.2

Conservation Tillage Field

This community phase is characterized by periodically alternating crops and utilizing various conservation tillage methods to promote soil health and reduce erosion. Conservation tillage methods include strip-till, ridge-till, vertical-till, or no-till planting operations. Strip-till keeps seedbed preparation to narrow bands less than one-third the width of the row where crop residue and soil consolidation are left undisturbed in-between seedbed areas. Strip-till planting may be completed in the fall and nutrient application either occurs simultaneously or at the time of planting. Ridge-till uses specialized equipment to create ridges in the seedbed and vegetative residue is left on the surface in between the ridges. Weeds are controlled with herbicides and/or cultivation, seedbed ridges are rebuilt during cultivation, and soils are left undisturbed from harvest to planting. Vertical-till operations employ machinery that lightly tills the soil and cuts up crop residue, mixing some of the residue into the top few inches of the soil while leaving a large portion on the surface. No-till management is the most conservative, disturbing soils only at the time of planting and fertilizer application. Compared to conventional tillage operations, conservation tillage methods can improve soil ecosystem function by reducing soil erosion, increasing organic matter and water availability, improving water quality, and reducing soil compaction.

Community 4.3

Conservation Tillage with Cover Crop Field

This community phase applies conservation tillage methods as described above as well as adds cover crop practices. Cover crops typically include nitrogen-fixing species (e.g., legumes), small grains (e.g., rye, wheat, oats), or forage covers (e.g., turnips, radishes, rapeseed). The addition of cover crops not only adds plant diversity but also promotes soil health by reducing soil erosion, limiting nitrogen leaching, suppressing weeds, increasing soil organic matter, and improving the overall soil ecosystem. In the case of small grain cover crops, surface cover and water infiltration are increased, while forage covers can be used to graze livestock or support local wildlife. Of the three community phases for this state, this phase promotes the greatest soil sustainability and improves ecological functioning within a row crop operation.

Pathway 4.1A

Community 4.1 to 4.2

Tillage operations are greatly reduced, alternating crops occurs on a regular interval, and crop residue remains on the soil surface.

Pathway 4.1B

Community 4.1 to 4.3

Tillage operations are greatly reduced or eliminated, alternating crops occurs on a regular interval, crop residue remains on the soil surface, and cover crops are planted following crop harvest.

Pathway 4.2A

Community 4.2 to 4.1

Intensive tillage is utilized, and monoculture row-cropping is established.

Pathway 4.2B

Community 4.2 to 4.3

Cover crops are implemented to minimize soil erosion.

Pathway 4.3B

Community 4.3 to 4.1

Intensive tillage is utilized, cover crops practices are abandoned, monoculture row-cropping is established on a more-or-less continuous basis.

Pathway 4.3A

Community 4.3 to 4.2

Cover crop practices are abandoned.

State 5

Reconstructed Marsh State

Marsh habitats provide multiple ecosystem services including flood abatement, water quality improvement, and biodiversity support (Mitsch and Gosselink 2007). However, many marsh communities have been eliminated as a result of type conversions to agricultural production, changes to the natural hydrologic regime, and invasion of non-native species, thereby significantly reducing these services (Annen et al. 2008). The extensive alterations of lands adjacent to Pondered Floodplain Marshes may not allow for restoration back to the historic reference condition. But ecological reconstruction can aim to aid the recovery of degraded, damaged, or destroyed functions. A successful reconstruction will have the ability to structurally and functionally sustain itself, demonstrate resilience to the natural ranges of stress and disturbance, and create and maintain positive biotic and abiotic interactions (SER 2002; Mitsch and Jørgensen 2004).

Community 5.1

Early Successional Reconstructed Marsh

This community phase represents the early community assembly from marsh habitat reconstruction and is highly dependent on invasive species control, hydroperiod repair, and planting (Adams and Galatowitsch 2006). In addition, adaptive restoration tactics that incorporate multiple restoration methods should be implemented to more clearly identify cause-effect relationships of vegetative development (Zedler 2005).

Community 5.2

Late Successional Reconstructed Marsh

Appropriately timed disturbance regimes (e.g. hydroperiod, invasive species control) and nutrient management applied to the early successional community phase can help increase the species richness and improve ecosystem function, pushing the site into a late successional community phase over time (Mitsch and Gosselink 2007).

Pathway 5.1A
Community 5.1 to 5.2

Maintenance of proper hydrology and nutrient balances in line with a developed wetland management plan.

Pathway 5.2A
Community 5.2 to 5.1

Reconstruction experiences a setback from extreme weather event or improper timing of management actions.

Transition T1A
State 1 to 2

Direct and indirect alterations to the landscape hydrology from human-induced land development transition the site to the hydrologically-altered state (2).

Transition T1B
State 1 to 3

Cultural treatments to enhance forage quality and yield transitions this site to the forage state (3).

Transition T1C
State 1 to 4

Installation of drain tiles, tillage, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

Transition T2A
State 2 to 3

Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

Transition T2B
State 2 to 4

Installation of drain tiles, tillage, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

Restoration pathway R2A
State 2 to 5

Hydroperiod restoration, site preparation, non-native species control, and seeding native species transition the site to the reconstructed marsh state (5).

Transition T3A
State 3 to 2

Land is abandoned and left fallow; natural succession by opportunistic species transition this site the hydrologically-altered state (2).

Transition T3B
State 3 to 4

Installation of drain tiles, tillage, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

Restoration pathway R3A

State 3 to 5

Hydroperiod restoration, site preparation, non-native species control, and seeding native species transition the site to the reconstructed marsh state (5).

Transition T4A

State 4 to 2

Agricultural production abandoned and left fallow; natural succession by opportunistic species transition this site to the hydrologically-altered state (2).

Transition T4B

State 4 to 3

Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

Restoration pathway R4A

State 4 to 5

Hydroperiod restoration, site preparation, non-native species control and seeding native species transition this site to the reconstructed marsh state (5).

Transition T5A

State 5 to 2

Land is abandoned and left fallow; natural succession by opportunistic species transition this site the hydrologically-altered state (2).

Transition T5B

State 5 to 3

Cultural treatments to enhance forage quality and yield transition the site to the forage state (3).

Transition T5C

State 5 to 4

Installation of drain tiles, seeding of agricultural crops, and non-selective herbicide transition the site to the cropland state (4).

Additional community tables

Inventory data references

Tier 3 Sampling Plots used to develop the reference state, community phases 1.1 and 1.2:

State County Ownership Easting Northing

Iowa Butler Big Marsh WMA – Iowa Department of Natural Resources 514739 4723158

Iowa Black Hawk Cedar Hills Sand Prairie State Preserve – The Nature Conservancy 536597 4715723

Iowa Linn Goose Pond Natural Area – Linn County Conservation Board 598148 4666650

Iowa Linn Behrens Pond & Woodland State Preserve – The Nature Conservancy 603013 4663503

Iowa Linn Chain of Lakes – Linn County Conservation Board 602345 4659136

Iowa Worth Ochee Yahola Park – Worth County Conservation Board 478294 4814762

Other references

Adams, C.R. and S.M. Galatowitsch. 2006. Increasing the effectiveness of reed canary grass (*Phalaris arundinacea*

L.) control in wet meadow restorations. *Restoration Ecology* 14: 441-451.

Annen, C.A., E.M. Kirsch, and R.W. Tyser. 2008. Reed canarygrass invasions alter succession patterns and may reduce habitat quality in wet meadows. *Ecological Restoration* 26: 190-193.

Baker, R.G., C.A. Chumbley, P.M. Witinok, and H.K. Kim. 1990. Holocene vegetational changes in eastern Iowa. *Journal of the Iowa Academy of Science* 97: 167-177.

Baker, R.G., L.J. Maher, C.A. Chumbley, and K.L. Van Zant. 1992. Patterns of Holocene environmental changes in the midwestern United States. *Quaternary Research* 37: 379-389.

Bernard, J.B. and T.E. Lauve. 1995. A comparison of growth and nutrient uptake in *Phalaris arundinacea* L. growing in a wetland and a constructed bed receiving landfill leachate. *Wetlands* 15: 176-182.

Bharati, L., K.-H. Lee, T.M. Isenhardt, and R.C. Schultz. 2002. Soil-water infiltration under crops, pasture, and established riparian buffer in Midwestern USA. *Agroforestry Systems* 56: 249-257.

Changes in Hydric Soils Database Selection Criteria. 77 Federal Register 12234 (29 February 2012), pp. 12234-12235.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Drobney, P.D., G.S. Wilhelm, D. Horton, M. Leoschke, D. Lewis, J. Pearson, D. Roosa, and D. Smith. 2001. Floristic Quality Assessment for the State of Iowa. Neal Smith National Wildlife Refuge and Ada Hayden Herbarium, Iowa State University, Ames, IA. 123 pps.

Eggers, S.D. and D.M. Reed. 2015. Wetland Plants and Plant Communities of Minnesota and Wisconsin, Version 3.2. U.S. Army Corps of Engineers, Regulatory Branch, St. Paul District. St. Paul, MN. 478 pps.

Federal Geographic Data Committee. 2013. Classification of Wetlands and Deepwater Habitats of the United States. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, D.C. 90 pps.

Franzluebbers, A.J., J.A. Stuedemann, H.H. Schomberg, and S.R. Wilkinson. 2000. Soil organic C and N pools under long-term pasture management in the Southern Piedmont USA. *Soil Biology and Biochemistry* 32:469-478.

Iowa Natural Areas Inventory [INAI]. 1984. An Inventory of Significant Natural Areas in Iowa: Two year Progress Report of the Iowa Natural Areas Inventory. Iowa Natural Areas Inventory, Iowa Department of Natural Resources, Des Moines, IA.

Franzluebbers, A.J., J.A. Stuedemann, H.H. Schomberg, and S.R. Wilkinson. 2000. Soil organic C and N pools under long-term pasture management in the Southern Piedmont USA. *Soil Biology and Biochemistry* 32:469-478.

Kercher, S.M. A. Herr-Turnoff, J.B. Zedler. 2007. Understanding invasion as a process: the case of *Phalaris arundinacea* in wet prairies. *Biological Invasions* 9: 657-665.

LANDFIRE. 2009. Biophysical Setting 4214710 Central Interior and Appalachian Floodplain Systems. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

Leake, J., D. Johnson, D. Donnelly, G. Muckle, L. Boddy, and D. Read. 2004. Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Canadian Journal of Botany* 82: 1016-1045.

Minnesota Department of Natural Resources [MDNR]. 2005. Field Guide to the Native Plant Communities of Minnesota: The Eastern Broadleaf Forest Province. Ecological Land Classification Program, Minnesota County Biological Survey, Natural Heritage and Nongame Research Program, Minnesota Department of Natural

Resources, St. Paul, MN.

Mitsch, W.J. and J.G. Gosselink. 2007. *Wetlands*, Fourth Edition. John Wiley & Sons, Inc. Hoboken, NJ. 582 pps.

Mitsch, W.J. and S.E. Jørgensen. 2004. *Ecological Engineering and Ecosystem Restoration*. John Wiley & Sons, Inc. Hoboken, NJ. 428 pps.

National Climate Data Center [NCDC]. 2006. *Climate of Iowa*. Central Region Headquarters, Climate Services Branch, National Climatic Data Center, Asheville, NC.

NatureServe. 2018. *NatureServe Explorer: An online encyclopedia of life* [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 26 November 2018).

Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.

Prior, J.C. 1991. *Landforms of Iowa*. University of Iowa Press for the Iowa Department of Natural Resources, Iowa City, IA. 153 pps.

Skinner, R.H. 2008. High biomass removal limits carbon sequestration potential of mature temperate pastures. *Journal for Environmental Quality* 37: 1319-1326.

Smith, R.D., A. Ammann, C. Bartoldus, and M.M. Brinson. 1995. *An Approach for Assessing Wetland Functions Using Hydrogeomorphic Classification, Reference Wetlands, and Functional Indices*. U.S. Army Corps of Engineers, Waterways Experiment Station, Wetlands Research Program Technical Report WRP-DE-9. 78 pps.

Smith, D.D. 1998. Iowa prairie: original extent and loss, preservation, and recovery attempts. *The Journal of the Iowa Academy of Sciences* 105: 94-108.

Teague, W.R., S.L. Dowhower, S.A. Baker, N. Haile, P.B. DeLaune, and D.M. Conover. 2011. Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems and Environment* 141: 310-322.

Tomer, M.D., D.W. Meek, and L.A. Kramer. 2005. Agricultural practices influence flow regimes of headwater streams in western Iowa. *Journal of Environmental Quality* 34:1547-1558.

Undersander, D., B. Albert, D. Cosgrove, D. Johnson, and P. Peterson. 2002. *Pastures for Profit: A Guide to Rotational Grazing (A3529)*. University of Wisconsin-Extension and University of Minnesota Extension Service. 43 pps.

U.S. Army Corps of Engineers [USACE]. 2010. *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Midwest Region (Version 2.0)*. U.S. Army Corps of Engineers, Wetlands Regulatory Assistance Program, U.S. Army Engineer Research and Development Center, Vicksburg, MS. 141 pps.

United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2003. *National Range and Pasture Handbook, Revision 1*. Grazing Lands Technology Institute. 214 pps.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*. U.S. Department of Agriculture Handbook 296. 682 pps.

United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2007. *Iowa NRCS Plant Community Species Lists*. Des Moines, IA. Available at https://www.nrcs.usda.gov/wps/portal/nrcs/detail/ia/technical/ecoscience/bio/?cid=nrcs142p2_008160. (Accessed 19 January 2018).

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2008. *Hydrogeomorphic Wetland Classification: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service*. Technical Note No. 190-8-76. Washington, D.C. 8 pps.

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>. (Accessed 1 March 2017).

Vitousek, P.M. 1995. Biological invasions and ecosystem processes: towards an integration of population biology and ecosystem studies. *Oikos* 57: 7-13.

Waggy, M.A. 2010. *Phalaris arundinacea*. In: Fire Effects Information System [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory. Available at <https://www.feis-crs.org/feis>. (Accessed 2 March 2018).

Wisconsin Department of Natural Resources (WDNR). 2015. The Ecological Landscapes of Wisconsin: An Assessment of Ecological Resources and a Guide to Planning Sustainable Management. Wisconsin Department of Natural Resources, PUB-SS-1131 2015, Madison, WI.

Zedler, J.B. 2005. Restoring wetland plant diversity: a comparison of existing and adaptive approaches. *Wetlands Ecology and Management* 13: 5-14.

Contributors

Lisa Kluesner
Ryan Dermody

Approval

Chris Tecklenburg, 5/18/2020

Acknowledgments

This project could not have been completed without the dedication and commitment from a variety of partners and staff. Team members supported the project by serving on the technical team, assisting with the development of state and community phases of the state-and-transition model, providing peer review and technical editing, and conducting quality control and quality assurance reviews.

Drake University:

Dr. Tom Rosburg, Professor of Ecology and Botany, Des Moines, IA

Iowa Department of Natural Resources:

John Pearson, Ecologist, Des Moines, IA

Greg Schmitt, Private Lands Biologist, West Union, IA

Conservation Districts of Iowa:

Sean Kluesner, Private Lands Wetland Easement Team Specialist, New Hampton, IA

LANDFIRE (The Nature Conservancy):

Randy Swaty, Ecologist, Evanston, IL

Natural Resources Conservation Service :

Rick Bednarek, Iowa State Soil Scientist, Des Moines, IA

Scott Brady, Acting Regional Ecological Site Specialist, Havre, MT

Leland Camp, Soil Scientist, Waverly, IA

Patrick Chase, Area Resource Soil Scientist, Fort Dodge, IA

Stacey Clark, Regional Ecological Site Specialist, St. Paul, MN

James Cronin, State Biologist, Des Moines, IA

Ryan Dermody, Soil Survey Leader, Waverly, IA

Tonie Endres, Senior Regional Soil Scientist, Indianapolis, IN
 Gregg Hadish, GIS Specialist, Des Moines, IA
 John Hammerly, Soil Data Quality Specialist, Indianapolis, IN
 Lisa Kluesner, Ecological Site Specialist, Waverly, IA
 Jeff Matthias, State Grassland Specialist, Des Moines, IA
 Louis Moran, Area Resource Soil Scientist, Sioux City, IA
 Kevin Norwood, Soil Survey Regional Director, Indianapolis, IN
 James Phillips, GIS Specialist, Des Moines, IA
 Neil Sass, Area Resource Soil Scientist, West Union, IA
 Jason Steele, Area Resource Soil Scientist, Fairfield, IA

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/05/2024
Approved by	Chris Tecklenburg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-
17. **Perennial plant reproductive capability:**
-