

# Ecological site R107XA208IA

## Ponded Upland Depression Sedge Meadow

Last updated: 5/21/2020  
Accessed: 05/05/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 107X—Iowa and Missouri Deep Loess Hills

The Iowa and Minnesota Loess Hills (MLRA 107A) includes the Northwest Iowa Plains, Inner Coteau, and Coteau Moraines landforms (Prior 1991; MDNR 2005). It spans two states (Iowa, 89 percent; Minnesota, 11 percent), encompassing approximately 4,470 square miles (Figure 1). The elevation ranges from approximately 1,700 feet above sea level (ASL) on the highest ridges to about 1,115 feet ASL in the lowest valleys. Local relief is mainly 10 to 100 feet. However, some valley floors can range from 80 to 200 feet, while some upland flats only range between 3 and 6 feet. The eastern half of the MLRA is underlain by Wisconsin-age till, deposited between 20,000 and 30,000 years ago and is known as the Sheldon Creek Formation. The western half is underlain by Pre-Illinoian glacial till, deposited more than 500,000 years ago and has since undergone extensive erosion and dissection. Both surfaces are covered by approximately 4 to 20 feet of loess on the hillslopes, and Holocene alluvium covers the till in the drainageways. Cretaceous bedrock, comprised of sandstone and shale, lies beneath the glacial material (USDA-NRCS 2006).

The vegetation in the MLRA has undergone drastic changes over time. Spruce forests dominated the landscape 30,000 to 21,500 years ago. As the last glacial maximum peaked 21,500 to 16,000 years ago, they were replaced with open tundras and parklands. The end of the Pleistocene Epoch saw a warming climate that initially prompted the return of spruce forests, but as the warming continued, spruce trees were replaced by deciduous trees (Baker et al. 1990). Not until approximately 9,000 years ago did the vegetation transition to prairies as climatic conditions continued to warm and subsequently dry. Between 4,000 and 3,000 years ago, oak savannas began intermingling within the prairie landscape, while the more wooded and forested areas maintained a foothold in sheltered areas. This prairie-forest transition ecosystem formed the dominant landscapes until the arrival of European settlers (Baker et al. 1992).

### Classification relationships

U.S. Forest Service Ecological Subregions: North Central Glaciated Plains (251B) Section, Outer Coteau des Prairies (251Bb), Northwest Iowa Plains (251Bd) Subsections (Cleland et al. 2007)

U.S. EPA Level IV Ecoregion: Loess Prairies (47a) (USEPA 2013)

National Vegetation Classification – Ecological System: Eastern Great Plains Wet Meadow, Prairie and Marsh (CES205.687) (NatureServe 2015)

National Vegetation Classification – Plant Associations: Carex atherodes Wet Meadow (CEGL002220)

(NatureServe 2015)

Biophysical Settings: Eastern Great Plains Wet Meadow-Prairie-Marsh (4214880) (LANDFIRE 2009)

Natural Resources Conservation Service – Iowa Plant Community Species List: Wet Meadow, Awned Sedge (USDA-NRCS 2007)

Iowa Department of Natural Resources: Prairie Marsh (INAI 1984)

Minnesota Department of Natural Resources: WMs92a Basin Meadow/Carr (MDNR 2005)

U.S. Army Corps of Engineers: Sedge Meadows (Eggers and Reed 2015)

### Ecological site concept

Ponded Upland Depression Sedge Meadows are located within the green areas on the map (Figure 1). They occur in upland depressions on slopes generally less than one percent. The soils are Mollisols that are very poorly-drained and deep, formed in loess. The site experiences occasional to frequent, brief to long ponding from precipitation, overland flow, and groundwater flow.

The historic pre-European settlement vegetation on this site was dominated by herbaceous species adapted to temporarily ponded conditions. Wheat sedge (*Carex atherodes* Spreng.) , common rivergrass (*Scolochloa festucacea* (Willd.) Link), and water knotweed (*Polygonum amphibium* L.) are the dominant species of Ponded Upland Depression Sedge Meadows. Other grasses and grass-likes that may occur include bluejoint (*Calamagrostis canadensis* (Michx.) P. Beauv), prairie cordgrass (*Spartina pectinata* Bosc ex Link), river bulrush (*Bolboschoenus fluviatilis* (Torr.) Soják), and common spikerush (*Eleocharis palustris* (L.) Roem & Schult.) (MDNR 2005; NatureServe 2015). Species typical of an undisturbed plant community associated with this ecological site include hemlock waterparsnip (*Sium suave* Walter), woollyfruit sedge (*Carex lasiocarpa* Ehrh.), and water sedge (*Carex aquatilis* Wahlenb.) (Drobney et al. 2001; MDNR 2005; NatureServe 2015). Depth and duration of ponding as well as periodic fire are the primary disturbance factors that maintain this site, while native large mammal grazing and drought are secondary factors (MDNR 2005; LANDFIRE 2009; NatureServe 2015).

### Associated sites

R107XA208IA	<b>Ponded Upland Depression Sedge Meadow</b> Loess or loamy sediments on broad upland flats that do not experience ponding, including Gillet Grove, Letri, Marcus, Rushmore, Spicer
-------------	--

### Similar sites


R107XA209IA	<b>Wet Upland Sedge Meadow</b> Wet Upland Sedge Meadows are similar in landscape position, but site is a MINERAL SOIL FLATS wetland
R107XA210IA	<b>Wet Upland Drainageway Prairie</b> Wet Upland Drainageway Prairies are similar in landscape position, but site is a SLOPE wetland

Table 1. Dominant plant species


Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Carex atherodes</i> (2) <i>Scolochloa festucacea</i>

### Physiographic features

Ponded Upland Depression Sedge Meadows occur on upland depressions (Figure 2). They are situated on elevations ranging from approximately 1000 to 1401 feet ASL. Ponding is occasional to frequent and lasts from two to less than 30 days. Ponded water depths can be as high as twelve inches.



**Figure 2. Figure 1. Location of Pondered Upland Depression Sedge Meadow ecological site within MLRA 107A.**



**Figure 3. Figure 2. Representative block diagram of Pondered Upland Depression Sedge Meadow and associated ecological sites.**

**Table 2. Representative physiographic features**

Slope shape across	(1) Concave
Slope shape up-down	(1) Concave
Landforms	(1) Upland > Depression
Runoff class	Negligible
Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	Occasional to frequent
Elevation	305–427 m
Slope	0–1%
Ponding depth	0–30 cm
Water table depth	0–183 cm

## Climatic features

The Iowa and Minnesota Loess Hills falls into the hot humid continental climate (Dfa) Köppen-Geiger climate classification (Peel et al. 2007). In winter, dry, cold air masses periodically shift south from Canada. As these air masses collide with humid air, snowfall and rainfall result. In summer, moist, warm air masses from the Gulf of Mexico migrate north, producing significant frontal or convective rains. Occasionally, hot, dry winds originating from the Desert Southwest will stagnate over the region, creating extended droughty periods in the summer from unusually high temperatures. Air masses from the Pacific Ocean can also spread into the region and dominate

producing mild, dry weather in the autumn known as Indian Summers (NCDC 2006).

The soil temperature regime of MLRA 107A is classified as mesic, where the mean annual soil temperature is between 46 and 59°F (USDA-NRCS 2006). Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south one travels. The average freeze-free period of this ecological site is about 159 days, while the frost-free period is about 144 days (Table 2). The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season. Average annual precipitation is about 32 inches, which includes rainfall plus the water equivalent from snowfall. The average annual low and high temperatures are 36 and 58°F, respectively (Table 3).

Climate data and analyses are derived from 30-year averages gathered from four National Oceanic and Atmospheric Administration (NOAA) weather stations contained within the range of this ecological site (Table 4).

**Table 3. Representative climatic features**

Frost-free period (characteristic range)	125-133 days
Freeze-free period (characteristic range)	148-149 days
Precipitation total (characteristic range)	762-813 mm
Frost-free period (actual range)	125-137 days
Freeze-free period (actual range)	147-149 days
Precipitation total (actual range)	737-838 mm
Frost-free period (average)	130 days
Freeze-free period (average)	148 days
Precipitation total (average)	787 mm


### **Climate stations used**

- (1) PRIMGHAR [USC00136800], Primghar, IA
- (2) ROCK RAPIDS [USC00137147], Rock Rapids, IA
- (3) CHEROKEE [USC00131442], Cherokee, IA
- (4) HOLSTEIN [USC00133909], Holstein, IA

### **Influencing water features**

Ponded Upland Depression Sedge Meadows are classified as a DEPRESSIONAL: Recharge, Ponded, Closed Depression, herbaceous wetland under the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008) and as a Palustrine, Persistent, Emergent Wetland under the National Wetlands Inventory (FGDC 2013). Precipitation, overland flow from adjacent uplands, and groundwater discharge are the main sources of water for this ecological site (Smith et al. 1995). Infiltration is very slow (Hydrologic Group D) for undrained soils, and surface runoff is negligible (Figure 5).

Primary wetland hydrology indicators for an intact Ponded Upland Depression Sedge Meadow may include: A1 Surface water, A2 High water table, A3 Saturation, and B7 Inundation visible on aerial photography. Secondary wetland hydrology indicators may include: C2 Dry-season water table, D2 Geomorphic position, and D5 FAC-neutral test (USACE 2010).




**Figure 8. Figure 5. Hydrologic cycling in Ponded Upland Depression Sedge Meadow ecological site.**

### Soil features

Soils of Ponded Upland Depression Sedge Meadows are in the Mollisols order, further classified as Typic Argialbolls that are impermeable with a negligible runoff potential. The soil series associated with this site includes Sperry (Figure 6). The parent material is loess, and the soils are very poorly-drained and deep with seasonal high-water tables. Soil pH classes are strongly acid to neutral (Table 5).

Soil map units in this ecological site, if not drained, may meet the definition of hydric soils and are listed as meeting criteria 2 and 3 of the hydric soils list (77 FR 12234).



**Figure 9. Figure 6. Profile sketches of soil series associated with Ponded Upland Depression Sedge Meadow.**

**Table 4. Representative soil features**

Parent material	(1) Loess
Family particle size	(1) Fine
Drainage class	Very poorly drained
Soil depth	203 cm

### Ecological dynamics

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

MLRA 107A is defined by a relatively low relief landscape that experiences lower rainfall amounts and available moisture compared to other MLRAs occurring to the south and east. As a result, prairie vegetation communities dominate the uplands, while forested communities are restricted to medium and large streams (Prior 1991; Eilers and Roosa 1994; MDNR 2017a, b). Ponded Upland Depression Sedge Meadows form an aspect of this vegetative continuum. This ecological site occurs on upland depressions on very-poorly drained soils. Species characteristic of this ecological site consist of hydrophytic herbaceous vegetation.


Ponding and fire are the most important ecosystem drivers for maintaining Ponded Upland Depression Sedge Meadows. The depth and duration of ponding affect species composition, cover, and vegetative production due to alternating aerobic and anaerobic surface substrate conditions. Periodic fires likely occurred on a two to five-year rotation interval and helped to reduce the accumulation of peat. The combination of fire and high-water levels prevented the establishment of shrubs for any significant amount of time (MDNR 2005; LANDFIRE 2009).

Drought has also played a role in shaping this ecological site. The periodic episodes of reduced soil moisture in conjunction with the very poorly-drained soils have favored the proliferation of plant species tolerant of such conditions. Drought can also slow the growth of plants and result in dieback of certain species. When coupled with fire, periods of drought can also eliminate or greatly reduce the occurrence of woody vegetation, substantially altering the extent of shrubs and trees (Pyne et al. 1996; MDNR 2005).

Today, Ponded Upland Depression Sedge Meadows have been virtually eliminated as the land has mostly been converted to agricultural production (Eilers and Roosa 1994; LANDFIRE 2009). Corn (*Zea mays* L.) and soybeans (*Glycine max* (L.) Merr.) are the dominant crops grown, but small patches of forage land may be present. A return to the historic plant community is likely not possible due to significant hydrologic and water quality changes in the watershed, but long-term conservation agriculture or habitat reconstruction efforts can help to restore some natural diversity and ecological functioning. The state-and-transition model that follows provides a detailed description of each state, community phase, pathway, and transition. This model is based on available experimental research, field observations, literature reviews, professional consensus, and interpretations.

## **State and transition model**

## R107AY208IA PONDED UPLAND DEPRESSION SEDGE MEADOW



Code	Process
T1A, T3A, T4A, T5A	Changes to natural hydroperiod and/or land abandonment
T1B, T2A, T4B, T5B	Cultural treatments are implemented to increase forage quality and yield
T1C, T2B, T3B, T5C	Agricultural conversion via tillage, seeding, and non-selective herbicide
1.1A	Reduction in average soil water levels
1.2A	Average soil water levels increase
2.1A	Increasing changes to hydrology and increasing sedimentation
R2A, R3A, R4A	Site preparation, non-native species control, and native seeding
3.1A	Mechanical harvesting is replaced with domestic livestock and continuous grazing
3.1B	Mechanical harvesting is replaced with domestic livestock and rest-rotational grazing
3.2A, 3.3B	Tillage, forage crop planting, and mechanical harvesting replace grazing
3.2B	Implementation of rest-rotational grazing
3.3A	Implementation of continuous grazing
4.1A	Less tillage, residue management
4.1B	Less tillage, residue management, and implementation of cover cropping
4.2B	Implementation of cover cropping
4.2A, 4.3B	Intensive tillage, remove residue, and reinitiate monoculture row cropping
4.3A	Remove cover cropping
5.1A	Invasive species control and implementation of disturbance regimes
5.2A	Drought or improper timing/use of management actions

### State 1 Reference State

The reference plant community is categorized as a sedge meadow, dominated by hydrophytic herbaceous vegetation. The two community phases within the reference state are dependent on ponding and fire. The depth and duration of ponding alters species composition, cover, and extent, while regular fire intervals keep woody

species from dominating. Drought has more localized impacts in the reference phases, but does contribute to overall species composition, diversity, cover, and productivity.

#### **Dominant plant species**

- wheat sedge (*Carex atherodes*), other herbaceous
- common rivergrass (*Scolochloa festucacea*), other herbaceous
- American mannagrass (*Glyceria grandis*), other herbaceous

### **Community 1.1**

#### **Wheat Sedge - Common Rivergrass**

Wheat Sedge – Common Rivergrass – Sites in this reference community phase are dominated by sedges and grasses that are interspersed with forbs. Mature plants typically range between 1.5 and 3 feet tall, and ground cover is continuous (75 to 100 percent). Wheat sedge is the dominant species of this reference community phase (sometimes forming monotypic stands), and common rivergrass is characteristic. Bluejoint, prairie cordgrass, river bulrush, and common spikerush can also occur. Common forbs include water knotweed, wild mint (*Mentha arvensis* L.), and white panicle aster (*Symphotrichum lanceolatum* (Willd.) G.L. Nesom ssp. lanceolatum var. lanceolatum) (MDNR 2005; LANDFIRE 2009; NatureServe 2015).

#### **Dominant plant species**

- wheat sedge (*Carex atherodes*), other herbaceous
- common rivergrass (*Scolochloa festucacea*), other herbaceous

### **Community 1.2**

#### **Wheat Sedge – American Mannagrass**

Wheat Sedge – American Mannagrass – This reference community phase can occur when ponding frequency and depth are reduced such as from periodic drought. Wheat sedge continues to form the dominant herbaceous component, but American mannagrass (*Glyceria grandis* S. Watson) replaces common rivergrass during this drier phase (Carey 1994; NatureServe 2015).

#### **Dominant plant species**

- wheat sedge (*Carex atherodes*), other herbaceous
- American mannagrass (*Glyceria grandis*), other herbaceous

### **Pathway 1.1A**

#### **Community 1.1 to 1.2**

Reduction in average soil water levels

### **Pathway 1.2A**

#### **Community 1.2 to 1.1**

Average soil water levels increase

## **State 2**

### **Hydrologically-altered State**

Hydrology is the most important determinant of wetlands and wetland processes. Hydrology modifies and determines the physiochemical environment (i.e., sediments, soil chemistry, water chemistry) which in turn directly affects the vegetation, animals, and microbes (Mitsch and Gosselink 2007). Human activities on landscape hydrology have greatly altered Ponded Upland Depression Sedge Meadows. Alterations such as agricultural tile draining and conversion to cropland on adjacent lands have changed the natural hydroperiod and rate of sedimentation as well as increased nutrient pollution (Werner and Zedler 2003; Mitsch and Gosselink 2007).

#### **Dominant plant species**



- reed canarygrass (*Phalaris arundinacea*), other herbaceous
- wheat sedge (*Carex atherodes*), other herbaceous

## Community 2.1

### Reed Canarygrass - Wheat Sedge

Reed Canarygrass – Wheat Sedge – This community phase represents the early changes to the natural wetland hydroperiod, sedimentation, and nutrient runoff. Sedimentation results in a reduction of soil organic matter and high dry bulk density. It also leads to a homogenization of the local microtopography, reducing the surface area and associated species diversity (Green and Galatowitsch 2002; Werner and Zedler 2002). Wheat sedge continues to form a component of the herbaceous layer, but the highly-invasive reed canarygrass (*Phalaris arundinacea* L.) co-dominates.

#### Dominant plant species

- reed canarygrass (*Phalaris arundinacea*), other herbaceous
- wheat sedge (*Carex atherodes*), other herbaceous

## Community 2.2

### Reed Canarygrass - Cattails

Reed Canarygrass – Cattails – Sites falling into this community phase have experienced significant sedimentation and are dominated by either a monoculture of reed canarygrass or a combination of reed canarygrass and cattails (*Typha* L.) (MDNR 2005). Reed canarygrass stands can significantly alter the physiochemical environment as well as the biotic communities, making the site only suitable to reed canarygrass. These monotypic stands create a positive feedback loop that perpetuates increasing sedimentation, altered hydrology, and dominance by this non-native species, especially in sites affected by nutrient enrichment from agricultural runoff (Vitousek 1995; Bernard and Lauve 1995; Green and Galatowitsch 2002; Werner and Zedler 2002; Kercher et al. 2007; Waggy 2010).

#### Dominant plant species

- reed canarygrass (*Phalaris arundinacea*), other herbaceous

## Pathway 2.1A

### Community 2.1 to 2.2

Increasing changes to hydrology and increasing sedimentation

## State 3

### Forage State

The forage state arises when the site is converted to a farming system that emphasizes domestic livestock production, known as grassland agriculture. Fire suppression, periodic cultural treatments (e.g., clipping, drainage, soil amendment applications, planting new species and/or cultivars, mechanical harvesting) and grazing by domesticated livestock transition and maintain this state (USDA-NRCS 2003). Early settlers seeded non-native species, as smooth brome (*Bromus inermis* Leyss.) and Kentucky bluegrass (*Poa pratensis* L.), to help extend the grazing season (Smith 1998). Over time, as lands were continuously harvested or grazed by herds of cattle, these species were able to spread and expand across the prairie ecosystem, reducing the native species diversity and ecological function.

## Community 3.1

### Hayfield

Hayfield – Sites in this community phase consist of forage plants that are planted and mechanically harvested. Mechanical harvesting removes much of the aboveground biomass and nutrients that feed the soil microorganisms (Franzluebbers et al. 2000; USDA-NRCS 2003). As a result, soil biology is reduced leading to decreases in nutrient uptake by plants, soil organic matter, and soil aggregation. Frequent biomass removal can also reduce the site's carbon sequestration capacity (Skinner 2008).

## **Community 3.2**

### **Continuous Pastured Grazing System**

Continuous Pastured Grazing System – This community phase is characterized by continuous grazing where domestic livestock graze a pasture for the entire season. Depending on stocking density, this can result in lower forage quality and productivity, weed invasions, and uneven pasture use. Continuous grazing can also increase the amount of bare ground and erosion and reduce soil organic matter, cation exchange capacity, water-holding capacity, and nutrient availability and retention (Bharati et al. 2002; Leake et al. 2004; Teague et al. 2011). Smooth brome, Kentucky bluegrass, and white clover (*Trifolium repens* L.) are common pasture species used in this phase. Their tolerance to continuous grazing has allowed these species to dominate, sometimes completely excluding the native vegetation.

## **Community 3.3**

### **Rest-Rotation Pastured Grazing System**

Rest-Rotation Pastured Grazing System – This community phase is characterized by rotational grazing where the pasture has been subdivided into several smaller paddocks. Through the development of a grazing plan, livestock utilize one or a few paddocks, while the remaining area is rested allowing plants to restore vigor and energy reserves, deepen root systems, develop seeds, as well as allow seedling establishment (Undersander et al. 2002; USDA-NRCS 2003). Rest-rotation pastured grazing systems include deferred rotation, rest rotation, high intensity – low frequency, and short duration methods. Vegetation is generally more diverse and can include orchardgrass (*Dactylis glomerata* L.), timothy (*Phleum pratense* L.), red clover (*Trifolium pratense* L.), and alfalfa (*Medicago sativa* L.). The addition of native prairie species can further bolster plant diversity and, in turn, soil function. This community phase promotes numerous ecosystem benefits including increasing biodiversity, preventing soil erosion, maintaining and enhancing soil quality, sequestering atmospheric carbon, and improving water yield and quality (USDA-NRCS 2003).

### **Pathway 3.1A**

#### **Community 3.1 to 3.2**

Mechanical harvesting is replaced with domestic livestock and continuous grazing

### **Pathway 3.1B**

#### **Community 3.1 to 3.3**

Mechanical harvesting is replaced with domestic livestock and rest-rotational grazing

### **Pathway 3.2A**

#### **Community 3.2 to 3.1**

Tillage, forage crop planting and mechanical harvesting replaces grazing

### **Pathway 3.2B**

#### **Community 3.2 to 3.3**

Implementation of rest-rotational grazing

### **Pathway 3.3B**

#### **Community 3.3 to 3.1**

Tillage, forage crop planting and mechanical harvesting replace grazing

### **Pathway 3.3A**

#### **Community 3.3 to 3.2**

Implementation of continuous grazing

## **State 4**

### **Cropland State**

The low topographic relief across the MLRA has resulted in nearly the entire area being converted to agriculture (Eilers and Roosa 1994). Agricultural tile drains used to lower the water table and the continuous use of tillage, row-crop planting, and chemicals (i.e., herbicides, fertilizers, etc.) have effectively eliminated the reference community and many of its natural ecological functions in favor of crop production. Corn and soybeans are the dominant crops for the site, and oats (*Avena L.*) and alfalfa (*Medicago sativa L.*) may be rotated periodically. These areas are likely to remain in crop production for the foreseeable future.

#### **Community 4.1**

##### **Conventional Tillage Field**

Conventional Tillage Field – Sites in this community phase typically consist of monoculture row-cropping maintained by conventional tillage practices. They are cropped in either continuous corn or corn-soybean rotations. The frequent use of deep tillage, low crop diversity, and bare soil conditions during the non-growing season negatively impacts soil health. Under these practices, soil aggregation is reduced or destroyed, soil organic matter is reduced, erosion and runoff are increased, and infiltration is decreased, which can ultimately lead to undesirable changes in the hydrology of the watershed (Tomer et al. 2005).

#### **Community 4.2**

##### **Conservation Tillage Field**

Conservation Tillage Field – This community phase is characterized by rotational crop production that utilizes various conservation tillage methods to promote soil health and reduce erosion. Conservation tillage methods include strip-till, ridge-till, vertical-till, or no-till planting systems. Strip-till keeps seedbed preparation to narrow bands less than one-third the width of the row where crop residue and soil consolidation are left undisturbed in-between seedbed areas. Strip-till planting may be completed in the fall and nutrient application either occurs simultaneously or at the time of planting. Ridge-till uses specialized equipment to create ridges in the seedbed and vegetative residue is left on the surface in between the ridges. Weeds are controlled with herbicides and/or cultivation, seedbed ridges are rebuilt during cultivation, and soils are left undisturbed from harvest to planting. Vertical-till systems employ machinery that lightly tills the soil and cuts up crop residue, mixing some of the residue into the top few inches of the soil while leaving a large portion on the surface. No-till management is the most conservative, disturbing soils only at the time of planting and fertilizer application. Compared to conventional tillage systems, conservation tillage methods can improve soil ecosystem function by reducing soil erosion, increasing organic matter and water availability, improving water quality, and reducing soil compaction.

#### **Community 4.3**

##### **Conservation Tillage with Cover Crop Field**

Conservation Tillage with Cover Crop Field – This community phase applies conservation tillage methods as described above as well as adds cover crop practices. Cover crops typically include nitrogen-fixing species (e.g., legumes), small grains (e.g., rye, wheat, oats), or forage covers (e.g., turnips, radishes, rapeseed). The addition of cover crops not only adds plant diversity but also promotes soil health by reducing soil erosion, limiting nitrogen leaching, suppressing weeds, increasing soil organic matter, and improving the overall soil ecosystem. In the case of small grain cover crops, surface cover and water infiltration are increased, while forage covers can be used to graze livestock or support local wildlife. Of the three community phases for this state, this phase promotes the greatest soil sustainability and improves ecological functioning within a cropland system.

#### **Pathway 4.1A**

##### **Community 4.1 to 4.2**

Less tillage, residue management

#### **Pathway 4.1B**

##### **Community 4.1 to 4.3**

Less tillage, residue management and implementation of cover cropping

### **Pathway 4.2A** **Community 4.2 to 4.1**

Intensive tillage, remove residue and reinitialize monoculture row cropping

### **Pathway 4.2B** **Community 4.2 to 4.3**

Implementation of cover cropping

### **Pathway 4.3B** **Community 4.3 to 4.1**

Intensive tillage, remove residue and reinitialize monoculture row cropping

### **Pathway 4.3A** **Community 4.3 to 4.2**

Remove cover cropping

## **State 5** **Reconstructed Sedge Meadow State**

Sedge meadow habitats provide multiple ecosystem services including flood abatement, water quality improvement, and biodiversity support. However, many sedge meadow communities have been stressed from watershed-scale changes in hydrology or eliminated as a result of type conversions to agricultural production, thereby significantly reducing these services (Zedler 2003). The extensive alterations of lands adjacent to Pondered Upland Depression Sedge Meadows may not allow for restoration back to the historic reference condition. But ecological reconstruction can aim to aid the recovery of degraded, damaged or destroyed functions. A successful reconstruction will have the ability to structurally and functionally sustain itself, demonstrate resilience to the natural ranges of stress and disturbance, and create and maintain positive biotic and abiotic interactions (SER 2002; Mitsch and Jørgensen 2004).

### **Community 5.1** **Early Successional Sedge Meadow**

Early Successional Sedge Meadow – This community phase represents the early community assembly from sedge meadow reconstruction and is highly dependent on seed viability, hydroperiod, soil organic matter content, and site preparation. Successful establishment of sedges can be maximized by using seed collected during the same growing season, utilizing genotypes adapted to the environmental location, ensuring soil is saturated at the time of seeding, and improving the water holding capacity and fertility of the soil (Budelsky and Galatowitsch 1999; van der Valk et al. 1999; Mitsch and Gosselink 2007; Hall and Zedler 2010). In addition, suppression and removal of non-native species is essential for reducing competition (Perry and Galatowitsch 2003).

### **Community 5.2** **Late Successional Sedge Meadow**

Late Successional Sedge Meadow – Appropriately timed disturbance regimes (e.g., hydroperiod, prescribed fire) and nutrient management applied to the early successional community phase can help increase the species richness, pushing the site into a late successional community phase over time (Mitsch and Gosselink 2007).

### **Pathway 5.1A** **Community 5.1 to 5.2**

Invasive species control and implementation of disturbance regimes

**Pathway 5.2A**  
**Community 5.2 to 5.1**

Drought or improper timing/use of management actions

**Transition T1A**  
**State 1 to 2**

Changes to natural hydroperiod and/or land abandonment

**Transition T1B**  
**State 1 to 3**

Cultural treatments are implemented to increase forage quality and yield

**Transition T1C**  
**State 1 to 4**

Agricultural conversion via tillage, seeding, and non-selective herbicide

**Transition T2A**  
**State 2 to 3**

Cultural treatments are implemented to increase forage quality and yield

**Transition T2B**  
**State 2 to 4**

Agricultural conversion via tillage, seeding and non-selective herbicide

**Transition R2A**  
**State 2 to 5**

Site preparation, non-native species control and native seeding

**Restoration pathway T3A**  
**State 3 to 2**

Changes to natural hydroperiod and/or land abandonment

**Transition T3B**  
**State 3 to 4**

Agricultural conversion via tillage, seeding and non-selective herbicide

**Transition R3A**  
**State 3 to 5**

Site preparation, non-native species control and native seeding

**Restoration pathway T4A**  
**State 4 to 2**

Changes to natural hydroperiod and/or land abandonment

## **Restoration pathway T4B**

### **State 4 to 3**

Cultural treatments are implemented to increase forage quality and yield

## **Transition R4A**

### **State 4 to 5**

Site preparation, non-native species control and native seeding

## **Restoration pathway T5A**

### **State 5 to 2**

Changes to natural hydroperiod and/or land abandonment

## **Restoration pathway T5B**

### **State 5 to 3**

Cultural treatments are implemented to increase forage quality and yield

## **Restoration pathway T5C**

### **State 5 to 4**

Agricultural conversion via tillage, seeding and non-selective herbicide

## **Additional community tables**

### **Inventory data references**

No field plots were available for this site. A review of the scientific literature and professional experience were used to approximate the plant communities for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on the sources identified in the ecological site description.

### **Other references**

Baker, R.G., C.A. Chumbley, P.M. Witinok, and H.K. Kim. 1990. Holocene vegetational changes in eastern Iowa. *Journal of the Iowa Academy of Science* 97: 167-177.

Baker, R.G., L.J. Maher, C.A. Chumbley, and K.L. Van Zant. 1992. Patterns of Holocene environmental changes in the midwestern United States. *Quaternary Research* 37: 379-389.

Bernard, J.B. and T.E. Lauve. 1995. A comparison of growth and nutrient uptake in *Phalaris arundinacea* L. growing in a wetland and a constructed bed receiving landfill leachate. *Wetlands* 15: 176-182.

Bharati, L., K.-H. Lee, T.M. Isenhardt, and R.C. Schultz. 2002. Soil-water infiltration under crops, pasture, and established riparian buffer in Midwestern USA. *Agroforestry Systems* 56: 249-257.

Budelsky, R.A. and S.M. Galatowitsch. 1999. Effects of moisture, temperature, and time on seed germination of five wetland Carices: implications for restoration. *Restoration Ecology* 7: 86-97.

Carey, J.H. 1994. *Scolochloa festucacea*. In: Fire Effects Information System [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available at <https://www.feis-crs.org/feis/>. (Accessed 23 January 2018).

Changes in Hydric Soils Database Selection Criteria. 77 Federal Register 12234 (29 February 2012), pp. 12234-12235.

- Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Coterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.
- Drobney, P.D., G.S. Wilhelm, D. Horton, M. Leoschke, D. Lewis, J. Pearson, D. Roosa, and D. Smith. 2001. Floristic Quality Assessment for the State of Iowa. Neal Smith National Wildlife Refuge and Ada Hayden Herbarium, Iowa State University, Ames, IA. 123 pps.
- Eggers, S.D. and D.M. Reed. 2015. Wetland Plants and Plant Communities of Minnesota and Wisconsin, Version 3.2. U.S. Army Corps of Engineers, Regulatory Branch, St. Paul District. St. Paul, MN. 478 pps.
- Eilers, L. and D. Roosa. 1994. The Vascular Plants of Iowa: An Annotated Checklist and Natural History. University of Iowa Press, Iowa City, IA. 319 pps.
- Federal Geographic Data Committee. 2013. Classification of Wetlands and Deepwater Habitats of the United States. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, D.C. 90 pps.
- Franzluebbers, A.J., J.A. Stuedemann, H.H. Schomberg, and S.R. Wilkinson. 2000. Soil organic C and N pools under long-term pasture management in the Southern Piedmont USA. *Soil Biology and Biochemistry* 32:469-478.
- Green, E.K. and S.M. Galatowitsch. 2002. Effects of *Phalaris arundinacea* and nitrate-N addition on the establishment of wetland plant communities. *Journal of Applied Ecology* 39: 134-144.
- Hall, S.J. and J.B. Zedler. 2010. Constraints on sedge meadow self-restoration in urban wetlands. *Restoration Ecology* 18: 671-680.
- Iowa Natural Areas Inventory [INAI]. 1984. An Inventory of Significant Natural Areas in Iowa: Two Year Progress Report of the Iowa Natural Areas Inventory. Iowa Natural Areas Inventory, Iowa Department of Natural Resources, Des Moines, IA.
- Kercher, S.M. A. Herr-Turnoff, J.B. Zedler. 2007. Understanding invasion as a process: the case of *Phalaris arundinacea* in wet prairies. *Biological Invasions* 9: 657-665.
- LANDFIRE. 2009. Biophysical Setting 4214880 Eastern Great Plains Wet Meadow-Marsh-Prairie System. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.
- Leake, J., D. Johnson, D. Donnelly, G. Muckle, L. Boddy, and D. Read. 2004. Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Canadian Journal of Botany* 82: 1016-1045.
- Minnesota Department of Natural Resources [MDNR]. 2005. Field Guide to the Native Plant Communities of Minnesota: The Prairie Parkland and Tallgrass Aspen Parklands Provinces. Ecological Land Classification Program, Minnesota County Biological Survey, and Natural Heritage and Nongame Research Program. Minnesota Department of Natural Resources, St. Paul, Minnesota.
- Minnesota Department of Natural Resources [MDNR]. 2017a. Coteau Moraines Subsection. Minnesota Department of Natural Resources: Ecological Classification System. Available at: <http://www.dnr.state.mn.us/ecs/251Bb/index.html>. (Accessed 10 October 2017).
- Minnesota Department of Natural Resources [MDNR]. 2017b. Inner Coteau Subsection. Minnesota Department of Natural Resources: Ecological Classification System. Available at: <http://www.dnr.state.mn.us/ecs/251Bc/index.html>. (Accessed 10 October 2017).
- Mitsch, W.J. and J.G. Gosselink. 2007. Wetlands, Fourth Edition. John Wiley & Sons, Inc. Hoboken, NJ. 582 pps.

- Mitsch, W.J. and S.E. Jørgensen. 2004. Ecological Engineering and Ecosystem Restoration. John Wiley & Sons, Inc. Hoboken, NJ. 428 pps.
- National Climate Data Center [NCDC]. 2006. Climate of Iowa. Central Region Headquarters, Climate Services Branch, National Climatic Data Center, Asheville, NC.
- NatureServe. 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 19 January 2018).
- Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.
- Perry, L.G. and S.M. Galatowitsch. 2003. A test of two annual cover crops for controlling *Phalaris arundinacea* invasion in restored sedge meadow wetlands. *Restoration Ecology* 11: 297-307.
- Prior, J.C. 1991. Landforms of Iowa. University of Iowa Press for the Iowa Department of Natural Resources, Iowa City, IA. 153 pps.
- Pyne, S.J., P.L. Andrews, and R.D. Laven. 1996. Introduction to Wildland Fire, Second Edition. John Wiley and Sons, Inc. New York, New York. 808 pps.
- Skinner, R.H. 2008. High biomass removal limits carbon sequestration potential of mature temperate pastures. *Journal for Environmental Quality* 37: 1319-1326.
- Smith, R.D., A. Ammann, C. Bartoldus, and M.M. Brinson. 1995. An Approach for Assessing Wetland Functions Using Hydrogeomorphic Classification, Reference Wetlands, and Functional Indices. U.S. Army Corps of Engineers, Waterways Experiment Station, Wetlands Research Program Technical Report WRP-DE-9. 78 pps.
- Smith, D.D. 1998. Iowa prairie: original extent and loss, preservation, and recovery attempts. *The Journal of the Iowa Academy of Sciences* 105: 94-108.
- Society for Ecological Restoration [SER] Science & Policy Working Group. 2002. The SER Primer on Ecological Restoration. Available at: <http://www.ser.org/>. (Accessed 28 February 2017).
- Teague, W.R., S.L. Dowhower, S.A. Baker, N. Haile, P.B. DeLaune, and D.M. Conover. 2011. Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems and Environment* 141: 310-322.
- Tomer, M.D., D.W. Meek, and L.A. Kramer. 2005. Agricultural practices influence flow regimes of headwater streams in western Iowa. *Journal of Environmental Quality* 34:1547-1558.
- Undersander, D., B. Albert, D. Cosgrove, D. Johnson, and P. Peterson. 2002. Pastures for Profit: A Guide to Rotational Grazing (A3529). University of Wisconsin-Extension and University of Minnesota Extension Service. 43 pps.
- U.S. Army Corps of Engineers [USACE]. 2010. Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Midwest Region (Version 2.0). U.S. Army Corps of Engineers, Wetlands Regulatory Assistance Program, U.S. Army Engineer Research and Development Center, Vicksburg, MS. 141 pps.
- United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2003. National Range and Pasture Handbook, Revision 1. Grazing Lands Technology Institute. 214 pps.
- United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. 682 pps.
- United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2007. Iowa NRCS Plant Community Species Lists. Des Moines, IA. Available at <https://www.nrcs.usda.gov/wps/>



portal/nrcs/detail/ia/technical/ecoscience/bio/?cid=nrcs142p2\_008160. (Accessed 19 January 2018).

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2008. Hydrogeomorphic Wetland Classification: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service. Technical Note No. 190-8-76. Washington, D.C. 8 pps.

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1: 3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-andiv-ecoregions-continental-united-states>. (Accessed 1 March 2017).

van der Valk, A.G., T.L. Bremholm, and E. Gordon. 1999. The restoration of sedge meadows: seed viability, seed germination requirements, and seedling growth of *Carex* species. *Wetlands* 19: 756-764.

Vitousek, P.M. 1995. Biological invasions and ecosystem processes: towards an integration of population biology and ecosystem studies. *Oikos* 57: 7-13.

Waggy, M.A. 2010. *Phalaris arundinacea*. In: Fire Effects Information System [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory. Available at: <https://www.feis-crs.org/feis/>. (Accessed 1 February 2017).

Werner, K.J. and J.B. Zedler. 2002. How sedge meadow soils, microtopography, and vegetation respond to sedimentation. *Wetlands* 3: 451-466.

Zedler, J.B. 2003. Wetlands at your service: reducing impacts of agriculture at the watershed scale. *Frontiers in Ecology and the Environment* 1: 65-72.

## Contributors

Lisa Kluesner  
Dan Pulido

## Approval

Chris Tecklenburg, 5/21/2020

## Acknowledgments

This project could not have been completed without the dedication and commitment from a variety of partners and staff (Table 6). Team members supported the project by serving on the technical team, assisting with the development of state and community phases of the state-and-transition model, providing peer review and technical editing, and conducting quality control and quality assurance reviews.

Table 6. List of primary contributors and reviewers.

Organization Name Title Location

Drake University:  
Dr. Tom Rosburg, Professor of Ecology and Botany, Des Moines, IA

Iowa Department of Natural Resources:  
John Pearson, Ecologist, Des Moines, IA

LANDFIRE (The Nature Conservancy):  
Randy Swaty, Ecologist, Evanston, IL

Natural Resources Conservation Service:  
Rick Bednarek, Iowa State Soil Scientist, Des Moines, IA

Patrick Chase, Area Resource Soil Scientist, Fort Dodge, IA  
 Stacey Clark, Regional Ecological Site Specialist, St. Paul, MN  
 James Cronin, State Biologist, Des Moines, IA  
 Tonie Endres, Senior Regional Soil Scientist, Indianapolis, IN  
 John Hammerly, Soil Data Quality Specialist, Indianapolis, IN  
 Lisa Kluesner, Ecological Site Specialist, Waverly, IA  
 Sean Kluesner, Earth Team Volunteer, Waverly, IA  
 Jeff Matthias, State Grassland Specialist, Des Moines, IA  
 Louis Moran, PhD, Area Resource Soil Scientist, Sioux City, IA  
 Kevin Norwood, Soil Survey Regional Director, Indianapolis, IN  
 Doug Oelmann, Soil Scientist, Des Moines, IA  
 James Phillips, GIS Specialist, Des Moines, IA  
 Dan Pulido, Soil Survey Leader, Atlantic, IA  
 Jason Steele, Area Resource Soil Scientist, Fairfield, IA  
 Doug Wallace, Ecological Site Specialist, Columbia, MO

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/05/2024
Approved by	Chris Tecklenburg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---