

# Ecological site R107XB007MO

## Loess Upland Prairie

Last updated: 5/21/2020  
Accessed: 02/08/2025

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 107X—Iowa and Missouri Deep Loess Hills

The Iowa and Missouri Deep Loess Hills (MLRA 107B) includes the Missouri Alluvial Plain, Loess Hills, Southern Iowa Drift Plain, and Central Dissected Till Plain landform regions (Prior 1991; Nigh and Schroeder 2002). It spans four states (Iowa, 53 percent; Missouri, 32 percent; Nebraska, 12 percent; and Kansas 3 percent), encompassing over 14,000 square miles (Figure 1). The elevation ranges from approximately 1,565 feet above sea level (ASL) on the highest ridges to about 600 feet ASL along the Missouri River near Glasgow in central Missouri. Local relief varies from 10 to 20 feet in the major river floodplains, to 50 to 100 feet in the dissected uplands, and loess bluffs of 200 to 300 feet along the Missouri River. Loess deposits cover most of the area, with deposits reaching a thickness of 65 to 200 feet in the Loess Hills and grading to about 20 feet in the eastern extent of the region. Pre-Illinoian till, deposited more than 500,000 years ago, lies beneath the loess and has experienced extensive erosion and dissection. Pennsylvanian and Cretaceous bedrock, comprised of shale, mudstones, and sandstones, lie beneath the glacial material (USDA-NRCS 2006).

The vegetation in the MLRA has undergone drastic changes over time. Spruce forests dominated the landscape 30,000 to 21,500 years ago. As the last glacial maximum peaked 21,500 to 16,000 years ago, they were replaced with open tundras and parklands. The end of the Pleistocene Epoch saw a warming climate that initially prompted the return of spruce forests, but as the warming continued, spruce trees were replaced by deciduous trees (Baker et al. 1990). Not until approximately 9,000 years ago did the vegetation transition to prairies as climatic conditions continued to warm and subsequently dry. Between 4,000 and 3,000 years ago, oak savannas began intermingling within the prairie landscape. This prairie-oak savanna ecosystem formed the dominant landscapes until the arrival of European settlers (Baker et al. 1992).

### Classification relationships

Major Land Resource Area (MLRA): Iowa and Missouri Deep Loess Hills (107B)

USFS Subregions: Central Dissected Till Plains Section (251C); Loess Hills (251Cb) Subsection (Cleland et al. 2007).

U.S. EPA Level IV Ecoregion: Steeply Rolling Loess Prairies (47e), Rolling Loess Prairies (47f), Nebraska/Kansas Loess Hills (47h)

Biophysical Setting (LANDFIRE 2009): Central Tallgrass Prairie (4214210)

Ecological Systems (National Vegetation Classification System, Nature Serve 2015): Central Tallgrass Prairie (CES205.683)

Eilers and Roosa (1994): Loess Hills

Iowa Department of Natural Resources (INAI nd): Southern Mesic Prairie

Missouri Natural Heritage Program (Nelson 2010): Mesic Loess/Glacial Till Prairie

Nebraska Game and Parks Commission (Steinauer and Rolfsmeier 2010): Upland Tall-Grass Prairie

Plant Associations (National Vegetation Classification System, Nature Serve 2015): *Andropogon gerardii* – *Sorghastrum nutans* – (*Sporobolus heterolepis*) – *Liatris* spp. – *Ratibida pinnata* Grassland (CEGL002203)

White (1983): Iowan Erosion Surface Upland Type

### Ecological site concept

Loess Upland Prairies are generally located within the green areas on the map (Figure 1). They occur on hillslopes, ridges, and interfluves. Soils are Inceptisols and Mollisols that are somewhat-poorly to well-drained and very deep, formed from loess with a slightly acidic (decreased pH) to slightly alkaline (increased pH) environment. These fine-silty, fertile soils have high soil uniformity resulting in increased nutrient- and water-holding capacity, increased organic matter retention, and good soil aeration that allows deep penetration by plant roots, which generally results in high plant productivity (Catt 2001).

The historic pre-European settlement vegetation on this site was dominated by a variety of tallgrass prairie species. Big bluestem (*Andropogon gerardii* Vitam) is the dominant monocot species, while leadplant (*Amorpha canescens* Pursh) and golden zizia (*Zizia aurea* (L.) W.D.J. Koch) are important indicators for the ecological site. Herbaceous species typical of an undisturbed plant community associated with this ecological site include Bicknell's sedge (*Carex bicknellii* Britton), prairie dropseed (*Sporobolus heterolepis* (A. Gray) A. Gray), stiff sunflower (*Helianthus pauciflorus* Nutt. ssp. *pauciflorus*), and stiff coreopsis (*Coreopsis palmata* Nutt.) (Drobney et al. 2001; Ladd and Thomas 2015). In addition to leadplant, Jersey tea (*Ceanothus herbaceus* Raf.) and prairie rose (*Rosa arkansana* Porter) are common shrubs that can be found scattered throughout the prairie. Fire was the primary disturbance factor that maintained this site, while drought and large mammal grazing were secondary factors (Kennedy 1970).

Relative to other prairie ecological sites in the MLRA, Loess Upland Prairies have high average clay content and overtop a somewhat restrictive paleosol which slows the rate of infiltration. As a result, Loess Upland Prairies are moister than associated prairie ecological sites as evidenced by the frequency of species with lower wetness coefficients (Drobney et al. 2001).

### Associated sites

R107XB027IA	<b>Calcareous Till Upland Prairie</b> Glacial till soils including Burchard, Liston, and Steinauer
R107XB008MO	<b>Loamy Footslope Savanna</b> Loess soils on footslopes, including Castana, Colo, Danbury, Deloit, Ely, Judson, Napier, Nodaway, Olmitz, Udarents, and Udorthents

### Similar sites

R107XB027IA	<b>Calcareous Till Upland Prairie</b> Calcareous Till Upland Prairies are similar in general species present but soils contains no carbonates, are derived from glacial till, and only occurs on shoulders and backslopes
R107XB012MO	<b>Calcareous Loess Upland Prairie</b> Calcareous Loess Upland Prairies are similar in landscape position but parent material is calcareous loess


R107XB006MO	<b>Calcareous Loess Exposed Backslope Prairie</b> Calcareous Loess Exposed Backslope Prairies are similar in landscape position but only occurs on south and west aspects, is drier as evidenced by a more xeric species composition, and parent material is calcareous loess
R107XB002MO	<b>Deep Loess Upland Prairie</b> Deep Loess Upland Prairies are similar in landscape position but species composition is less mesic

**Table 1. Dominant plant species**


Tree	Not specified
Shrub	(1) <i>Amorpha canescens</i>
Herbaceous	(1) <i>Andropogon gerardii</i> (2) <i>Zizia aurea</i>

## Physiographic features

Loess Upland Prairies occur on hillslopes, ridges, and interfluves on uplands on dissected till plains (Figure 2). This ecological site, situated on elevations ranging from approximately 600 to 1,560 feet ASL, is unique to the Loess Hills landform. This site does not experience flooding but rather generates runoff to adjacent, downslope ecological sites.



**Figure 2. Figure 1. Location of Loess Upland Prairie ecological site within MLRA 107B.**



**Figure 3. Figure 2. Representative block diagram of Loess Upland Prairie and associated ecological sites.**

**Table 2. Representative physiographic features**

Hillslope profile	(1) Summit (2) Shoulder (3) Backslope
Slope shape across	(1) Convex
Slope shape up-down	(1) Convex
Landforms	(1) Ridge (2) Interfluve (3) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	607–1,559 ft
Slope	4–22%
Water table depth	80 in
Aspect	Aspect is not a significant factor

### Climatic features

The Iowa and Missouri Deep Loess Hills falls into two Köppen-Geiger climate classifications (Peel et al. 2007): hot humid continental climate (Dfa) dominates the majority of the MLRA with small portions in the south falling into the humid subtropical climate (Cfa). In winter, dry, cold air masses periodically shift south from Canada. As these air masses collide with humid air, snowfall and rainfall result. In summer, moist, warm air masses from the Gulf of Mexico migrate north, producing significant frontal or convective rains (Decker 2017). Occasionally, high pressure will stagnate over the region, creating extended droughty periods. These periods of drought have historically occurred on 22-year cycles (Stockton and Meko 1983).

The soil temperature regime of MLRA 107B is classified as mesic, where the mean annual soil temperature is between 46 and 59°F (USDA-NRCS 2006). Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south you travel. The average freeze-free period of this ecological site is about 172 days, while the frost-free period is about 150 days (Table 2). The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season. Average annual precipitation is 35 inches, which includes rainfall plus the water equivalent from snowfall (Table 3). The average annual low and high temperatures are 38 and 60°F, respectively.

Climate data and analyses are derived from 30-year average gathered from seven National Oceanic and Atmospheric Administration (NOAA) weather stations contained within the range of this ecological site (Table 4).

**Table 3. Representative climatic features**

Frost-free period (characteristic range)	131-136 days
Freeze-free period (characteristic range)	152-166 days
Precipitation total (characteristic range)	34-36 in
Frost-free period (actual range)	125-137 days
Freeze-free period (actual range)	146-172 days
Precipitation total (actual range)	33-37 in
Frost-free period (average)	133 days
Freeze-free period (average)	161 days
Precipitation total (average)	35 in

### Climate stations used

- (1) CARROLL [USC00131233], Carroll, IA
- (2) DENISON [USC00132171], Denison, IA
- (3) ATLANTIC 1 NE [USC00130364], Atlantic, IA
- (4) AUDUBON [USC00130385], Audubon, IA
- (5) HARLAN 1N [USC00133632], Harlan, IA
- (6) SHENANDOAH [USC00137613], Shenandoah, IA
- (7) RED OAK [USC00136940], Red Oak, IA

## Influencing water features

Loess Upland Prairies are not influenced by wetland or riparian water features. Precipitation is the main source of water for this ecological site. Infiltration is moderate to very slow (Hydrologic Groups B, C and D), and surface runoff is medium to very high. Precipitation infiltrates the soil surface and percolates downward through the horizons. As it reaches the less permeable underlying paleosols, percolation is slowed but perching is not common (White 1983). The Dakota bedrock aquifer in the northern region of this ecological site is typically deep and confined, leaving it generally unaffected by recharge. However, there are surficial aquifers in the Pennsylvanian strata in the southern extent of the ecological site that are shallow and allow some recharge (Prior et al. 2003). Surface runoff contributes some water to downslope ecological sites. Evapotranspiration rates occur on a latitudinal gradient, with the northern end of the ecological site receiving a greater number of days with sun and high winds resulting in a higher average evapotranspiration rate compared to the southern end (Visher 1954).




Figure 10. Figure 5. Hydrologic cycling in Loess Upland Prairie ecological site.

## Soil features

Soils of Loess Upland Prairies are in the Inceptisol and Mollisol orders, further classified as Dystric Eutrudepts, Aquertic Argiudolls, Aquic Pachic Hapludolls, Typic Argiudolls, or Typic Hapludolls. They were formed under prairie vegetation. The soil series associated with this site includes Arisburg, Arispe, Arthur, Exira, Marshall, Minden, and Polo. The parent material is loess, and the soils are somewhat-poorly to well-drained and very deep with no coarse fragments. Soil pH classes are very strongly acid to moderately alkaline. No rooting restrictions are noted for the soils of this ecological site. Average clay content is between 27 and 35 percent limiting extreme compaction, but erosion from wind and water can be high.



Figure 11. Polo series




Figure 12. Figure 6. Profile sketches of soil series associated with Loess Upland Prairie.

Table 4. Representative soil features

Parent material	(1) Loess
Surface texture	(1) Silt loam
Family particle size	(1) Fine-silty
Drainage class	Somewhat poorly drained to well drained
Permeability class	Very slow
Soil depth	80 in
Available water capacity (0-40in)	7-8 in
Calcium carbonate equivalent (0-40in)	0-30%
Electrical conductivity (0-40in)	0-2 mmhos/cm
Sodium adsorption ratio (0-40in)	0
Soil reaction (1:1 water) (0-40in)	4.5-8.4

## Ecological dynamics

Prairie ecosystems are regarded as the most endangered ecosystem in North America where an estimated four percent of the tallgrass prairie habitat remains (Steinauer and Collins 1996). The Loess Hills region of MLRA 107B

were once dominated by tall and mixed-grass prairies, extending across more than 90 percent of the area (Rosburg 1994; Farnsworth 2009). However, by the early twenty-first century much of the land had been converted to agriculture, leaving an estimated 20 percent of the region to be classified as “grassland” and another three percent classified as “remnant prairie” (Farnsworth 2009).

Loess Upland Prairies form a vegetative continuum throughout the Loess Hills, where soil moisture serves as the primary influence on community composition (White 1983; White and Glenn-Lewin 1984). This ecological site can occur on nearly any aspect. Species characteristic of this ecological site are sun-loving, fire- and drought-adapted plants.

Fire is the most important ecosystem driver for maintaining this ecological site (Vogl 1974; Anderson 1990; Eilers and Roosa 1994). Fire intensity typically consisted of periodic, low-intensity surface fires (Stambaugh et al. 2006; LANDFIRE 2009). Ignition sources included summertime lightning strikes from convective storms and bimodal, human ignitions during the spring and fall seasons. Native Americans regularly set fires to improve sight lines for hunting, driving large game, improving grazing and browsing habitat, agricultural and village clearing, and enhancing vital ethnobotanical plants (Day 1953; Barrett 1980; White 1994). Fire frequency has been estimated to occur on average every 6.6 years in the Loess Hills region (Stambaugh et al. 2006). This continuous disturbance provided critical conditions for perpetuating the native prairie ecosystem.

Grazing by native ungulates is often cited as an important disturbance regime of North American grasslands, with bison (*Bison bison*), prairie elk (*Cervus elaphus*), and white-tailed deer (*Odocoileus virginianus*) serving as the dominant herbivores of the area. However, plant community succession in the Loess Hills region does not necessarily follow this hypothesis. The steep and rugged topography of the Loess Hills has been considered an impediment to grazing by large ungulates such as bison. Any role bison played in the area was most likely relegated to the northwestern extent where the terrain is milder (Dinsmore 1994). Elk and deer are believed to have played a relatively significant role in keeping woody vegetation at bay in the prairies of the Loess Hills (Farnsworth 2009; LANDFIRE 2009).

Drought has also played a role in shaping the prairie ecosystems in the Loess Hills. The periodic episodes of reduced soil moisture in conjunction with the well-drained soils have favored the proliferation of plant species tolerant of such conditions (Stambaugh et al. 2006). In addition, drought can also slow the growth of plants and result in dieback of certain species. When coupled with fire, periods of drought can also greatly delay the recovery of woody vegetation, substantially altering the extent of shrubs and trees (Pyne et al. 1996).

Today, Loess Upland Prairies are limited in their extent, having been converted to pasture or agricultural land. What remnants do exist show evidence of indirect anthropogenic influence as some non-native species (e.g., Kentucky bluegrass (*Poa pratensis* L.), smooth brome (*Bromus inermis* Leyss.)) are present in the community composition. A return to the historic plant community is likely not possible, but long-term restoration efforts can help to restore some natural diversity and ecological functioning.

## **State and transition model**

## R107BY007MO LOESS UPLAND PRAIRIE




Figure 13. STM

### State 1 Reference State

The reference plant community is categorized as a mesic tallgrass prairie and includes grasses, forbs, and varying components of shrubs. The reference community phase is dependent on a fire frequency of every one to six years. Shorter fire intervals maintain dominance by grasses, while less frequent intervals allow woody vegetation to increase their importance in the plant canopy. Grazing and drought disturbances have less impact in the reference phases, but do contribute to overall species composition, diversity, cover, and productivity.

#### Dominant plant species

- leadplant (*Amorpha canescens*), shrub
- big bluestem (*Andropogon gerardii*), grass

### Community 1.1 Leadplant/Big Bluestem – Golden Zizia

Many species are present on the site, but leadplant, big bluestem, and golden zizia are the diagnostic species. Characteristic forbs include white heath aster (*Symphotrichum ericoides* (L.) G.L. Nesom var. *ericoides*), flowering spurge (*Euphorbia corollata* L.), and smooth oxeye (*Heliopsis helianthoides* (L.) Sweet). Jersey tea, American plum (*Prunus americana* Marshall), black cherry (*Prunus serotina* Ehrh.), and prairie rose are common associates of the shrub component in this community.



## **Dominant plant species**

- leadplant (*Amorpha canescens*), shrub
- little bluestem (*Schizachyrium scoparium*), grass
- golden zizia (*Zizia aurea*), other herbaceous

## **State 2**

### **Cropland State**

Loess is the main contributing factor to the Midwest's highly-productive agricultural soils, and as a result, much of the MLRA has been converted to cropland, including significant portions of this ecological site (USGS 1999). The continuous use of tillage, row-crop planting, and chemicals (i.e., herbicides, fertilizers, etc.) have effectively eliminated the reference community and many of its natural ecological functions in favor of crop production. Corn (*Zea mays* L.) and soybeans (*Glycine max* (L.) Merr.) are the dominant crops for the site. These areas are likely to remain in crop production for the foreseeable future.

### **Community 2.1**

#### **Conventional Tillage Field**

Sites in this community phase typically consist of monoculture row-cropping maintained by conventional tillage practices. They are cropped in either continuous corn or corn-soybean rotations. The frequent use of deep tillage, low crop diversity, and bare soil conditions during the non-growing season negatively impact soil health. Under these practices, soil aggregation is reduced or destroyed, soil organic matter is reduced, erosion and runoff are increased, and infiltration is decreased, which can ultimately lead to undesirable changes in the hydrology of the watershed (Tomer et al. 2005).

### **Community 2.2**

#### **Conservation Tillage Field**

This community phase is characterized by rotational crop production that utilizes various conservation tillage methods to promote soil health and reduce erosion. Conservation tillage methods include strip-till, ridge-till, vertical-till, or no-till planting systems. Strip-till keeps seedbed preparation to narrow bands less than one-third the width of the row where crop residue and soil consolidation are left undisturbed in-between seedbed areas. Strip-till planting may be completed in the fall and nutrient application either occurs simultaneously or at the time of planting. Ridge-till uses specialized equipment to create ridges in the seedbed and vegetative residue is left on the surface in between the ridges. Weeds are controlled with herbicides and/or cultivation, seedbed ridges are rebuilt during cultivation, and soils are left undisturbed from harvest to planting. Vertical-till systems employ machinery that lightly tills the soil and cuts up crop residue, mixing some of the residue into the top few inches of the soil while leaving a large portion on the surface. No-till management is the most conservative, disturbing soils only at the time of planting and fertilizer application. Compared to conventional tillage system, conservation tillage methods can reduce soil erosion, increase organic matter and water availability, improve water quality, and reduce soil compaction.

### **Community 2.3**

#### **Conservation Tillage Field/Alternative Crop Field**

This condition applies conservation tillage methods as described above as well as adds cover crop practices. Cover crops typically include nitrogen-fixing species (e.g., legumes), small grains (e.g., rye, wheat, oats), or forage covers (e.g., turnips, radishes, rapeseed). The addition of cover crops not only adds plant diversity but also promotes soil health by reducing soil erosion, limiting nitrogen leaching, suppressing weeds, increasing soil organic matter, and improving the overall soil. In the case of small grain cover crops, surface cover and water infiltration are increased, while forage covers can be used to graze livestock or support local wildlife. Of the three community phases for this state, this phase promotes the greatest soil sustainability and improves ecological functioning within a cropland system.

## **Pathway P2.1A**

### **Community 2.1 to 2.2**

Tillage operations are greatly reduced, crop rotation occurs on a regular schedule, and crop residue is allowed to remain on the soil surface.

### **Pathway P2.1B** **Community 2.1 to 2.3**

Cover crops are implemented to prevent soil erosion

### **Pathway P2.2A** **Community 2.2 to 2.1**

Intensive tillage is utilized and monoculture row-cropping is established.

### **Pathway P2.2B** **Community 2.2 to 2.3**

Cover crops are implemented to prevent soil erosion.

### **Pathway P2.3B** **Community 2.3 to 2.1**

Intensive tillage is utilized, cover crops practices are abandoned, monoculture row-cropping is established, and crop rotation is reduced or eliminated.

### **Pathway P2.3A** **Community 2.3 to 2.2**

Cover crop practices are abandoned.

## **State 3** **Cool Season Grasses**

The cool-season pasture state occurs when the reference state has been anthropogenically-altered for livestock production. Fire suppression, seeding of non-native cool-season grasses, removal of woody vegetation, periodic cultural treatments, and grazing by domesticated livestock transition and maintain this simplified grassland state (Rosburg 1994; USDA-NRCS 2003). Early settlers seeded such non-native cool-season species as smooth brome (*Bromus inermis* Leyss.) and Kentucky bluegrass in order to help extend the grazing season (Smith 1998). Over time, as lands were continually grazed by large herds of cattle, the non-native species were able to spread and expand across the prairie habitat, reducing the native species diversity.

#### **Dominant plant species**

- smooth brome (*Bromus inermis*), grass
- Kentucky bluegrass (*Poa pratensis*), grass

### **Community 3.1** **Smooth Brome – Kentucky Bluegrass**

Species characteristic of this community phase include big bluestem, smooth brome, and Kentucky bluegrass. While the native big bluestem forms the dominant component of the canopy, smooth brome and Kentucky bluegrass occur in higher frequencies across the site. Annuals and biennials are important components of this community phase and are indicative of the disturbed nature of the site (Rosburg 1994).

#### **Dominant plant species**

- smooth brome (*Bromus inermis*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- big bluestem (*Andropogon gerardii*), grass

## State 4

### Oldfield State

Agricultural lands that have been abandoned and left fallow undergo ecological succession to the oldfield state. Two community phases define this state and are determined by time and human activities. Initially, the site is inhabited by ruderal, herbaceous annuals and perennials. As the land remains untouched from natural or anthropogenic disturbances, woody species begin to dominate eventually forming a near continuous canopy.

#### Dominant plant species

- eastern redcedar (*Juniperus virginiana*), tree
- roughleaf dogwood (*Cornus drummondii*), shrub
- lambsquarters (*Chenopodium album*), other herbaceous
- red clover (*Trifolium pratense*), other herbaceous
- goldenrod (*Solidago*), other herbaceous

### Community 4.1

#### Weedy Herbaceous Annuals - Perennials

This community phase represents the early stages of land abandonment characterized by bare soil and lack of human land modification. During these first few years, the phase is dominated by herbaceous annual (e.g. lambsquarters (*Chenopodium album* L.)) and perennial (e.g., red clover (*Trifolium pratense* L.) and goldenrod (*Solidago* L.)) species. Annuals are the dominant life form comprising nearly 85 percent of the species richness but within a few years perennial species increase their importance on the site. Non-natives and natives are relatively equally dominant throughout this phase, with natives contributing around 40 percent of the species richness initially and increasing to 60 percent (Brown et al. 1987).

### Community 4.2

#### Slippery Elm-Eastern Redcedar/Roughleaf Dogwood

Sites falling into this community phase are strongly dominated by a limited number of early-successional woody species. Plant community composition is largely shaped by abiotic effects of patch size, dispersal dynamics, and distance effects. Generally, roughleaf dogwood (*Cornus drummondii* C.A. Mey.) forms the most frequent or abundant woody species across the site due to its clonal stem production and high seed yield. Slippery elm (*Ulmus rubra* Muhl.) is a frequent invader of abandoned agricultural or pasture lands with a roughly linear colonization rate over the years. Eastern redcedar (*Juniperus virginiana* L.) is a pioneer species and can be a co-dominant component of the tree canopy, with densities typically higher in smaller fields (Yao et al. 1999). Herbaceous species that can be found in the understory include common blue violet (*Viola sororia* Willd.), white snakeroot (*Ageratina altissima* (L.) R.M. King & H. Rob. Var. *altissima*), and creeper (*Parthenocissus* Planch.) (Rosburg 1994).

#### Dominant plant species

- slippery elm (*Ulmus rubra*), tree
- eastern redbud (*Cercis canadensis*), tree
- roughleaf dogwood (*Cornus drummondii*), shrub

### Pathway P4.1A

#### Community 4.1 to 4.2

Soil is left undisturbed for more than six years.

### Pathway P4.2A

#### Community 4.2 to 4.1

Fire or other disturbance factor is used to control and eliminate the woody plant cover.

## State 5

## **Reconstructed Prairie State**

Prairie reconstructions have become an important tool for repairing natural ecological functioning and providing habitat protection for numerous grassland-dependent species. The historic plant community of the tallgrass prairie was extremely diverse and complex, and prairie replication is not considered to be possible once the native vegetation has been altered by post-European settlement land uses. Therefore ecological restoration should aim to aid the recovery of degraded, damaged, or destroyed ecosystems. A successful restoration will have the ability to structurally and functionally sustain itself, demonstrate resilience to the natural ranges of stress and disturbance, and create and maintain positive biotic and abiotic interactions (SER 2002). The reconstructed prairie state is the result of a long-term commitment involving a multi-step, adaptive management process. Diverse, species-rich seed mixes are important to utilize as they allow the site to undergo successional stages that exhibit changing composition and dominance over time (Smith et al. 2010). On-going management via prescribed fire and/or light grazing will help the site progress from an early successional community dominated by annuals and some weeds to a later seral stage composed of native perennial grasses, forbs, and shrubs. Establishing a prescribed fire regimen that mimics natural disturbance patterns can increase native species cover and diversity while reducing cover of non-native forbs and grasses. Light grazing alone can help promote species richness, while grazing accompanied with fire can control the encroachment of woody vegetation (Brudvig et al. 2007).

### **Community 5.1**

#### **Early Successional Reconstructed Prairie**

This community phase represents the early community assembly from prairie reconstruction and is highly dependent on the seed mix utilized and the timing and priority of planting operations. The seed mix should look to include a diverse mix of native cool-season and warm-season annual and perennial grasses and forbs typical of the reference state. Cool-season annuals can help to provide litter that promotes cool, moist soil conditions to the benefit of the other species in the seed mix. The first season following site preparation and seeding will typically result in annuals and other volunteer species forming the vegetative cover. Control of non-native species, particularly perennial species, is crucial at this point in order to ensure they do not establish before the native vegetation (Martin and Wilsey 2012). After the first season, native C4 grasses should begin to become more prominent on the landscape and over time close the canopy.

### **Community 5.2**

#### **Late Successional Reconstructed Prairie**

Appropriately timed disturbance regimes (e.g., prescribed fire) applied to the early successional community phase can help increase the beta diversity, pushing the site into a late successional community phase over time. While prairie communities are dominated by grasses, these species can suppress forb establishment and reduce overall diversity and ecological functioning (Martin and Wilsey 2006; Williams et al. 2007). Reducing accumulated plant litter from perennial bunchgrasses allows more light and nutrients to become available for forb recruitment, allowing for greater ecosystem complexity (Wilsey 2008).

### **Pathway P5.1A**

#### **Community 5.1 to 5.2**

Application of stand improvement practices in line with a developed management plan.

### **Pathway P5.2A**

#### **Community 5.2 to 5.1**

Reconstruction experiences a setback from extreme weather event or improper timing of management actions.

### **Transition T1A**

#### **State 1 to 2**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (2).

### **Transition T1B**

### **State 1 to 3**

Interseeding non-native cool-season grasses and brush control transition this site to the cool-season pasture state (3).

### **Transition T2A**

#### **State 2 to 3**

Non-selective herbicide and seeding of non-native cool-season grasses transitions the site to the cool-season pasture state (3).

### **Transition T2B**

#### **State 2 to 4**

Land is abandoned and left fallow; natural succession by opportunistic species transitions this site to the oldfield state (4).

### **Transition R2A**

#### **State 2 to 5**

Land is entered into a CRP or WRP contract or other restoration initiative and site preparation, invasive species control (native and non-native), and seeding native species transition this site to the reconstructed prairie state (5).

### **Transition T3A**

#### **State 3 to 2**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (2).

### **Transition T3B**

#### **State 3 to 4**

Land is abandoned and left fallow; natural succession by opportunistic species transitions this site to the oldfield state (4).

### **Restoration pathway R3A**

#### **State 3 to 5**

Site preparation, invasive species control (native and non-native), and seeding native species transition this site to the reconstructed prairie state (5).

### **Transition T4B**

#### **State 4 to 2**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (2).

### **Transition T4A**

#### **State 4 to 3**

Brush control and interseeding of non-native, cool-season grasses transition this site to the cool-season pasture state (3).

### **Restoration pathway R4A**

#### **State 4 to 5**

Restoration 4A – Site preparation, invasive species control (native and non-native), and seeding native species transition this site to the reconstructed prairie state (5).

## **Transition T5A**

### **State 5 to 2**

Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (2).

## **Transition T5B**

### **State 5 to 3**

Land is converted to the cool-season pasture state through the use of non-selective herbicide and seeding of non-native cool-season grasses (3).

## **Transition T5C**

### **State 5 to 4**

Transition 5C – Active management of the restored prairie is ceased and woody encroachment transitions this site to the oldfield state (4).

## **Additional community tables**

### **Animal community**

#### Wildlife\*

Game species that utilize this ecological site include:

Northern Bobwhite will utilize this ecological site for food (seeds, insects) and cover needs (escape, nesting and roosting cover).

Cottontail rabbits will utilize this ecological site for food (seeds, soft mast) and cover needs.

Turkey will utilize this ecological site for food (seeds, green browse, soft mast, and insects) and nesting and brood-rearing cover. Turkey poults feed heavily on insects provided by this site type.

White-tailed Deer will utilize this ecological site for browse (plant leaves in the growing season, seeds and soft mast in the fall/winter). This site type also can provide escape cover.

Bird species associated with this ecological site's reference state condition:

Breeding birds as related to vegetation structure (related to time since fire, grazing, haying, and mowing):

Vegetation Height Short (< 1.5 feet, low litter levels, bare ground visible):

Grasshopper Sparrow, Horned Lark, Upland Sandpiper, Greater Prairie Chicken, Northern Bobwhite

Mid-Vegetation Height (1.5 – 3 feet, moderate litter levels, some bare ground visible):

Eastern Meadowlark, Dickcissel, Field Sparrow, Upland Sandpiper, Greater Prairie Chicken, Northern Bobwhite, Eastern Kingbird, Bobolink, Lark Sparrow

Tall Vegetation Height (> 3 feet, moderate-high litter levels, little bare ground visible):

Henslow's Sparrow, Dickcissel, Greater Prairie Chicken, Field Sparrow, Northern Bobwhite, Sedge Wren, Northern Harrier

Brushy – mix of grasses, forbs, native shrubs (e.g., *Rhus copallina*, *Prunus americana*, *Rubus* spp., *Rosa carolina*) and small trees (e.g., *Cornus racemosa*): Bell's Vireo, Yellow-Breasted Chat, Loggerhead Shrike, Brown Thrasher, Common Yellowthroat

Winter Resident: Short-Eared Owl, Le Conte's Sparrow

Amphibian and reptile species associated with this ecological site's reference state condition: prairies with or nearby to fishless ponds/pools (may be ephemeral) may have Eastern Tiger Salamander (*Ambystoma tigrinum tigrinum*) and Western Chorus Frog (*Pseudacris triseriata triseriata*); prairies with crawfish burrows may have Northern

Crawfish Frog (*Rana areolata circulosa*); other species include Northern Prairie Skink (*Eumeces septentrionalis septentrionalis*), Ornate Box Turtle (*Terrapene ornata ornata*), Western Slender Glass Lizard (*Ophisaurus attenuatus attenuatus*), Eastern Yellow-bellied Racer (*Coluber constrictor flaviventris*), Prairie Ring-necked Snake (*Diadophis punctatus arnyi*), and Bullsnake (*Pituophis catenifer sayi*).

Small mammals associated with this ecological site's reference state condition: Least Shrew (*Cryptotis parva*), Franklin's Ground Squirrel (*Spermophilus franklinii*), Plains Pocket Gopher (*Geomys bursarius*), Prairie Vole (*Microtus ochrogaster*), Southern Bog Lemming (*Synaptomys cooperi*), Meadow Jumping Mouse (*Zapus hudsonius*), Thirteen-lined Ground Squirrel (*Spermophilus tridecemlineatus*) and Badger (*Taxidea taxus*).

Invertebrates: Many native insect species are likely associated with this ecological site, especially native bees, ants, beetles, butterflies and moths, and crickets, grasshoppers and katydids. However information on these groups is often lacking enough resolution to assign them to individual ecological sites.

Insect species known to be associated with this ecological site's reference state condition: Regal Fritillary butterfly (*Speyeria idalia*) whose larvae feed primarily on native prairie violets (*Viola pedata*, *V. pedatifida*, and *V. sagittata*); Mottled Dusky Wing butterfly (*Erynnis martialis*), Golden Byssus butterfly (*Problema byssus kumskaka*), Delaware Skipper butterfly (*Atryone logan logan*), and Crossline Skipper butterfly (*Polites origenes*). The larvae of the moth *Eucosma bipunctella* bore into compass plant (*Silphium laciniatum*) roots and feed and the larvae of the moth *Eucosma giganteana* bore into a number of *Silphium* species roots and feed. Native bees, important pollinators, that may be associated with this ecological site's reference condition include: *Colletes brevicornis*, *Andrena beameri*, *A. helianthiformis*, *Protandrena rudbeckiae*, *Halictus parallelus*, *Lasioglossum albipennis*, *L. coreopsis*, *L. disparilis*, *L. nymphaeum*, *Ashmeadiella buconis*, *Megachile addenda*, *Anthidium psoraleae*, *Eucera hamata*, *Melissodes coloradensis*, *M. coreopsis*, and *M. vernoniae*. The Short-winged Katydid (*Amblycorypha parvipennis*), Green Grasshopper (*Hesperotettix speciosus*) and Two-voiced Conehead katydid (*Neoconcephalus bivocatus*) are possible orthopteran associates of this ecological site.

Other invertebrate associates include the Grassland Crayfish (*Procambarus gracilis*).

\*This section prepared by Mike Leahy, Natural Areas Coordinator, Missouri Department of Conservation, 2013

## Other information

### Forestry

Management: This ecological site is not recommended for traditional timber management activity. Historically this site was dominated by a ground cover of native prairie grasses and forbs. Some scattered open grown trees may have also been present. May be suitable for non-traditional forestry uses such as windbreaks, environmental plantings, alley cropping (a method of planting, in which rows of trees or shrubs are interspersed with rows of crops) or woody bio-fuels.

## Inventory data references

Tier 3 Sampling Plot(s) used to develop the reference state, community phase 1.1:

State County Ownership Legal Description Easting Northing  
Iowa Crawford King Cemetery – Milford Township T80N R38W S28 315410 4663401  
Iowa Shelby Dinesen Prairie State Preserve T84N R38W S11 310253 4619895

## Other references

Anderson, R.C. 1990. The historic role of fire in the North American grassland. In: S.L. Collins and L.L. Wallace, eds. *Fire in North American Tallgrass Prairie*. University of Oklahoma Press, Norman, OK.

Baker, R.G., C.A. Chumbley, P.M. Witinok, and H.K. Kim. 1990. Holocene vegetational changes in eastern Iowa. *Journal of the Iowa Academy of Science* 97: 167-177.

Baker, R.G., L.J. Maher, C.A. Chumbley, and K.L. Van Zant. 1992. Patterns of Holocene environmental changes in

the midwestern United States. *Quaternary Research* 37: 379-389.

Barrett, S.W. 1980. Indians and fire. *Western Wildlands* Spring: 17-20.

Brown, V.K., S.D. Hendrix, and H. Dingle. 1987. Plants and insects in early old-field succession: comparison on an English site and an American site. *Biological Journal of the Linnean Society* 31: 59-74.

Brudvig, L.A., C.M. Mabry, J.R. Miller, and T.A. Walker. 2007. Evaluation of central North American prairie management based on species diversity, life form, and individual species metrics. *Conservation Biology* 21: 864-874.

Catt, J. 2001. The agricultural importance of loess. *Earth-Science Reviews* 54: 213-229.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. *Ecological Subregions: Sections and Subsections of the Conterminous United States*. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Day, G. 1953. The Indian as an ecological factor in the northeastern forest. *Ecology* 34: 329-346.

Decker, W.L. 2017. *Climate of Missouri*. University of Missouri, Missouri Climate Center, College of Agriculture, Food and Natural Resources. Available at <http://climate.missouri.edu/climate.php>. (Accessed 24 February 2017).

Dinsmore, J.J. 1994. *A Country So Full of Game: The Story of Wildlife in Iowa*. University of Iowa Press, Iowa City, Iowa. 261 pps.

Drobney, P.D., G.S. Wilhelm, D. Horton, M. Leoschke, D. Lewis, J. Pearson, D. Roosa, and D. Smith. 2001. *Floristic Quality Assessment for the State of Iowa*. Neal Smith National Wildlife Refuge and Ada Hayden Herbarium, Iowa State University, Ames, IA.

Eilers, L. and D. Roosa. 1994. *The Vascular Plants of Iowa: An Annotated Checklist and Natural History*. University of Iowa Press, Iowa City, IA. 319 pps.

Farnsworth, D.A. 2009. *Establishing restoration baselines for the Loess Hills region*. M.S. Thesis. Iowa State University, Ames, IA. 123 pps.

Iowa Natural Areas Inventory [INAI]. No date. *Vegetation Classification of Iowa*. Iowa Natural Areas Inventory, Iowa Department of Natural Resources, Des Moines, IA.

Kennedy, R.F. 1970. *Sheeder Prairie Preserve: A Natural Prairie Landscape in Guthrie County, Iowa*. Report to the Iowa State Conservation Commission and the Iowa State Preserves Advisory Board. Department of Botany and Plant Pathology, Iowa State University, Ames, Iowa. 37 pps.

Ladd, D. and J.R. Thomas. 2015. Ecological checklist of the Missouri flora for Floristic Quality Assessment. *Phytoneuron* 12: 1-274.

LANDFIRE. 2009. *Biophysical Setting 4214210 Central Tallgrass Prairie*. In: *LANDFIRE National Vegetation Dynamics Models*. USDA Forest Service and US Department of Interior. Washington, DC.

Martin, L.M. and B.J. Wilsey. 2006. Assessing grassland restoration success: relative roles of seed additions and native ungulate activities. *Journal of Applied Ecology* 43: 1098-1110.

Martin, L.M. and B.J. Wilsey. 2012. Assembly history alters alpha and beta diversity, exotic-native proportions and functioning of restored prairie plant communities. *Journal of Applied Ecology* 49: 1436-1445.

NatureServe. 2015. *NatureServe Explorer: An online encyclopedia of life* [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 13 February 2017).

Nelson, P. 2010. *The Terrestrial Natural Communities of Missouri, Revised Edition*. Missouri Natural Areas



Committee, Department of Natural Resources and the Department of Conservation, Jefferson City, MO. 500 pps.

Nigh, T.A. and W.A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.

Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.

Prior, J.C. 1991. Landforms of Iowa. University of Iowa Press for the Iowa Department of Natural Resources, Iowa City, IA. 153 pps.

Prior, J.C., J.L. Boekhoff, M.R. Howes, R.D. Libra, and P.E. VanDorpe. 2003. Iowa's Groundwater Basics: A Geological Guide to the Occurrence, Use, & Vulnerability of Iowa's Aquifers. Iowa Department of Natural Resources, Iowa Geological Survey Educational Series 6. 92 pps.

Pyne, S.J., P.L. Andrews, and R.D. Laven. 1996. Introduction to Wildland Fire, Second Edition. John Wiley and Sons, Inc. New York, New York. 808 pps.

Rosburg, T. 1994. Community and Physiological Ecology of Native Grasslands in the Loess Hills of Western Iowa. PhD Dissertation. Iowa State University, Ames, IA. 228 pps.

Smith, D.D. 1998. Iowa prairie: original extent and loss, preservation, and recovery attempts. *The Journal of the Iowa Academy of Sciences* 105: 94-108.

Smith, D.D., D. Williams, G. Houseal, and K. Henderson. 2010. The Tallgrass Prairie Center Guide to Prairie Restoration in the Upper Midwest. University of Iowa Press, Iowa City, IA. 338 pps.

Society for Ecological Restoration [SER] Science & Policy Working Group. 2002. The SER Primer on Ecological Restoration. Available at: <http://www.ser.org/>. (Accessed 28 February 2017).

Stambaugh, M.C., R.P. Guyette, E.R. McMurry, and D.C. Dey. 2006. Fire history at the Eastern Great Plains Margin, Missouri River Loess Hills. *Great Plains Research* 16: 149-59.

Steinauer, E.M. and L. Collins. 1996. Prairie ecology: the tallgrass prairie. In: Samson, F.B. and F.L. Knopf, eds. *Prairie Conservation: Preserving North America's Most Endangered Ecosystem*. Island Press, Washington, D.C. 351 pps.

Steinauer, G. and S. Rolfsmeier. 2010. Terrestrial Natural Communities of Nebraska, Version IV. Unpublished report of the Nebraska Game and Parks Commission. Lincoln, NE. 224 pps.

Stockton, C.W. and D.M. Meko. 1983. Drought recurrence in the Great Plains as reconstructed from long-term tree-ring records. *Journal of Climate and Applied Meteorology* 22: 17-29.

Tomer, M.D., D.W. Meek, and L.A. Kramer. 2005. Agricultural practices influence flow regimes of headwater streams in western Iowa. *Journal of Environmental Quality* 34: 1547-1558.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2003. National Range and Pasture Handbook, Revision 1. Grazing Lands Technology Institute. 214 pps.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. 682 pps.

U.S. Environmental Protection Agency [EPA]. 2013. Level III and Level IV Ecoregions of the Continental United States. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>. (Accessed 1 March 2017).

U.S. Geological Survey. 1999. Geology of the Loess Hills, Iowa. Information Handout, July. U.S. Department of the Interior, U.S. Geological Survey. Available at <https://pubs.usgs.gov/info/loess/>. (Accessed 27 February 2017).

Visher, S.S. 1954. Climatic Atlas of the United States. Harvard University Press, Cambridge, MA. 403pps.

Vogl, R.J. 1974. Effects of fire on grasslands. In: T.T. Kozlowski and C.E. Ahlgren, eds. Fire and Ecosystems. Academic Press, New York, New York.

White, J.A. 1983. Regional and Local Variation in Composition and Structure of the Tallgrass Prairie Vegetation of Iowa and Eastern Nebraska. M.S. Thesis. Iowa State University, Ames, IA. 168 pps.

White, J. 1994. How the terms savanna, barrens, and oak openings were used in early Illinois. In: J. Fralisch, ed. Proceedings of the North American Conference on Barrens and Savannas. Illinois State University, Normal, IL.

White, J.A. and S.C. Glenn-Lewin. 1984. Regional and local variation in tallgrass prairie remnants of Iowa and eastern Nebraska. *Vegetatio* 57: 65-78.

Williams, D.A., L.L. Jackson, and D.D. Smith. 2007. Effects of frequent mowing on survival and persistence of forbs seeded into a species-poor grassland. *Restoration Ecology* 15: 24-33.

Wilsey, B.J. 2008. Productivity and subordinate species response to dominant grass species and seed source during restoration. *Restoration Ecology* 18: 628-637.

Yao, J., R.D. Holt, P.M. Rich, and W.S. Marshall. 1999. Woody plant colonization in an experimentally fragmented landscape. *Ecography* 22: 715-728.

## **Approval**

Chris Tecklenburg, 5/21/2020

## **Acknowledgments**

This project could not have been completed without the dedication and commitment from a variety of partners and staff (Table 6). Team members supported the project by serving on the technical team, assisting with the development of state and community phases of the state-and-transition model, providing peer review and technical editing, and conducting quality control and quality assurance reviews.

Organization Name Title Location

Drake University:

Dr. Tom Rosburg Professor of Ecology and Botany Des Moines, IA

Iowa Department of Natural Resources:

Lindsey Barney District Forester Oakland, IA

John Pearson Ecologist Des Moines, IA

LANDFIRE (The Nature Conservancy):

Randy Swaty Ecologist Evanston, IL

Natural Resources Conservation Service:

Rick Bednarek IA State Soil Scientist Des Moines, IA

Stacey Clark Regional Ecological Site Specialist St. Paul, MN

Tonie Endres Senior Regional Soil Scientist Indianapolis, IA

John Hammerly Soil Data Quality Specialist Indianapolis, IN

Lisa Kluesner Ecological Site Specialist Waverly, IA

Sean Kluesner Earth Team Volunteer Waverly, IA

Jeff Matthias State Grassland Specialist Des Moines, IA

Kevin Norwood Soil Survey Regional Director Indianapolis, IN

Doug Oelmann Soil Scientist Des Moines, IA

James Phillips GIS Specialist Des Moines, IA  
Dan Pulido Soil Survey Leader Atlantic, IA  
Melvin Simmons Soil Survey Leader Gallatin, MO  
Tyler Staggs Ecological Site Specialist Indianapolis, IN  
Jason Steele Area Resource Soil Scientist Fairfield, IA  
Doug Wallace Ecological Site Specialist Columbia, MO

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Lisa Kluesner
Contact for lead author	
Date	02/08/2025
Approved by	Chris Tecklenburg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 
17. **Perennial plant reproductive capability:**
-