

Ecological site R109XY023MO

Shallow Limestone Backslope Glade/Woodland

Accessed: 05/09/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 109X—Iowa and Missouri Heavy Till Plain

The Iowa and Missouri Heavy Till Plain (area outlined in red on the map) is an area of rolling hills interspersed with interfluvial divides and alluvial valleys. Elevation ranges from about 660 feet along the lower reaches of rivers, to about 980 feet on stable interfluvial summits in southern Iowa. Relief is about 80 to 160 feet between major streams and adjacent interfluvial summits. Most of the till plain drains south to the Missouri River via the Grand and Chariton River systems, but the northeastern portion drains southeast to the Mississippi River. Loess caps the pre-Illinoian aged till on interfluvial divides, whereas the till is exposed on side slopes. Mississippian aged limestone and Pennsylvanian aged sandstone and shale crop out on lower slopes in some areas.

Classification relationships

Terrestrial Natural Community Type in Missouri (Nelson, 2010):

The reference state for this ecological site is most similar to a Limestone Glade.

Missouri Department of Conservation Forest and Woodland Communities (Missouri Department of Conservation, 2006):

The reference state for this ecological site is most similar to a Limestone/Dolomite Woodland.

National Vegetation Classification System Vegetation Association (NatureServe, 2010):

The reference state for this ecological site is most similar to *Schizachyrium scoparium* - *Bouteloua curtipendula* - *Rudbeckia missouriensis* - *Mentzelia oligosperma* Wooded Herbaceous Vegetation (CEGL002251).

Geographic relationship to the Missouri Ecological Classification System (Nigh & Schroeder, 2002):
 This ecological site occurs in several Land Type Associations, primarily within the following Subsections:
 Grand River Hills
 Loess Hills

Ecological site concept

NOTE: This is a “provisional” Ecological Site Description (ESD) that is under development. It contains basic ecological information that can be used for conservation planning, application and land management. As additional information is collected, analyzed and reviewed, this ESD will be refined and published as “Approved”.

Shallow Limestone Backslope Glade/Woodlands are within the green areas on the map. These sites are in scattered locations mainly in the western and northeastern parts of the MLRA and adjacent areas. Soils are shallow to limestone bedrock. The reference plant community ranges from open areas of grasses and forbs interspersed with bare bedrock, to areas with shrubs and widely scattered chinkapin and post oak.

Associated sites

F109XY030MO	Loamy Floodplain Forest Loamy Floodplain Forests and other floodplain ecological sites are downslope.
R109XY008MO	Till Backslope Savanna Till Backslope Savannas are upslope, on steep lower backslopes.
R109XY046MO	Till Upland Savanna Till Upland Savannas are upslope, on gently sloping upper backslopes.

Similar sites

R109XY023MO	Shallow Limestone Backslope Glade/Woodland There are no similar ecological sites in this MLRA.
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Quercus muehlenbergii</i>
Shrub	(1) <i>Rhus copallina</i>
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Bouteloua curtipendula</i>

Physiographic features

This site is on upland backslopes with slopes of 14 to 50 percent . The site receives runoff from upslope summit and shoulder sites, and generates runoff to adjacent, downslope ecological sites. This site does not flood.

The following figure (adapted from Minor & Davis, 1983) shows the typical landscape position of this ecological site, and landscape relationships among the major ecological sites of the uplands and adjacent floodplains. The site is within the area labeled “4”, and is typically downslope from Till Upland and Backslope Savanna ecological sites. In most areas, Upland Drainageway or Terrace and Floodplain ecological sites are directly downslope.




Figure 2. Landscape relationships for this ecological site

Table 2. Representative physiographic features

Landforms	(1) Hill
Flooding frequency	None
Ponding frequency	None
Slope	14–50%
Aspect	Aspect is not a significant factor

Climatic features

The Iowa and Missouri Heavy Till Plain MLRA has a continental type of climate marked by strong seasonality. In winter, dry-cold air masses, unchallenged by any topographic barriers, periodically swing south from the northern plains and Canada. If they invade reasonably humid air, snowfall and rainfall result. In summer, moist, warm air masses, equally unchallenged by topographic barriers, swing north from the Gulf of Mexico and can produce abundant amounts of rain, either by fronts or by convectional processes. In some summers, high pressure stagnates over the region, creating extended droughty periods. Spring and fall are transitional seasons when abrupt changes in temperature and precipitation may occur due to successive, fast-moving fronts separating contrasting air masses.

This MLRA experiences small regional differences in climates that grade inconspicuously into each other. The basic gradient for most climatic characteristics is along a line from north to south. Both mean annual temperature and precipitation exhibit fairly minor gradients along this line.

Mean January minimum temperature follows the north-to-south gradient. However, mean July maximum temperature shows hardly any geographic variation in the region. Mean July maximum temperatures have a range of only two to three degrees across the region.

Mean annual precipitation varies along the same gradient as temperature – lower annual precipitation in the north, higher in the south. Seasonality in precipitation is very pronounced due to strong continental influences. June precipitation, for example, averages four to five times greater than January precipitation.

During years when precipitation comes in a fairly normal manner, moisture is stored in the top layers of the soil during the winter and early spring, when evaporation and transpiration are low. During the summer months the loss of water by evaporation and transpiration is high, and if rainfall fails to occur at frequent intervals, drought will result. Drought directly influences ecological communities by limiting water supplies, especially at times of high temperatures and high evaporation rates. Drought indirectly affects ecological communities by increasing plant and animal susceptibility to the probability and severity of fire. Frequent fires encourage the development of grass/forb dominated communities and understories.

Superimposed upon the basic MLRA climatic patterns are local topographic influences that create topoclimatic, or microclimatic variations. For example, air drainage at nighttime may produce temperatures several degrees lower in valley bottoms than on side slopes. At critical times during the year, this phenomenon may produce later spring or

earlier fall freezes in valley bottoms. Slope orientation is an important topographic influence on climate. Summits and south-and-west-facing slopes are regularly warmer and drier, supporting more grass dominated communities than adjacent north- and-east-facing slopes that are cooler and moister that support more woody dominated communities. Finally, the cooler microclimate within a canopied forest is measurably different from the climate of a more open and warmer grassland or savanna area.

Source: University of Missouri Climate Center - <http://climate.missouri.edu/climate.php>; Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin, United States Department of Agriculture Handbook 296 - <http://soils.usda.gov/survey/geography/mlra/>

Table 3. Representative climatic features

Frost-free period (average)	163 days
Freeze-free period (average)	186 days
Precipitation total (average)	1,067 mm

Climate stations used

- (1) DONNELLSON [USC00132299], Donnellson, IA
- (2) BETHANY [USC00230608], Bethany, MO
- (3) KEARNEY 3E [USC00234382], Kearney, MO

Influencing water features

This ecological site is not influenced by wetland or riparian water features.

Soil features

These soils are underlain with limestone bedrock at less than 20 inches. The soils were formed under prairie vegetation, and have dark, organic-rich surface horizons. Parent material is limestone residuum. These soils are clayey and are skeletal, with high amounts of limestone gravel, channers and flagstones. They are not affected by seasonal wetness. Soil series associated with this site include Gasconade, Nordness, and Sogn.

Table 4. Representative soil features

Parent material	(1) Residuum–limestone
Surface texture	(1) Flaggy silty clay loam
Family particle size	(1) Clayey
Drainage class	Somewhat excessively drained
Permeability class	Very slow to slow
Soil depth	10–51 cm
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	5–40%
Available water capacity (0-101.6cm)	2.54–5.08 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0

Soil reaction (1:1 water) (0-101.6cm)	6.1–7.8
Subsurface fragment volume <=3" (Depth not specified)	0–5%
Subsurface fragment volume >3" (Depth not specified)	35–50%

Ecological dynamics

Information contained in this section was developed using historical data, professional experience, field reviews, and scientific studies. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The Reference Plant Community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

Limestone glades are open, rocky areas with very shallow soils dominated by drought-adapted herbaceous flora, generally occurring on south-and west-facing slopes of otherwise wooded sites (Nelson 2010). One of the most striking aspects of dolomite glades is their unique and characteristic flora. Many plants have an affinity for calcareous dolomite substrates. Glade plants in general possess many adaptations enabling them to survive in a harsh environment often subject to widely fluctuating extremes of temperature and moisture. The following conditions are general characteristic of most limestone glades (Nelson and Ladd 1983; Nelson et al. 2013):

- Calcareous limestone bedrock at or near the surface as a result of major erosional activity and resistance to weathering;
- Moderate to steep slopes in deeply dissected drainages or hilly to mountainous terrain with a southern or western exposure with intense solar radiation;
- Extremely thin soil cover interspersed with abundant rock fragments and rock outcrops;
- Exceptionally dry conditions throughout much of the growing season, although soils may be seasonally saturated in spring, winter, and fall;
- Peripheral areas and sometimes large expanses of the glades themselves characterized by a mosaic of stunted, often gnarled trees and shrubs.

Shallow Limestone Backslope Glade/Woodland ecological sites harbor a wide diversity of plants and animals. Grasses such as little bluestem, Indian grass, and sideoats grama, are also found on upland prairies. The shallow soils of this ecological site limit the growth and abundance of trees and support the native grasses and forbs that dominate these systems. Fire played an important role in the maintenance of these systems, as well. It is likely that these sites burned at least once every five years.


These periodic fires removed the litter and stimulated the growth and flowering of the grasses and forbs. They also further limited the growth and dominance of trees, especially eastern redcedar. Fire tolerant chinkapin oak and post oak occupied islands and edges where the deeper range of the soil component occurred, creating a complex mosaic of open glade and low-density woodland. During fire-free intervals, woody species increased, but not to densities on over-grazed glades.

These sites are not productive. In the absence of fire, woody species, especially eastern redcedar, quickly occupy the site. This is especially true after grazing has reduced grass cover and exposed more surface to the dispersal of cedar seeds by birds. Once established, cedars can quickly fill in a glade/woodland system, especially if grazing has diminished the vigor of the diverse flora. Many glades have been heavily grazed and suffer substantial redcedar invasion. Removal of the redcedar and the application of prescribed fire have proven to be an effective way to manage these systems.

A State and Transition Diagram follows. Detailed descriptions of each state, transition, plant community, and pathway follow the model. This model is based on available experimental research, field observations, professional consensus, and interpretations. It is likely to change as knowledge increases.

State and transition model

Shallow Limestone Backslope Glade/Woodland, R109XY023MO



Code	Event/Activity
T1A	Fire suppression (> 20 years)
T1B	Uncontrolled grazing; fire suppression
T3A	Uncontrolled grazing
R2A	Cedar removal; prescribed fire
R3A	Grazing exclusion; prescribed fire; woody removal
1.1A	Fire-free interval (10-20 years)
1.2A	Fire interval (3-10 years)

Figure 7. State & Transition Model for this ecological site.

State 1

Reference

Glade/Woodland Reference sites harbor a wide diversity of plants and animals. Many, like little bluestem, Indian grass, and sideoats grama, are also found on upland prairies. The glade/woodland complexes range from wide open grassy areas with shallow soils and bare bedrock, to areas with widely scattered chinkapin and post oaks on locations with soil depths at the deeper extreme of the range for this soil component. On protected slopes, open woodlands with chinkapin and post oaks are more common. Here the deeper soil depth range for this soil component and protected aspects allow more woody components to dominate. As fire frequencies decrease invasion by woody species such as eastern redcedar will begin to increase and occupy more space on the site. Return to normal historical fire frequencies will reduce this woody increase and re-open up the site. While many glades have suffered from grazing and fire suppression, good examples can still be found.

Community 1.1

Chinkapin Oak-Eastern Redcedar/Winged Sumac/Little Bluestem

This phase has widely scattered chinkapin oak and post oak with little bluestem, side oats grama, and dropseeds dominating the open ground layer. Numerous forbs such as Missouri coneflower and lichens are also present and locally abundant. Bedrock outcropping is common.

Forest overstory. The Forest Overstory Species list is based on commonly occurring species listed in Nelson

Forest understory. The Forest Understory list is based on commonly occurring species listed in Nelson (2010).

Community 1.2

Chinkapin Oak/Winged Sumac-Eastern Redcedar/Little Bluestem-Sideoats Grama

This phase is similar to community phase 1.1 but post oak, chinkapin oak and numerous shrubs are increasing due to longer periods of fire suppression. Some displacement of grasses and forbs may be occurring due to shading and competition from the increased densities of shrubs and oaks.

State 2

Woody Invaded Glade/Woodland

This state is dominated by eastern redcedar with large increases of oak density due to extended periods of fire suppression. This State can form relatively even-age stands, dating to when fire suppression became the dominant management characteristic on the site. Canopy closures can approach 100% with little or no ground flora. Transition back to the Reference State may require a number of prescribed fire events and thinning out of excess woody species. This state also can transition to a grazed state (State 3) with the introduction of domestic livestock.

Community 2.1

Chinkapin Oak-Eastern Redcedar/Winged Sumac/Little Bluestem

State 3

Grazed Glade/Woodland

The Grazed Glade/Woodland State has reduced cover, diversity and vigor of native glade/woodland flora. Woody species encroachment, particularly by eastern redcedar, has also increased in this State. Potential physical site damage by uncontrolled livestock grazing may further degrade this State.

Community 3.1

Chinkapin Oak-Eastern Redcedar/Winged Sumac/Broomsedge

Additional community tables

Table 5. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)	Diameter (Cm)	Basal Area (Square M/Hectare)
Tree							
chinquapin oak	QUMU	<i>Quercus muehlenbergii</i>	Native	–	5–20	–	–
post oak	QUST	<i>Quercus stellata</i>	Native	–	5–20	–	–
eastern redcedar	JUVI	<i>Juniperus virginiana</i>	Native	–	5–20	–	–

Table 6. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	Native	–	10–30
little bluestem	SCSC	<i>Schizachyrium scoparium</i>	Native	–	10–30
hairy woodland brome	BRPU6	<i>Bromus pubescens</i>	Native	–	10–20
slender woodland sedge	CADI5	<i>Carex digitalis</i>	Native	–	10–20
Indiangrass	SONU2	<i>Sorghastrum nutans</i>	Native	–	5–20
puffsheath dropseed	SPNE2	<i>Sporobolus neglectus</i>	Native	–	5–10
Mead's sedge	CAME2	<i>Carex meadii</i>	Native	–	5–10
Forb/Herb					
western silver aster	SYSE2	<i>Symphotrichum sericeum</i>	Native	–	10–20
Carolina larkspur	DECA3	<i>Delphinium carolinianum</i>	Native	–	10–20
butterfly milkweed	ASTU	<i>Asclepias tuberosa</i>	Native	–	10–20
slimflower scurfpea	PSTE5	<i>Psoraleidium tenuiflorum</i>	Native	–	5–20
tall blazing star	LIAS	<i>Liatris aspera</i>	Native	–	5–20
common goldstar	HYHI2	<i>Hypoxis hirsuta</i>	Native	–	5–20
diamondflowers	STNIN	<i>Stenaria nigricans var. nigricans</i>	Native	–	5–20
prairie rosinweed	SITE	<i>Silphium terebinthinaceum</i>	Native	–	5–20
Michaux's croton	CRMIE	<i>Croton michauxii var. ellipticus</i>	Native	–	5–20
prairie tea	CRMO6	<i>Croton monanthogynus</i>	Native	–	5–20
hoary puccoon	LICA12	<i>Lithospermum canescens</i>	Native	–	10–20
hairy sunflower	HEHI2	<i>Helianthus hirsutus</i>	Native	–	10–20
Shrub/Subshrub					
winged sumac	RHCO	<i>Rhus copallinum</i>	Native	–	5–20
fragrant sumac	RHAR4	<i>Rhus aromatica</i>	Native	–	5–10
dwarf hackberry	CETE	<i>Celtis tenuifolia</i>	Native	–	5–10
Nonvascular					
fishscale lichen	PSDE60	<i>Psora decipiens</i>	Native	–	5–20
sarcogyne lichen	SARE18	<i>Sarcogyne regularis</i>	Native	–	5–20
Russell's fishscale lichen	PSRU3	<i>Psora russellii</i>	Native	–	5–20
	THCO12	<i>Thyrea confusa</i>	Native	–	5–20

Table 7. Community 2.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)
Grass/grass-like (Graminoids)					
little bluestem	SCSC	<i>Schizachyrium scoparium</i>	Native	–	–
Indiangrass	SONU2	<i>Sorghastrum nutans</i>	Native	–	–
sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	Native	–	–
Forb/Herb					
Missouri orange coneflower	RUMI	<i>Rudbeckia missouriensis</i>	Native	–	–

Animal community

Wildlife

Oaks provide hard mast; scattered shrubs provide soft mast; frequent bedrock outcrops provide reptile habitat and a patchier groundflora; sedges and native cool-season grasses provide green browse;

Native warm-season grasses on dry sites provide cover and nesting habitat; and a diversity of forbs provides a diversity and abundance of insects.

Post-burn areas can provide temporary bare-ground – herbaceous cover habitat important for turkey poults and quail chicks.

Game species that utilize this ecological site include: Northern Bobwhite will utilize this ecological site for food (seeds, insects), cover needs (escape, nesting and roosting cover) and brood-rearing habitat.

Cottontail rabbits will utilize this ecological site for food (seeds, soft mast) and cover needs.

Turkey will utilize this ecological site for food (seeds, green browse, soft mast, and insects) and nesting and brood-rearing cover. Turkey poults feed heavily on insects provided by this site type.

White-tailed Deer will utilize this ecological site for browse (plant leaves in the growing season, seeds and soft mast in the fall/winter). This site type also can provide escape cover.

Bird species associated with this ecological site's reference state condition: Field Sparrow, Yellow-breasted Chat, Brown Thrasher, Indigo Bunting, Red-headed Woodpecker, Eastern Bluebird, Northern Bobwhite, and Eastern Wood-Pewee.

Amphibian and reptile species that may be associated with this ecological site's reference state: Five-lined Skink (*Eumeces fasciatus*), Six-lined Racerunner (*Cnemidophorus sexlineatus*), and Prairie Ring-necked Snake (*Diadophis punctatus arnyi*).

Invertebrates – Many native insect species are likely associated with this phase of this ecological site's reference state condition, especially native bees, ants, beetles, butterflies and moths, and crickets, grasshoppers and katydids. However we don't have enough information on these groups to assign them to this ecological site's reference state condition at this time.

(References used in this section: Fitzgerald and Pashley. 2000b; Heitzman and Heitzman 1996; Jacobs 2001; Johnson 2000; Pitts and McGuire 2000; Schwartz and others 2001; MDC 2006)

Other information

Forestry

Management: Site index values are less than 30 for eastern redcedar and generally less than 40 for oak.

Productivity is very low. Very limited timber management opportunities exist. These sites are valuable for wildlife purposes and watershed protection. Severely reduced rooting depth restricts tree growth. These sites respond well to prescribed fire as a management tool.

Limitations: Surface stones and surface rock; very shallow soil depth. Surface stones and rocks are problems for

efficient and safe equipment operation. Severe seedling mortality due to high soil surface temperatures and low available water holding capacity is possible. Machine planting and mechanical site preparation is not recommended. Hard bedrock at shallow depths may interfere with equipment operation. Rock outcrops may cause breakage of timber when harvesting. Surface stones and rocks will make equipment use extremely difficult. Erosion is a hazard when slopes exceed 15 percent. On steep slopes greater than 35 percent, traction problems increase and equipment use is not recommended.

Inventory data references

EXTINCT in reference state - no plots

Other references

Fitzgerald, J.A. and D.N. Pashley. 2000a. Partners in Flight bird conservation plan for the Ozark/Ouachitas. American Bird Conservancy.

Fitzgerald, J.A. and D.N. Pashley. 2000b. Partners in Flight bird conservation plan for the Dissected Till Plains. American Bird Conservancy.

Heitzman, J.R. and J.E. Heitzman. 1996. Butterflies and moths of Missouri. 2nd ed. Missouri Department of Conservation, Jefferson City.

Jacobs, B. 2001. Birds in Missouri. Missouri Department of Conservation, Jefferson City.

Johnson, T.R. 2000. The amphibians and reptiles of Missouri. 2nd ed. Missouri Department of Conservation, Jefferson City.

Minor, Paul E., & Keith O. Davis. 1983. Soil Survey of Clinton County, Missouri. U.S. Dept. of Agric. Soil Conservation Service.

Missouri Department of Conservation. 2006. Missouri Forest and Woodland Community Profiles. Missouri Department of Conservation, Jefferson City, Missouri.

NatureServe. 2010. Vegetation Associations of Missouri (revised). NatureServe, St. Paul, Minnesota.

Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.

Nelson, Paul W and Douglas Ladd. 1980. "Preliminary report on the identification, distribution and classification of Missouri glades".

Nelson, P. W., J. A. Fitzgerald, K. Larson, R. McCoy, A. Schotz, J. Taft, T. Witsell, B. Yahn. 2013. Central Hardwoods Joint Venture Glade Conservation Assessment for the Interior Highlands and Interior Low Plateaus of the Central Hardwoods Region. Central Hardwoods Joint Venture.

Nigh, Timothy A., and Walter A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.

Pitts, D.E. and W.D. McGuire. 2000. Wildlife management for Missouri landowners. 3rd ed. Missouri Department of Conservation, Jefferson City.

Schwartz, C.W., E.R. Schwartz and J.J. Conley. 2001. The wild mammals of Missouri. University of Missouri Press, Columbia and Missouri Department of Conservation, Jefferson City.

Contributors

Doug Wallace
Fred Young

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
