

Ecological site F111XA018IN

Shallow Restricted

Last updated: 4/20/2020
Accessed: 05/05/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 111X–Indiana and Ohio Till Plain

A PROVISIONAL ECOLOGICAL SITE is a conceptual grouping of soil map unit components within a Major Land Resource Area (MLRA) based on the similarities in response to management. Although there may be wide variability in the productivity of the soils grouped into a Provisional Site, the soil vegetation interactions as expressed in the State and Transition Model are similar and the management actions required to achieve objectives, whether maintaining the existing ecological state or managing for an alternative state, are similar. Provisional Sites are likely to be refined into more precise group during the process of meeting the APPROVED ECOLOGICAL SITE DESCRIPTION criteria.

This PROVISIONAL ECOLOGICAL SITE has been developed to meet the standards established in the National Ecological Site Handbook. The information associated with this ecological site does not meet the Approved Ecological Site Description Standard, but it has been through a Quality Control and Quality Assurance processes to assure consistency and completeness. Further investigations, reviews and correlations are necessary before it becomes an Approved Ecological Site Description.

111A – Indiana and Ohio Till Plain, Central Part. This area is in the Till Plains Section of the Central Lowland Province of the Interior Plains. It is dominated by broad, nearly level ground moraines that are broken in some areas by kames, outwash plains, and stream valleys along the leading edge of the moraines. Narrow, shallow valleys commonly are along the few large streams in the area. Elevation ranges from 680 to 1,250 feet (205 to 380 meters), increasing gradually from west to east. Relief is mainly a few meters, but in some areas hills rise as much as 100 feet (30 meters) above the adjoining plains.

The extent of the major Hydrologic Unit Areas (identified by four-digit numbers) that make up this MLRA is as follows: Wabash (0512), 46 percent; Great Miami (0508), 30 percent; Scioto (0506), 22 percent; and the Middle Ohic (0509), 2 percent. The major rivers in the area include the East and West Forks of the White River and the Whitewater River in Indiana and the Great Miami, Stillwater, Big Darby, Scioto, and Big Walnut Rivers in Ohio.

Surface deposits in this area include glacial deposits of till, lacustrine sediments, and outwash from Wisconsin and older glacial periods. A moderately thick mantle of loess covers much of the area. Most of this MLRA is underlain by Silurian and Devonian limestone and dolostone. Also, some areas of Late Ordovician shale and limestone are in the western part of the MLRA (USDA, 2006).

Classification relationships

Major Land Resource Area (USDA-Natural Resources Conservation Service, 2006)

USFS Ecological Regions (USDA, 2007):

Sections – Southern Unglaciaded Allegheny Plateau (221E), Central Till Plains, Beech Maple (222H), Interior Low Plateau-Transition Hills (223B), Interior Low Plateau-Bluegrass (223F)

Subsections - Lower Scioto River Plateau (221Eg), Bluffton Till Plains (222Ha), Miami-Scioto Plain-Tipton Till Plain

(222Hb), Little Miami Old Drift Plain (222Hc), Mad River Interlobate Plains (222Hd), Darby Plains (222He), Brown County Hills (223Ba), Northern Bluegrass (223Fd), Muscatatuck Flats and Valleys (223Fe), Scottsburg Lowlands (223Ff)

NatureServe Systems anticipated (NatureServe, 2011): Agriculture - Cultivated Crops and Irrigated Agriculture, Agriculture - Pasture/Hay, Allegheny-Cumberland Dry Oak Forest and Woodland, Appalachian (Hemlock)-Northern Hardwood Forest, Central Appalachian Pine-Oak Rocky Woodland, Central Interior Acidic Cliff and Talus, Central Interior Highlands Calcareous Glade and Barrens, Central Tallgrass Prairie, Clearcut - Grassland/Herbaceous, Introduced Upland Vegetation – Treed, Managed Tree Plantation, Mississippi River Riparian Forest, North-Central Interior and Appalachian Acidic Peatland, North-Central Interior Beech-Maple Forest, North-Central Interior Dry-Mesic Oak Forest and Woodland, North-Central Interior Floodplain, North-Central Interior Freshwater Marsh, North-Central Interior Oak Savanna, North-Central Interior Wet Flatwoods, North-Central Interior Wet Meadow-Shrub Swamp, North-Central Oak Barrens, Northeastern Interior Dry-Mesic Oak Forest, Ruderal Forest, Ruderal Upland - Old Field, South-Central Interior / Upper Coastal Plain Wet Flatwoods, South-Central Interior Large Floodplain, South-Central Interior Mesophytic Forest, South-Central Interior Small Stream and Riparian, Southern Appalachian Oak Forest, Southern Interior Low Plateau Dry-Mesic Oak Forest, Southern Ridge and Valley / Cumberland Dry Calcareous Forest, Successional Shrub/Scrub

LANDFIRE Biophysical Settings anticipated (USGS, 2010): Allegheny-Cumberland Dry Oak Forest and Woodland, Appalachian (Hemlock-) Northern Hardwood Forest, Central Interior and Appalachian Floodplain Systems, Central Interior and Appalachian Riparian Systems, Central Interior and Appalachian Shrub-Herbaceous Wetland Systems, Central Interior and Appalachian Swamp Systems, Central Interior Highlands Calcareous Glade and Barrens, Central Interior Highlands Dry Acidic Glade and Barrens, Central Tallgrass Prairie, Great Lakes Coastal Marsh Systems, North-Central Interior Beech-Maple Forest, North-Central Interior Dry-Mesic Oak Forest and Woodland, North-Central Interior Dry Oak Forest and Woodland, North-Central Interior Oak Savanna, North-Central Interior Wet Flatwoods, South-Central Interior Mesophytic Forest, South-Central Interior/Upper Coastal Plain Flatwoods, Southern Appalachian Oak Forest, Southern Interior Low Plateau Dry-Mesic Oak Forest

Ecological site concept

This site is an upland site formed on clayey residium overlaying limestone and shale. The depth to the restrictive layer is less than 20 inches. Drainage is well drained or drier with slopes up generally from 25-80%. Landscape position, shallow soil depth, and slope provide a level of protection for the site from fire. This combination of factors allow the eastern red cedar to become a co-dominant on the site with oak species. The dark soil surface color and depth of the color, along with the drainage of the site indicate that the understory was comprised largely of herbaceous, mostly graminoid, species. A large portion of the site in agricultural use is used for pasture or hay. The majority of the site is in natural vegetation, with an appreciable amount being in second growth woodlands or brush.

Associated sites

F111XA019IN	Moderately Deep Restricted Site is generally lower on the landscape; soil depth to restrictive layer is between 20 and 40 inches
F111XA020IN	Deep Restricted Site is generally lower on the landscape; soil depth to restrictive layer is 40 inches or greater




Figure 1. Block diagram showing soil series on the landscape.

Table 1. Dominant plant species

Tree	(1) <i>Quercus</i> (2) <i>Juniperus virginiana</i>
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This ecosite is found in unspecified landscape in MLRA 111A: Indiana and Ohio Till Plain, Central Part. This site is shallow (>20 inches) to impermeable layer.




Figure 2. Block diagram showing soil series on the landscape.

Table 2. Representative physiographic features

Landforms	(1) Hill
Flooding frequency	None
Ponding frequency	None
Elevation	91–488 m
Slope	6–80%
Ponding depth	0 cm
Water table depth	152 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

The average annual precipitation in this area is 36 to 43 inches (915 to 1,090 millimeters). Most of the rainfall occurs as convective thunderstorms during the growing season. About half or more of the precipitation occurs during the freeze-free period. Snowfall is common in winter. The average annual temperature is 49 to 53 degrees F (9 to 12 degrees C). The freeze-free period averages about 195 days and ranges from 175 to 215 days.

Table 3. Representative climatic features

Frost-free period (characteristic range)	145-158 days
Freeze-free period (characteristic range)	174-191 days
Precipitation total (characteristic range)	1,016-1,092 mm

Frost-free period (actual range)	142-165 days
Freeze-free period (actual range)	171-198 days
Precipitation total (actual range)	991-1,118 mm
Frost-free period (average)	152 days
Freeze-free period (average)	183 days
Precipitation total (average)	1,041 mm




Figure 3. Monthly precipitation range




Figure 4. Monthly minimum temperature range




Figure 5. Monthly maximum temperature range




Figure 6. Monthly average minimum and maximum temperature




Figure 7. Annual precipitation pattern




Figure 8. Annual average temperature pattern

Climate stations used

- (1) WASHINGTON COURT HOUSE [USC00338794], Washington Court House, OH
- (2) FRANKLIN [USC00332928], Franklin, OH
- (3) INDIANAPOLIS SE SIDE [USC00124272], Indianapolis, IN
- (4) KOKOMO 3 WSW [USC00124662], Russiaville, IN
- (5) COLUMBUS [USC00121747], Columbus, IN
- (6) RICHMOND WTR WKS [USC00127370], Richmond, IN
- (7) CHILLICOTHE MOUND CITY [USC00331528], Chillicothe, OH
- (8) COLUMBUS OHIO STATE UNIV AP [USW00004804], Dublin, OH

Influencing water features

This ecological site is not influenced by wetland or riparian water features.

Soil features

The soil series associated with this site are: Weikert, Rohan, Opequon, Gasconade, Corydon. They are shallow, well drained to somewhat excessively drained, and very slow to rapid permeable soils, with very strongly acidic to

neutral soil reaction, that formed in Residuum from Limestone, Limestone and shale, Sandstone and shale, Shale.




Figure 9. Mapunit location in the MLRA.

Table 4. Representative soil features

Parent material	(1) Residuum–limestone and shale
Surface texture	(1) Channery clay (2) Flaggy silt loam (3) Stony silty clay loam
Family particle size	(1) Clayey
Drainage class	Well drained to somewhat excessively drained
Permeability class	Very slow to rapid
Soil depth	38–48 cm
Surface fragment cover <=3"	0–3%
Surface fragment cover >3"	3%
Available water capacity (0-101.6cm)	3.56–6.86 cm
Calcium carbonate equivalent (0-101.6cm)	0–15%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	5.2–7.2
Subsurface fragment volume <=3" (Depth not specified)	1–33%
Subsurface fragment volume >3" (Depth not specified)	1–40%

Ecological dynamics

The historic plant community of the Shallow Restricted ecological site is that of a barrens that is somewhat protected from fire. The site is mostly wooded with various oak species and eastern red cedar being characteristic of the site. The understory is dominated by herbaceous species, particularly little bluestem and sideoats grama. Low intensity surface fires maintain the herbaceous understory and less frequent, more severe fires keep the site from being dominated by eastern red cedar. Much of this site is currently still in a wooded or old field state. Areas used for agriculture are mostly used for growing forage or pasture.

State and transition model




Figure 10. STM

**Shallow Restricted, F111AY018IN
Diagram Legend**

T1A	Remove woody species, plant forage/pasture, management
T2A	Plant desired species, low intensity fire
T2B	No management
T3A	Plant desired species, low intensity fire
T3B	Remove woody species, plant forage/pasture, management

Figure 11. legend

**State 1
Shallow Restricted woodland**

This is the reference or diagnostic plant community for this site. The site is mostly wooded with various oak species and eastern red cedar being characteristic of the site. The understory is dominated by herbaceous species, particularly little bluestem and sideoats grama.

Dominant plant species

- oak (*Quercus*), tree
- eastern redcedar (*Juniperus virginiana*), tree

Community 1.1

oaks/eastern red cedar/little bluestem

Low intensity surface fires maintain the herbaceous understory and less frequent, more severe fires keep the site from being dominated by eastern red cedar.

Dominant plant species

- oak (*Quercus*), tree
- eastern redcedar (*Juniperus virginiana*), tree
- little bluestem (*Schizachyrium*), grass

State 2

Agriculture

The portion of this site used for agriculture is largely used for pasture or cool season forage production.

Community 2.1

cool season forage/pasture

This phase is characterized by forage or grazing agriculture. Different mixes of, generally, cool season grasses and forbs, largely clovers, are grown.

State 3

Old Field State

This state is characterized by interrupted drainage and removal of historical vegetation. Both contribute to site being comprised of largely weedy species that will progress to tree dominance.

Community 3.1

tall fescue/eastern red cedar

Tall fescue and eastern red cedar colonize the site after agricultural abandonment.

Transition T1A

State 1 to 2

Remove the woody species, plant desired forage species, and pasture/forage management will move the site to and maintain state #2.

Restoration pathway R2A

State 2 to 1

Plant appropriate species, low intensity fires.

Transition T2B

State 2 to 3

Abandonment or no management on agricultural field.

Transition R3A

State 3 to 1

Removal of existing vegetation, planting of desired species, and low intensity fire will move this state towards the reference community.

Restoration pathway T3B

State 3 to 2

Clear woody vegetation, plant crop species, and management that allows the area to be in agriculture.

Additional community tables

Inventory data references

Site concept developed through expert opinion and review of literature.

Other references

Braun, E. Lucy. 2001. Deciduous forests of eastern North America. Caldwell, N.J.: Blackburn Press.

Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. Indiana Academy of Science , 94, 245-269.

NatureServe. (2011). An online encyclopedia of life [web application]. NatureServe, Arlington, VA, USA [Online: [www. natureserve. org/explorer](http://www.natureserve.org/explorer)] .

Jackson, Marion T. 1997. The Natural heritage of Indiana. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

Johnson, Paul S., Stephen R. Shifley, and Robert Rogers. 2002. The ecology and silviculture of oaks. Wallingford, Oxon: CABI

USDA. (2007). Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

USDA. (2006). Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U. S. Department of Agriculture, Natural Resources Conservation Service. U. S. Department of Agriculture Handbook 296.

USGS. (2010). LANDFIRE Biophysical Settings. Retrieved from <http://www.landfire.gov>

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

Approval

Chris Tecklenburg, 4/20/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	TYLER STAGGS
Contact for lead author	

Date	05/05/2024
Approved by	Chris Tecklenburg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-