

Ecological site F111XD013IN

Wet Lacustrine Forest

Last updated: 5/28/2020
Accessed: 05/03/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 111X–Indiana and Ohio Till Plain

A PROVISIONAL ECOLOGICAL SITE is a conceptual grouping of soil map unit components within a Major Land Resource Area (MLRA) based on the similarities in response to management. Although there may be wide variability in the productivity of the soils grouped into a Provisional Site, the soil vegetation interactions as expressed in the State and Transition Model are similar and the management actions required to achieve objectives, whether maintaining the existing ecological state or managing for an alternative state, are similar. Provisional Sites are likely to be refined into more precise group during the process of meeting the APPROVED ECOLOGICAL SITE DESCRIPTION criteria.

This PROVISIONAL ECOLOGICAL SITE has been developed to meet the standards established in the National Ecological Site Handbook. The information associated with this ecological site does not meet the Approved Ecological Site Description Standard, but it has been through a Quality Control and Quality Assurance processes to assure consistency and completeness. Further investigations, reviews and correlations are necessary before it becomes an Approved Ecological Site Description.

111D – Indiana and Ohio Till Plain, Western Part. This MLRA occurs in two separate areas. One area is in the west-central part of Indiana (73 percent), and the other is in southwestern Ohio (27 percent). The MLRA makes up 5,355 square miles (13,880 square kilometers). It includes the towns of Crawfordsville, Delphi, Frankfort, Lafayette, and Liberty, Indiana, and Hamilton, Lebanon, Middletown, and Wilmington, Ohio. Interstates 65 and 74 cross the part of this area in Indiana, and Interstates 71 and 75 cross the part in Ohio. Shades and Turkey Run State Parks are in the part in Indiana, and Caesar Creek and Hueston Woods State Parks are in the part in Ohio. A small portion of the Wright-Patterson Air Force Base, in Ohio, is in the northern part of the area.

This area is in the Till Plains Section of the Central Lowland Province of the Interior Plains. It is dominated by loess hills and flats that are broken in places by moraines, kames, outwash plains, and stream terraces. Narrow, shallow valleys commonly are along the few large streams in the area. Elevation ranges from 530 to 1,050 feet (160 to 320 meters), increasing gradually from southwest to northeast. Relief is mainly a few meters, but in some areas hills rise as much as 100 feet (30 meters) above the adjoining plains.

The extent of the major Hydrologic Unit Areas (identified by four-digit numbers) that make up this MLRA is as follows: Wabash (0512), 68 percent; Great Miami (0508), 15 percent; Middle Ohio (0509), 14 percent; Scioto (0506), 2 percent; and Upper Illinois (0712), 1 percent. Wildcat Creek in Indiana and the Little Miami River in Ohio have been designated as National Wild and Scenic Rivers. Sugar Creek and Walnut Creek occur in the part of the area in northern Indiana, and the Whitewater River is in the part in southeastern Indiana. The Sevenmile, Fourmile, and Great Miami Rivers cross the part of the area in Ohio.

Most of the eastern part of this MLRA is underlain by Late Ordovician shale and limestone. The western part is underlain by shale, siltstone, sandstone, limestone, and dolostone ranging in age from Middle Pennsylvanian to Silurian. Surficial materials include glacial deposits of till, outwash, and lacustrine sediments from Wisconsin and

older glacial periods. A thin or moderately thick mantle of loess overlies much of the area.

Classification relationships

Major Land Resource Area (USDA-Natural Resources Conservation Service, 2006)

USFS Ecological Regions (USDA, 2007):

Sections –Central Till Plains, Beech Maple (222H), Interior Low Plateau-Shawnee Hills (223D), Interior Low Plateau-Bluegrass (223F), Central Till Plains-Oak Hickory (223G), Central Till Plains and Grand Prairies (251D)

Subsections -Bluffton Till Plains (222Ha), Miami-Scioto Plain-Tipton Till Plain (222Hb), Little Miami Old Drift Plain (222Hc), Mad River Interlobate Plains (222Hd), Crawford Uplands (223De), Crawford Escarpment (223Df), Northern Bluegrass (223Fd), Lower Wabash Alluvial Plain (223Gc), Southwest Indiana Glaciated Lowlands (223Ge), Eastern Grand Prairie (253Dd).

NatureServe Systems anticipated (NatureServe, 2011): Agriculture - Cultivated Crops and Irrigated Agriculture, Agriculture - Pasture/Hay, Allegheny-Cumberland Dry Oak Forest and Woodland, Central Interior Acidic Cliff and Talus, Central Interior Highlands Calcareous Glade and Barrens, Central Tallgrass Prairie, Clearcut - Grassland/Herbaceous, Introduced Upland Vegetation – Treed, Managed Tree Plantation, Mississippi River Riparian Forest, North-Central Interior and Appalachian Acidic Peatland, North-Central Interior Beech-Maple Forest, North-Central Interior Dry-Mesic Oak Forest and Woodland, North-Central Interior Dry Oak Forest and Woodland, North-Central Interior Floodplain, North-Central Interior Freshwater Marsh, North-Central Interior Maple-Basswood Forest, North-Central Interior Oak Savanna, North-Central Interior Wet Flatwoods, North-Central Interior Wet Meadow-Shrub Swamp, North-Central Oak Barrens, Northern Atlantic Coastal Plain Hardwood Forest, Ruderal Forest, Ruderal Upland - Old Field, South-Central Interior / Upper Coastal Plain Wet Flatwoods, South-Central Interior Large Floodplain, South-Central Interior Mesophytic Forest, South-Central Interior Small Stream and Riparian, Southern Appalachian Oak Forest, Southern Interior Low Plateau Dry-Mesic Oak Forest, Successional Shrub/Scrub

LANDFIRE Biophysical Settings anticipated (USGS, 2010): Allegheny-Cumberland Dry Oak Forest and Woodland, Bluegrass Savanna and Woodland, Central Interior and Appalachian Floodplain Systems, Central Interior and Appalachian Riparian Systems, Central Interior and Appalachian Shrub-Herbaceous Wetland Systems, Central Interior and Appalachian Swamp Systems, Central Interior Highlands Calcareous Glade and Barrens, Central Interior Highlands Dry Acidic Glade and Barrens, Central Tallgrass Prairie, Great Lakes Coastal Marsh Systems, Mississippi River Alluvial Plain Dry-Mesic Loess Slope Forest, North-Central Interior Beech-Maple Forest, North-Central Interior Dry-Mesic Oak Forest and Woodland, North-Central Interior Dry Oak Forest and Woodland, North-Central Interior Maple-Basswood Forest, North-Central Interior Oak Savanna, North-Central Interior Wet Flatwoods, Paleozoic Plateau Bluff and Talus, Pennyroyal Karst Plain Prairie and Barrens, South-Central Interior Mesophytic Forest, South-Central Interior/Upper Coastal Plain Flatwoods, Southern Appalachian Oak Forest, Southern Interior Low Plateau Dry-Mesic Oak Forest

Ecological site concept

This site is a wetland site formed on lacustrine parent materials. It is located on lake plains, glacial lake relicts, and flats. The soils have very dark surfaces with a subsurface that is in the loamy textural group, and are very poorly or poorly drained. These soils generally occur on nearly level to depressional areas of stream terraces, till plains, and glacial lake plains. The site is occasionally flooded for a brief duration, but ponding occurs frequently to a depth of 30 inches for up to more than a month.

The characteristic vegetation of the site is that of a flatwoods type dominated by pin oak and swamp white oak. Silver maple and green ash are also common in the canopy. The large, seasonal fluctuation of water on the site allows for the co-existence of upland and lowland trees. Inundation of the site generally occurs in the spring which leads to a sparse and patchy understory. Fire did occur on the site, but high intensity fires were rare. Low intensity surface fires were more common on sites that are adjacent to more fire prone sites such as prairies and savannas. Ponding in the spring followed by summer drought along with windthrow were the most dominant disturbance factors. A large portion of this site has been drained and is in agricultural production.

Associated sites

F111XD014IN	Lacustrine Forest Located on similar landscape positions; soils are somewhat poorly to moderately well drained.
-------------	---

Similar sites

F111XD015IN	Wet Loess Upland Located on glacial till parent material; soil surface color is 3/2 Munsell or darker.
F111XD008IN	Till Depression Flatwood Located on loess parent material; soils are taxonomically mollisols.

Table 1. Dominant plant species

Tree	(1) <i>Quercus palustris</i> (2) <i>Quercus bicolor</i>
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This ecosite is found in till plain in MLRA 111D: Indiana and Ohio Till Plain, Western Part

list of unique landform positions: Feetslope, Summit, Toeslope

list of unique landforms: depression, flat, glacial drainage channel, glacial lake (relict), ground moraine, lake plain, outwash plain, outwash terrace, pothole, stream terrace, till plain

Soils in this site have one or more characteristics that are severely limiting to the planting and growth of trees and shrubs. Soil depth is less than 20 inches; available water capacity is less than 3 inches; depth to a water table during the growing season is less than 0.5 feet; pH is less than 4.0 or greater than 8.5, sodium adsorption rate is greater than 25; CaCO₃ percentages equal or exceed 40%; flooding duration is very long. This group also includes urban land and water.

Table 2. Representative physiographic features

Landforms	(1) Lake plain (2) Till plain (3) Flat
Flooding duration	Extremely brief (0.1 to 4 hours) to brief (2 to 7 days)
Flooding frequency	None to occasional
Ponding duration	Brief (2 to 7 days) to very long (more than 30 days)
Ponding frequency	None to frequent
Elevation	91–351 m
Slope	0–2%
Ponding depth	0–76 cm
Water table depth	0–30 cm
Aspect	SE, W

Climatic features

The average annual precipitation in this area is 36 to 43 inches (915 to 1,090 millimeters). Most of the rainfall occurs as convective thunderstorms during the growing season. About half or more of the precipitation occurs during the freeze-free period. Snowfall is common in winter.

The average annual temperature is 49 to 54 degrees F (10 to 12 degrees C). The freeze-free period averages about 200 days and ranges from 180 to 215 days.

Table 3. Representative climatic features

Frost-free period (characteristic range)	133-150 days
Freeze-free period (characteristic range)	174-182 days
Precipitation total (characteristic range)	991-1,092 mm
Frost-free period (actual range)	132-158 days
Freeze-free period (actual range)	171-187 days
Precipitation total (actual range)	991-1,118 mm
Frost-free period (average)	144 days
Freeze-free period (average)	178 days
Precipitation total (average)	1,041 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) FRANKFORT DISPOSAL PLT [USC00123082], Frankfort, IN
- (2) YOUNG AMERICA [USC00129905], Kokomo, IN
- (3) FRANKLIN [USC00332928], Franklin, OH
- (4) TERRE HAUTE INDIANA ST [USC00128723], Terre Haute, IN
- (5) BOSWELL 4WNW [USC00120858], Fowler, IN
- (6) WABASH [USC00129138], Wabash, IN
- (7) XENIA 6 SSE [USC00339361], Xenia, OH
- (8) FAIRFIELD [USC00332651], Hamilton, OH

Influencing water features

This site is a depressional wetland that receives water via precipitation and runoff from surrounding landscapes. Flooding occurs occasionally with a brief (2 to 7 day) duration. The soils are mostly very poorly to poorly drained. Drainage and permeability (very slow to rapid) along with landscape position lead much of the site being frequently ponded, most often in the spring and occasionally in the fall. Duration of ponding varies from brief (2 to 7 days) to very long (> 30 days). Ponding depth has an average maximum of 30 inches.

Wetland description

The hydrogeographic model classification of this site is DEPRESSIONAL: Lake Plain, Ponded, Swale; forested. This site has a Cowardin Classification of PFO6An; it is a deciduous forested palustrine system that is temporarily ponded on mineral soil.

Soil features

The soil series associated with this site are: Zipp, Patton, Montgomery, Milford, Belleville. They are very deep, very poorly drained to poorly drained, and very slow to rapid permeable soils, with moderately acidic to moderately alkaline soil reaction, that formed in Glaciofluvial deposits, Glaciolacustrine deposits, and Lacustrine deposits

Table 4. Representative soil features

Parent material	(1) Glaciolacustrine deposits (2) Glaciofluvial deposits (3) Lacustrine deposits
Surface texture	(1) Loamy sand (2) Silt loam (3) Silty clay loam
Family particle size	(1) Loamy
Drainage class	Very poorly drained to poorly drained
Permeability class	Very slow to rapid
Soil depth	76 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	10.92–20.57 cm
Calcium carbonate equivalent (Depth not specified)	0–20%
Electrical conductivity (Depth not specified)	0 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0
Soil reaction (1:1 water) (Depth not specified)	6.5–7.9
Subsurface fragment volume <=3" (Depth not specified)	2%
Subsurface fragment volume >3" (Depth not specified)	2%

Ecological dynamics

The historic plant community of this ecological site is a flatwood forest. The dominant species in the canopy are pin oak and swamp white oak with silver maple and green ash being common as well. The soils of this site are very poorly or poorly drained and characterized by seasonal ponding in the spring and much drier in the summer. This seasonal change in water presence limited seed germination and diversity of shrubs and ground layer species.

State and transition model

Wet Lacustrine Forest, F111DY013IN



Wet Lacustrine Forest, F111DY013IN

Diagram Legend

T1A	Selective tree harvest
T1B	Remove woody species, drainage, site preparation, planting, management
T2A	Timber stand improvement and tree planting
T2B	Remove woody species, drainage, site preparation, planting, management
T3A	Drainage removal, planting, TSI management
3.1A	Pasture/forage planting and maintenance
3.2A	Tillage/no-till planting and management of row crops.

State 1 Flatwood Forest

This is the diagnostic plant community of the site. The dominant species in the canopy are pin oak and swamp white oak with green ash and silver maple being common as well. The seasonal change in water presence limited seed germination and diversity of shrubs and ground layer species.

Community 1.1 pin oak/swamp white oak

This phase is characterized by a closed to partially open canopy dominated by pin oak and swamp white oak.

State 2

High Graded State

This phase is characterized by the removal of the more marketable tree species, primarily oak species. The resulting tree species, maples, ash, black gum, become the dominant species in the canopy.

Community 2.1

maple/ash/black gum

This phase is characterized by the removal of the oaks and the maple/ash/black gum dominate the canopy.

State 3

Agriculture State

This state is characterized by the conversion of the site to agricultural use. Most common practice is a corn and soybean rotation of various types. A small portion of the historic acres are used for forage and pasture.

Community 3.1

corn/soybeans

This phase is characterized by row crop agriculture of small grains, primarily corn and soybeans.

Community 3.2

cool season pasture/forage

This phase is characterized by forage or grazing agriculture. Different mixes of, generally, cool season grasses and forbs, largely clovers, are grown.

Pathway P3.1

Community 3.1 to 3.2

Establishment of forage / pasture species.

Pathway P3.2

Community 3.2 to 3.1

Establishment of row crops

Transition T 1-2

State 1 to 2

Selective tree harvest of the more marketable species, primarily oaks.

Transition T 1-3

State 1 to 3

Removal off trees and other wood species. Install drainage system (if warranted), prepare the site for planting the agricultural crop, and regular agricultural practices.

Restoration pathway R 2-1

State 2 to 1

Timber stand improvement practices and planting (if warranted) of desired species.

Transition T 2-3 State 2 to 3

Removal of trees and other wood species. Install drainage system (if warranted), prepare the site for planting the agricultural crop, and regular agricultural practices.

Restoration pathway R 3-1 State 3 to 1

Removal of drainage system (if warranted), site preparation, and tree planting.

Additional community tables

Inventory data references

Site concept developed through expert opinion, review of the literature, and field reconnaissance.

Other references

Braun, E. Lucy. 2001. Deciduous forests of eastern North America. Caldwell, N.J.: Blackburn Press.

Federal Geographic Data Committee. 2013. Classification of wetlands and deepwater habitats of the United States. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, DC.

Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. Indiana Academy of Science , 94, 245-269.

NatureServe. (2011). An online encyclopedia of life [web application]. NatureServe, Arlington, VA, USA [Online: www.natureserve.org/explorer].

Jackson, Marion T. 1997. The Natural heritage of Indiana. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

Johnson, Paul S., Stephen R. Shifley, and Robert Rogers. 2002. The ecology and silviculture of oaks. Wallingford, Oxon: CABI

USDA. (2007). Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

USDA. (2006). Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U. S. Department of Agriculture, Natural Resources Conservation Service. U. S. Department of Agriculture Handbook 296.

USDA-NRCS. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service. Technical Note No. 190–8–76. Washington D.C.

USGS. (2010). LANDFIRE Biophysical Settings. Retrieved from <http://www.landfire.gov>

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

Approval

Chris Tecklenburg, 5/28/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Tyler Staggs
Contact for lead author	
Date	05/03/2024
Approved by	Chris Tecklenburg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial**

distribution on infiltration and runoff:

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-